

# A contribution to the understanding of ocular and cerebrospinal fluid dynamics in astronauts during long-lasting spaceflight

---

Orešković, Darko; Radoš, Milan; Klarica, Marijan

Source / Izvornik: **Croatian Medical Journal, 2021, 62, 420 - 421**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.3325/cmj.2021.62.420>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:829000>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International/Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-03-05**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine  
Digital Repository](#)



## A contribution to the understanding of ocular and cerebrospinal fluid dynamics in astronauts during long-lasting spaceflight

Darko Orešković<sup>1</sup>, Milan Radoš<sup>2</sup>, Marijan Klarica<sup>2</sup>

<sup>1</sup>Department of Molecular Biology, Ruđer Bošković Institute, Zagreb, Croatia  
[doresk@irb.hr](mailto:doresk@irb.hr)

<sup>2</sup>Department of Pharmacology and Croatian Institute for Brain Research, Zagreb University School of Medicine, Zagreb, Croatia

Zhang and Hargens (1) reviewed the knowledge on visual impairment intracranial pressure (VIIP) syndrome in astronauts, a condition that arises as a consequence of long-lasting spaceflight. However, the etiology and pathophysiology of VIIP syndrome remain unclear and only hypothetical. The main reason is the absence of data regarding astronauts' intracranial (ICP) and intraocular pressure (IOP) during long-lasting spaceflights. It is believed that a crucial role in the pathogenesis of VIIP syndrome is played by chronic ICP/IOP mismatch, which is a consequence of chronically elevated ICP and which results in translaminal pressure gradient. A direct transmission of elevated ICP from the intracranial to the intraocular compartment through the perioptic subarachnoid space should lead to optic nerve sheath distension and disk edema. Therefore, elevated pressure gradient across the lamina cribrosa caused by chronic, gently elevated ICP should finally cause posterior-globe flattening, disk edema, choroid folds, and a hyperoptic shift (1).

Zhang et al (1) explain ICP regulation by the classic hypothesis of cerebrospinal fluid (CSF) physiology, which describes CSF circulation as a third circulation (1-5). However, we believe that the presented hypothesis of VIIP syndrome could be partially elucidated by our new approach to CSF physiology. Based on our CSF physiology hypothesis, another hypothesis of ICP and IOP regulation was derived, in which the total pressure is expressed by the equation (1):

$$P_o = P_v + FxR$$

where  $P_o$  is ICP or IOP,  $P_v$  is the superior sagittal sinus pressure or episcleral venous pressure; while  $F$  and  $R$

are the CSF or aqueous humor system formation rate and the outflow resistance (4-6). This equation shows that both pressures directly depend on the CSF secretion, circulation, and absorption, and that the pressures will be higher if there is greater resistance to the CSF circulation and absorption. It should be emphasized that there are no principle differences regarding IOP and ICP regulation explanation.

We have recently abandoned the classic hypothesis of CSF physiology and formed a new one (7-9). The hypothesis of ICP regulation has been derived from the new CSF hypothesis and from experimental results on animals and a phantom, together with theoretical analyses of equations describing fluid mechanics. Hence, the ICP regulation appears not to be a product of CSF formation, circulation, and absorption relationship, as is generally believed (1,4-6), but depends on the laws of fluid mechanics and on the anatomical characteristics inside the cranial and spinal CSF space (10-13). ICP is a hydrostatic pressure, which directly depends on the gravitational force. In the human brain ventricles in the sitting position ICP is about -10 cm H<sub>2</sub>O, while in the lumbar space it is about +60 cm H<sub>2</sub>O (10-13). According to the laws of fluid mechanics, if the gravitation is zero or almost zero, the pressure gradient should disappear (from -10 to +60), and negative ICP values should become positive, which should permanently increase the ventricular ICP (10-13).

Furthermore, our experiments on cats (14) have shown that significant ICP changes caused by changes in body position did not significantly change IOP. Thus, while during a body position shift from horizontal to vertical, the ICP

in ventricles decreased significantly, for 21.5 cm H<sub>2</sub>O (from 17.4 to -4.1 cm H<sub>2</sub>O), the IOP decreased insignificantly, for 4.2 cm H<sub>2</sub>O (from 18.5 to 14.3 cm H<sub>2</sub>O). Our results clearly show that in the vertical body position there is a (physiological) gradient between ICP and IOP, predominantly in the eyes.

This means that regulation mechanisms of ICP and IOP are fundamentally different, rather than the same as was traditionally believed (1,4-6). Therefore, micro-gravity, which directly affects ICP, should not have such an effect on IOP. A different way of pressure regulation is likely what causes an altered pressure gradient. Accordingly, a significantly higher ICP should reduce the (physiological) gradient between ICP and IOP.

To conclude, according to the new hypothesis of ICP regulation, an ICP increase is expected in microgravity, and represents an additional confirmation of the new CSF physiology hypothesis. Additionally, due to different regulatory mechanisms of IOP and ICP, a non-physiological gradient between these pressures should exist. However, the etiology of VIIP syndrome is expected to remain only hypothetical as long as human ICP and IOP are not measured in long-lasting microgravity conditions.

**Acknowledgments** This work has been supported by the Croatian Science Foundation and Ministry of Science and Education of the Republic of Croatia (Projects: 1. Pathophysiology of cerebrospinal fluid and intracranial pressure. No. 108-1080231-0023; and 2. Serotonergic modulation of obesity: cross-talk between regulatory molecules and pathways. No. IP-2014-09-7827). Research was co-financed by the Scientific Centre of Excellence for Basic, Clinical and Translational Neuroscience (project: Experimental and clinical research of hypoxic-ischemic damage in perinatal and adult brain; GA KK01.1.1.01.0007 funded by the European Union).

## References

- Zhang L, Hargens AR. Spaceflight-induced intracranial hypertension and visual impairment: pathophysiology and countermeasures. *Physiol Rev*. 2018;98:59-87. [Medline:29167331](#) [doi:10.1152/physrev.00017.2016](#)
- Cushing H. The third circulation and its channels (Cameron Lecture). *Lancet*. 1925;2:851-7.
- Dandy WE. Experimental hydrocephalus. *Ann Surg*. 1919;70:129-42. [Medline:17864139](#) [doi:10.1097/0000658-191908000-00001](#)
- Davson H, Domer FR, Hollingsworth JR. The mechanism of drainage of the cerebrospinal fluid. *Brain*. 1973;96:329-36. [Medline:4197586](#) [doi:10.1093/brain/96.2.329](#)
- Davson H, Welch K, Segal MB. *Physiology and pathophysiology of the cerebrospinal fluid*. Edinburgh: Churchill-Livingstone; 1987.
- Goldman H. Out-flow pressure, minute volume and resistance of the anterior chamber flow in man. *Doc Ophthalmol*. 1951;5-6:278-356. [Medline:14896880](#)
- Bulat M, Klarica M. Recent insights into a new hydrodynamics of the cerebrospinal fluid. *Brain Res Brain Res Rev*. 2011;65:99-112. [Medline:20817024](#) [doi:10.1016/j.brainresrev.2010.08.002](#)
- Orešković D, Klarica M. Development of hydrocephalus and classical hypothesis of cerebrospinal fluid hydrodynamics: facts and illusions. *Prog Neurobiol*. 2011;94:238-58. [Medline:21641963](#) [doi:10.1016/j.pneurobio.2011.05.005](#)
- Orešković D, Klarica M. The formation of cerebrospinal fluid: nearly a hundred years of interpretations and misinterpretations. *Brain Res Brain Res Rev*. 2010;64:241-62. [Medline:20435061](#) [doi:10.1016/j.brainresrev.2010.04.006](#)
- Klarica M, Radoš M, Erceg G, Petošić A, Jurjević I, Orešković D. The influence of body position on cerebrospinal fluid pressure gradient and movement in cats with normal and impaired craniospinal communication. *PLoS One*. 2014;9:e95229. [Medline:24748150](#) [doi:10.1371/journal.pone.0095229](#)
- Klarica M, Radoš M, Draganić P, Erceg G, Orešković D, Maraković J, et al. Effect of head position on cerebrospinal fluid pressure in cats: comparison with artificial model. *Croat Med J*. 2006;47:233-8. [Medline:16625687](#)
- Orešković D, Radoš M, Klarica M. Role of choroid plexus in cerebrospinal fluid hydrodynamics. *Neurosci*. 2017;354:69-87. [Medline:28456718](#) [doi:10.1016/j.neuroscience.2017.04.025](#)
- Orešković D, Radoš M, Klarica M. The recent state of a hundred years old classic hypothesis of the physiology of cerebrospinal fluid. *Croat Med J*. 2017;58:381-3. [Medline:29308829](#) [doi:10.3325/cmj.2017.58.381](#)
- Kuzman T, Jurjević I, Mandac I, Radoš M, Orešković D, Jednačak H, et al. The effect of body position on intraocular and CSF pressures in the lateral ventricle, cortical and lumbar subarachnoid spaces in cats. *Acta Neuroch*. 2012;114:357-61. [Medline:22327723](#) [doi:10.1007/978-3-7091-0956-4\\_69](#)