

# Possible mechanisms mediating complete regrowth of hair following scalp tattooing in alopecia universalis

---

Homolak, Jan

Source / Izvornik: **JAAD Case Reports, 2023, 34, 10 - 11**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1016/j.jdcr.2023.01.031>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:540770>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International/Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-03-14**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



### Possible mechanisms mediating complete regrowth of hair following scalp tattooing in alopecia universalis



*To the Editor:* Ramnot et al<sup>1</sup> recently described a fascinating case of the complete regrowth of hair following scalp tattooing in a patient with alopecia universalis unresponsive to previous treatment (local and systemic glucocorticoids, squaric acid dibutyl ester immunotherapy) despite high compliance. The causality between scalp tattooing and alopecia regression cannot be determined with certainty; however, the timeline of events, location, and size of the observed effect suggest that the regrowth of hair might have been triggered by tattoo ink and/or the process of tattooing via unknown mechanisms. This letter aims to propose possible mechanisms by which the process of tattooing and the presence of tattoo ink might promote hair growth in alopecia.

In the process of tattooing, insoluble pigments are deposited into the dermal skin layer with a specialized needle where they are either engulfed by dermal macrophages<sup>2</sup> or transported to the draining lymph nodes.<sup>3,4</sup> Tattooing is inherently traumatic as the skin is repeatedly punctured with multiple closely spaced needles which disrupt the epidermal basement membrane and damage epidermal and dermal cells that consequently undergo necrosis.<sup>4</sup> Traumatic injury of tattooed skin has the potential to activate hair follicles as physiological mechanisms regulating wound healing and hair growth are highly intertwined.<sup>5</sup> The aforementioned is considered to be the underlying principle of microneedling that has shown some promising effects on the skin,<sup>6</sup> as well as on hair loss.<sup>7-10</sup> Accumulating evidence show that hair follicles are involved in wound healing, and that wounding promotes the transition of follicles to the anagen phase, possibly by activating Wnt/ $\beta$ -catenin signaling and promoting follicle (re)vascularization by stimulating vascular endothelial growth factor.<sup>11-13</sup>

Tattooing-induced hair growth may also be mediated by the effects of bioactive components of tattoo inks (organic or inorganic pigments, binders, solvents, additives, impurities, and microorganisms<sup>14-16</sup>) on hair follicles and/or alopecia-associated pathophysiological processes (eg, autoimmunity,<sup>17-19</sup>

oxidative stress<sup>20,21</sup>). The direct effects of tattoo ink components on hair follicles are unexplored; however, there are several reports of henna (surface) tattoo-induced hypertrichosis.<sup>22</sup> Furthermore, some tattoo inks demonstrate immunomodulatory and antioxidant properties. Functional consequences of sequestration of tattoo ink pigments in dermal macrophages are still unexplored,<sup>2</sup> however, Devcic et al recently reported that tattoo ink pigment-exposed macrophages secrete less interleukin 6, tumor necrosis factor, and chemoattractant cytokines (monocyte chemoattractant protein-1, macrophage inflammatory protein-1 $\alpha$ ) after stimulation with lipopolysaccharide, suggesting that tattoo pigments might act as immunosuppressants.<sup>23</sup> Polycyclic aromatic hydrocarbons, often found in black tattoo inks,<sup>24</sup> have been proposed as mediators of the immunomodulatory effects of crude coal tar used in the treatment of psoriasis.<sup>25</sup> An N-of-1 study reported antioxidant properties of a blue tattoo in vivo<sup>26</sup> and an in vitro study reported that copper phthalocyanine-containing blue tattoo ink demonstrates catalase and superoxide dismutase mimetic properties.<sup>15</sup> The aforementioned results suggest that some tattoo ink constituents might theoretically promote hair growth by inhibiting etiopathogenetic factors of alopecia (inflammation and oxidative stress).

To conclude, the reported tattoo-induced regrowth of hair growth might be explained by stimulation of the transition of hair follicles to the anagen phase by the process of tattooing (by stimulating wound healing mechanisms) and/or immunomodulatory and antioxidant properties of some tattoo ink constituents.

Jan Homolak, MD<sup>a,b</sup>

*From the Department of Pharmacology, University of Zagreb School of Medicine, Zagreb, Croatia<sup>a</sup>; and Croatian Institute for Brain Research, University of Zagreb School of Medicine, Zagreb, Croatia.<sup>b</sup>*

*Funding sources:* None.

*IRB approval status:* Not applicable.

*Key words:* alopecia; hair; hair growth; tattoo; tattoo ink.

*Correspondence to:* Jan Homolak, MD, Department of Pharmacology, University of Zagreb School of Medicine, Šalata 11, 10 000, Zagreb, Croatia

*E-mail's:* [homolakjan@gmail.com](mailto:homolakjan@gmail.com), [jan.homolak@mef.br](mailto:jan.homolak@mef.br)

### Conflicts of interest

None disclosed.

### REFERENCES

1. Ramnot A, Resnik S, Resnik B. Complete regrowth of hair following scalp tattooing in a patient with alopecia universalis. *JAAD Case Rep.* 2023;31:102-104. <https://doi.org/10.1016/j.jdc.2022.10.027>
2. Baranska A, Shawket A, Jouve M, et al. Unveiling skin macrophage dynamics explains both tattoo persistence and strenuous removal. *J Exp Med.* 2018;215(4):1115-1133. <https://doi.org/10.1084/jem.20171608>
3. Schreiber I, Hesse B, Seim C, et al. Synchrotron-based  $\nu$ -XRF mapping and  $\mu$ -FTIR microscopy enable to look into the fate and effects of tattoo pigments in human skin. *Sci Rep.* 2017; 7(1):11395. <https://doi.org/10.1038/s41598-017-11721-z>
4. Islam PS, Chang C, Selmi C, et al. Medical complications of tattoos: a comprehensive review. *Clin Rev Allergy Immunol.* 2016;50(2):273-286. <https://doi.org/10.1007/s12016-016-8532-0>
5. Ansell DM, Kloeppe JE, Thomason HA, Paus R, Hardman MJ. Exploring the "hair growth-wound healing connection": anagen phase promotes wound re-epithelialization. *J Invest Dermatol.* 2011;131(2):518-528. <https://doi.org/10.1038/jid.2010.291>
6. Iriarte C, Awosika O, Rengifo-Pardo M, Ehrlich A. Review of applications of microneedling in dermatology. *Clin Cosmet Investig Dermatol.* 2017;10:289-298. <https://doi.org/10.2147/CCID.S142450>
7. Dhurat R, Sukesh M, Avhad G, Dandale A, Pal A, Pund P. A randomized evaluator blinded study of effect of microneedling in androgenetic alopecia: a pilot study. *Int J Trichology.* 2013;5(1):6-11. <https://doi.org/10.4103/0974-7753.114700>
8. Fertig RM, Gamret AC, Cervantes J, Tosti A. Microneedling for the treatment of hair loss? *J Eur Acad Dermatol Venereol.* 2018; 32(4):564-569. <https://doi.org/10.1111/jdv.14722>
9. English RS, Ruiz S, DoAmaral P. Microneedling and its use in hair loss disorders: a systematic review. *Dermatol Ther (Heidelb).* 2022;12(1):41-60. <https://doi.org/10.1007/s13555-021-00653-2>
10. Gupta AK, Quinlan EM, Venkataraman M, Bamimore MA. Microneedling for hair loss. *J Cosmet Dermatol.* 2022;21(1): 108-117. <https://doi.org/10.1111/jocd.14525>
11. Yano K, Brown LF, Detmar M. Control of hair growth and follicle size by VEGF-mediated angiogenesis. *J Clin Invest.* 2001; 107(4):409-417.
12. Myung PS, Takeo M, Ito M, Atit RP. Epithelial Wnt ligand secretion is required for adult hair follicle growth and regeneration. *J Invest Dermatol.* 2013;133(1):31-41. <https://doi.org/10.1038/jid.2012.230>
13. Kim YS, Jeong KH, Kim JE, Woo YJ, Kim BJ, Kang H. Repeated microneedle stimulation induces enhanced hair growth in a murine model. *Ann Dermatol.* 2016;28(5):586-592. <https://doi.org/10.5021/ad.2016.28.5.586>
14. Giullbudagian M, Schreiber I, Singh AV, Laux P, Luch A. Safety of tattoos and permanent make-up: a regulatory view. *Arch Toxicol.* 2020;94(2):357-369. <https://doi.org/10.1007/s00204-020-02655-z>
15. Homolak J. In vitro analysis of catalase and superoxide dismutase mimetic properties of blue tattoo ink. *Free Radic Res.* 2022;56(5-6):343-357. <https://doi.org/10.1080/10715762.2022.2102976>
16. Nho SW, Kim M, Kweon O, et al. Microbial contamination of tattoo and permanent makeup inks marketed in the US: a follow-up study. *Lett Appl Microbiol.* 2020;71(4):351-358. <https://doi.org/10.1111/lam.13353>
17. Ito T. Recent advances in the pathogenesis of autoimmune hair loss disease alopecia areata. *Clin Dev Immunol.* 2013;2013: 348546. <https://doi.org/10.1155/2013/348546>
18. Ito T, Kageyama R, Nakazawa S, Honda T. Understanding the significance of cytokines and chemokines in the pathogenesis of alopecia areata. *Exp Dermatol.* 2020;29(8):726-732. <https://doi.org/10.1111/exd.14129>
19. Bain KA, McDonald E, Moffat F, et al. Alopecia areata is characterized by dysregulation in systemic type 17 and type 2 cytokines, which may contribute to disease-associated psychological morbidity. *Br J Dermatol.* 2020;182(1):130-137. <https://doi.org/10.1111/bjd.18008>
20. Acharya P, Mathur MC. Oxidative stress in alopecia areata: a systematic review and meta-analysis. *Int J Dermatol.* 2020; 59(4):434-440. <https://doi.org/10.1111/jid.14753>
21. Sachdeva S, Khurana A, Goyal P, Sardana K. Does oxidative stress correlate with disease activity and severity in alopecia areata? An analytical study. *J Cosmet Dermatol.* 2022;21(4): 1629-1634. <https://doi.org/10.1111/jocd.14253>
22. El Habr C, Mégarbané H. Temporary henna tattoos and hypertrichosis: a case report and review of the literature. *J Dermatol Case Rep.* 2015;9(2):36-38. <https://doi.org/10.3315/jdc.2015.1204>
23. Devic J, Dussol M, Collin-Faure V, et al. Immediate and sustained effects of cobalt and zinc-containing pigments on macrophages. *Front Immunol.* 2022;13:865239. <https://doi.org/10.3389/fimmu.2022.865239>
24. Lehner K, Santarelli F, Vasold R, et al. Black tattoos entail substantial uptake of genotoxic polycyclic aromatic hydrocarbons (PAH) in human skin and regional lymph nodes. *PLoS One.* 2014;9(3):e92787. <https://doi.org/10.1371/journal.pone.0092787>
25. Sekhon S, Jeon C, Nakamura M, et al. Review of the mechanism of action of coal tar in psoriasis. *J Dermatolog Treat.* 2018;29(3):230-232. <https://doi.org/10.1080/09546634.2017.1369494>
26. Homolak J. The effect of a color tattoo on the local skin redox regulatory network: an N-of-1 study. *Free Radic Res.* 2021;55(3): 221-229. <https://doi.org/10.1080/10715762.2021.1912340>

<https://doi.org/10.1016/j.jdc.2023.01.031>