

# Enigma of cerebrospinal fluid dynamics

---

**Klarica, Marijan; Orešković, Darko**

*Source / Izvornik:* **Croatian Medical Journal, 2014, 55, 287 - 298**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

<https://doi.org/10.3325/cmj.2014.55.287>

*Permanent link / Trajna poveznica:* <https://um.nsk.hr/um:nbn:hr:105:345991>

*Rights / Prava:* [In copyright](#)/[Zaštićeno autorskim pravom.](#)

*Download date / Datum preuzimanja:* **2025-03-31**



*Repository / Repozitorij:*

[Dr Med - University of Zagreb School of Medicine  
Digital Repository](#)



## Enigma of cerebrospinal fluid dynamics

Marijan Klarica<sup>1</sup>, Darko Orešković<sup>2</sup>

<sup>1</sup>Department of Pharmacology and Croatian Institute for Brain Research, University of Zagreb School of Medicine, Zagreb, Croatia

[marijan.klarica@mef.hr](mailto:marijan.klarica@mef.hr)

<sup>2</sup>Department of Molecular Biology, Ruđer Bošković Institute, Zagreb, Croatia

### TWO HYPOTHESES ON CSF PHYSIOLOGY

Cerebrospinal fluid (CSF) is a major part of the central nervous system (CNS) extracellular fluid, and fine regulation of its composition is vital to the brain's health. Although CSF dynamics has been studied for an entire century, many of its aspects are still insufficiently understood. Today there are two hypotheses (1,2) on CSF physiology: a) traditional hypothesis and b) microcirculatory/microvessel hypothesis.

According to the traditional hypothesis, CSF is formed inside the brain ventricles, mostly by secretion from the choroid plexuses, and it circulates along the ventricles and subarachnoid space to be absorbed across the arachnoid villi into the dural venous sinuses, and/or viacrinal and spinal nerves paraneural sheaths into the lymphatic system. Since substance exchange occurs between the CNS extracellular interstitial fluid (ISF) and CSF, it is assumed that CSF serves as a sink for the removal of various metabolites out of the CNS by its unidirectional pulsatile flow and absorption (3-5). This traditional hypothesis, with minor modifications, represents a common point of reference in scientific papers, review articles, and textbooks on the issue (6,7). Additionally, this hypothesis has been used to explain the yet clinically unsolved pathological states such as increase in intracranial pressure and hydrocephalus.

The emerging concept of CSF physiology described as microcirculatory/microvessel hypothesis suggests that CNS microvessels are instrumental in fluid filtration and reabsorption inside the brain and spinal cord parenchyma, as well as inside the CSF system (2,8-10). It seems that the osmotic pressure change inside the CNS capillary network is crucial in regulation of ISF and CSF volumes, which are

continuously mixed by to-and-fro fluid pulsations (9). Thus, distribution of water, which constitutes the bulk of ISF and CSF, is very limited due to its rapid turnover across the microvascular walls. It is important to emphasize that microcirculatory/microvessel hypothesis interconnects the physiology of all craniospinal fluids (plasma, intracellular, extracellular, and cerebrospinal fluid). This new concept of CSF and other fluids dynamics opens many possibilities in the investigation of severe clinical problems such as normal pressure, arrested or acute hypertensive hydrocephalus, intracranial hypertension, and focal and generalized brain edema.

This issue of the *Croatian Medical Journal* (CMJ) is dedicated to the CSF dynamics, and we hope that it will provide the readers with new data and views that can help them in their research projects. Due to the significance of the new CSF dynamics concept, this issue's editorial is supplemented by a reprint of the article by Bulat and Klarica published in the *Periodicum Biologorum* in 2005, which first presented the new hypothesis on fluids physiology (11) ([Supplementary material](#)). This issue also offers several comprehensive reviews on different aspects of CSF functioning. Gato et al describe embryonic CSF before the development of the choroid plexuses and show different novel concepts of embryonic CSF (eCSF) functioning, while Bueno et al in their extensive and detailed review of a very early protective barrier (embryonic blood-CSF barrier) explain the control of internal milieu (components of eCSF) in a developing brain. Orešković and Klarica describe methodological errors of traditional and most widely accepted perfusion method for measuring CSF formation and absorption. The article by Nakada discusses a new concept of CSF physiology, describing the importance of Virchow-

Robin space and aquaporin 4 in the functioning of CNS, while that by Yamada describes novel aspects of CSF physiological and pathophysiological movement visualized by a new non-invasive MRI technique (Time-SLIP). Babić et al discuss the current status of a great variety of CSF biomarkers for the use in Alzheimer disease diagnostics and Krishnamurthy et al summarize the data about pathophysiological mechanisms of hydrocephalus development induced by an application of hyperosmolar solution into the CSF space, and hypothesize that impaired efflux at the blood-brain barrier will result in an increased concentration of different substances inside the brain interstitial and ventricular fluid, leading to hydrocephalus development.

This issue also presents reports on several cases that can in no way be explained by the generally accepted traditional hypothesis. Such is the case of idiopathic CSF "hypersecretion" (Trevisi et al) in a 6-month-old infant, which cannot be controlled using standard operative drainage techniques (the article also gives an overview of the available data regarding CSF formation). There is also the case of the oldest living patient with hydranencephaly, which raises some questions regarding the CSF turnover and homeostasis in a person with no brain parenchyma inside the supratentorial space (Radoš et al). Also, this issue includes a case of a severe aqueductal stenosis lasting for 5 years without any detectable CSF movements, which is not accompanied

with hydrocephalus development (Radoš et al). In addition, Bechter and Shmitz describe time dynamics of contrast distribution from lumbar subarachnoid space into the psoas muscle tissue in one patient, and discuss the importance of their findings for pain research.

#### MARIN BULAT'S CONTRIBUTION TO THE PHYSIOLOGY AND PATHOPHYSIOLOGY OF THE CEREBROSPINAL FLUID

Due to the importance of his research for disputing the traditional CSF physiology hypothesis, we dedicate a part of this editorial to the work of Prof. emeritus Marin Bulat (1936-2012), who started the CSF physiology research in Croatia (Figure 1, Figure 2).

Marin Bulat began his scientific career at the Ruđer Bošković Institute under the mentorship of Prof. Zlatko Supek. In his master's (1964) and PhD thesis (1966), which determined the course of his scientific work, he explored the role of serotonin and its metabolites inside the CSF system and the brain. His early papers describe the serotonin passage from CSF into the brain by means of diffusion process (without any restrictions) depending on the concentration gradient (12,13). Serotonin is very quickly metabolized after its passage from CSF into the brain tissue and its metabolite 5-HIAA quickly disappears from the site of its formation. He showed that serotonin did not distribute inside the CSF



FIGURE 1. Professor emeritus Marin Bulat (1936-2012)




FIGURE 2. A blackboard in Prof. Bulat's study after many hours of discussion about the dynamics of cerebrospinal fluid and other intracranial fluids.

and that its lumbar CSF concentration merely reflected the changes inside the surrounding spinal cord tissue (14,15). Namely, the 5-HIAA concentration changes after its application into the cisterna magna did not influence its concentration inside the lumbar subarachnoid space. However, based on the classic hypothesis of CSF physiology, increased concentration inside the lumbar subarachnoid space would be reasonably expected. As he noticed that an increase in metabolic transformation inside the spinal cord tissue led to an increase in the concentration inside the surrounding CSF, he concluded that the changes of 5-HIAA concentration inside the lumbar CSF reflected only the local metabolism inside the spinal cord tissue (14,16).

On the basis of these observations, an idea developed that substances were not distributed throughout the CSF according to the classic hypothesis of CSF physiology. When molecules with different molecular weight were monitored, such as radioactive water, organic acids (5-HIAA,  $^3\text{H}$ -benzylpenicillin, phenolsulphonphthalein) or inulin inside the CSF system and CNS, it was observed that they were distributed in all directions and that the distribution intensity depended on the rate of their elimination into the CNS capillaries (9,17,18). According to the generally accepted hypothesis, the CSF is secreted inside the brain ventricles and flows unidirectionally along the subarachnoid spaces to be absorbed into the dural venous sinuses. However, a small molecule like water, which constitutes 99% of CSF bulk, does not flow unidirectionally along the CSF spaces since it is rapidly absorbed into the adjacent microvessels (19).

The distribution of substances with long residence time (inulin, proteins, etc) caused by to-and fro pulsations will always, after their application into the lateral ventricles (LV), create an illusion of a unidirectional bulk CSF circulation (from LV to cisterna magna, cortical and spinal subarachnoid space). On the contrary, after application of these substances into other parts of the CSF system, they are distributed in all directions (as well as into the LV, which is contradictory to the classical hypothesis) (9). Besides this, it can be observed that the arrival of inulin into the area of lumbar CSF is much faster than into the cortical subarachnoid space. All of this implies that the CSF moves much differently than what was previously conceived. After a series of experiments on the influence of CSF and blood hydrostatic and osmotic pressure on the CSF volume and pressure had been performed, a new hypothesis was suggested, according to which interstitial fluid and CSF make a single functional unit, while CNS microvessels are crucial for the fluid absorption and filtration (9).

In his fight against the conventional notions, Prof. Bulat found inspiration and strenght in the thought of Claude Bernard, one of the most distinguished physiologists and medical scientists of the nineteenth century: "When we meet a fact which contradicts a prevailing theory, we must accept the fact and abandon the theory, even when the theory is supported by great names and generally accepted." He never gave up, and he managed to show his students and colleagues the way to persevere in science. He used to say that science is like a candle that burns low, and that this flame should be preserved and carried on with extreme attention, scientific integrity, and hard persistent work. He passed on this difficult task to all of us who are now, together with our foreign colleagues, trying to save that flame of science and to pass it on to new generations.

Instead of conclusion let's thank Prof. emeritus Marin Bulat with another famous sentence by Claude Bernard: "A man of science rises ever, in seeking truth; and if he never finds it in its wholeness, he discovers nevertheless very significant fragments; and these fragments of universal truth are precisely what constitutes science."

## References

- 1 Bateman GA, Brown KM. The measurement of CSF flow through the aqueduct in normal and hydrocephalic children: from where does it come, to where does it go. *Childs Nerv Syst.* 2012;28:55-63. [Medline:22038153](#) [doi:10.1007/s00381-011-1617-4](#)
- 2 Naidich TP, Castillo M, Cha S, Smirniotopoulos JG. *Imaging of the brain.* Philadelphia: Saunders Elsevier; 2013.
- 3 Davson H, Welch K, Segal MB. *The physiology and pathophysiology of the cerebrospinal fluid.* 1st edition. Edinburgh: Churchill Livingstone; 1987.
- 4 Fishman AR. *Cerebrospinal fluid in disease of the nervous system.* Philadelphia: W B Saunders Co; 1991.
- 5 Irani DN. *Cerebrospinal fluid in clinical practice.* Philadelphia: Saunders Elsevier; 2009.
- 6 Johanson CE, Duncan JA, Klinge PM, Brinker T, Stopa EG, Silveberg GD. Multiplicity of cerebrospinal fluid functions: new challenges in health and disease. *Cerebrospinal Fluid Res.* 2008;5:10. [Medline:18479516](#) [doi:10.1186/1743-8454-5-10](#)
- 7 Brinker T, Stopa E, Morrison J, Klinge P. A new look at cerebrospinal fluid circulation. *Fluids Barriers CNS.* 2014;11:10. [Medline:24817998](#) [doi:10.1186/2045-8118-11-10](#)
- 8 Orešković D, Klarica M. The formation of cerebrospinal fluid: nearly a hundred years of interpretations and misinterpretations. *Brain Res Rev.* 2010;64:241-62. [Medline:20435061](#) [doi:10.1016/j.brainresrev.2010.04.006](#)
- 9 Bulat M, Klarica M. Recent insights into a new hydrodynamics of the cerebrospinal fluid. *Brain Res Rev.* 2011;65:99-112.

- [Medline:20817024](#) [doi:10.1016/j.brainresrev.2010.08.002](#)
- 10 Orešković D, Klarica M. Development of hydrocephalus and classical hypothesis of cerebrospinal fluid hydrodynamics: facts and illusions. *Prog Neurobiol.* 2011;94:238-58. [Medline:21641963](#) [doi:10.1016/j.pneurobio.2011.05.005](#)
  - 11 Bulat M, Klarica M. Fluid filtration and reabsorption across microvascular walls: control by oncotic or osmotic pressure? *Period Biol.* 2005;107:147-52.
  - 12 Bulat M, Supek Z. Fate of intracisternally injected 5-hydroxytryptamine in the rat brain. *Nature.* 1966;211:637-8. [Medline:5968736](#) [doi:10.1038/211637b0](#)
  - 13 Bulat M, Supek Z. Mechanism of 5-hydroxytryptamine penetration through the cerebrospinal fluid-brain barrier. *Nature.* 1968;219:72-3. [Medline:5659622](#) [doi:10.1038/219072a0](#)
  - 14 Bulat M, Živković B. Origin of 5-hydroxyindoleacetic acid in the spinal fluid. *Science.* 1971;173:738-40. [Medline:5109594](#) [doi:10.1126/science.173.3998.738](#)
  - 15 Bulat M, Lacković Z, Jakupčević M, Damjanov I. 5-Hydroxyindoleacetic acid in the lumbar fluid – a specific indicator of spinal cord injury. *Science.* 1974;185:527-8. [Medline:4841583](#) [doi:10.1126/science.185.4150.527](#)
  - 16 Bulat M, Živković B. Exchange of 5-hydroxyindoleacetic acid between spinal cord and lumbar cerebrospinal fluid. *J Physiol.* 1978;275:191-7. [Medline:633105](#)
  - 17 Vladić A, Strikić N, Jurčić D, Zmajević M, Klarica M, Bulat M. Homeostatic role of the active transport in elimination of 3H-benzylpenicillin out of the cerebrospinal fluid system. *Life Sci.* 2000;67:2375-85. [Medline:11065184](#) [doi:10.1016/S0024-3205\(00\)00823-7](#)
  - 18 Zmajević M, Klarica M, Varda R, Kudelić N, Bulat M. Elimination of phenolsulfonphthalein from the cerebrospinal fluid via capillaries in the central nervous system in cats by active transport. *Neurosci Lett.* 2002;321:123-5. [Medline:11872271](#) [doi:10.1016/S0304-3940\(01\)02526-5](#)
  - 19 Bulat M, Lupret V, Orešković D, Klarica M. Transventricular and transpial absorption of cerebrospinal fluid into cerebral microvessels. *Coll Antropol.* 2008;32:43-50. [Medline:18405057](#)

## Fluid filtration and reabsorption across microvascular walls: control by oncotic or osmotic pressure? (secondary publication)

The article represents a secondary publication identical to previously published paper Bulat M, Klarica M. Fluid filtration and reabsorption across microvascular walls: control by oncotic or osmotic pressure? *Period Biol.* 2005;107:147-52. Published with premission from *Periodicum biologorum*.

**Aim.** Relationships between hydrostatic and oncotic (colloid osmotic) pressures in both capillaries and interstitium are used to explain fluid filtration and reabsorption across microvascular walls. These pressures are incorporated in the Starling oncotic hypothesis of capillaries which fails, however, to explain fluid homeostasis when hydrostatic capillary pressure is high (in feet during orthostasis) and low (in lungs), or when oncotic plasma pressure is significantly decreased in experiments and some clinical states such as genetic analbuminaemia.

**Methods.** To explain fluid homeostasis we propose osmotic counterpressure hypothesis of capillaries which claims: 1) during water filtration across microvascular wall in arterial capillary, the plasma osmolytes are sieved (retained) so that plasma osmotic counterpressure is generated, 2) this osmotic counterpressure rises along the length of capillary and when it reaches capillary hydrostatic pressure the water filtration is halted, and 3) in venous capillaries and postcapillary venules where hydrostatic pressure is low, the osmotic counterpressure is instrumental in water reabsorption from interstitium what leads to dissipation of osmotic counterpressure. According to modified van't Hoff's equation the generation of osmotic counterpressure depends on plasma concentration of osmolytes and their restricted passage (reflection coefficient) across microvascular wall in comparison to water.

**Results.** Plasma NaCl makes 83% of plasma osmolarity and shows restricted passage across the walls of cerebral and peripheral continuous capillaries, so that Na and Cl are the most important osmolytes for generation of osmotic counterpressure. Our calculation indicates that at various rates of water filtration the osmotic counterpressure of

NaCl acts as negative feedback control: higher hydrostatic pressure and water filtration rate create higher osmotic counterpressure which opposes filtration and leads to higher water reabsorption rate. Furthermore, our analysis indicates that fluid volume changes in arterial capillaries are proportionally 100 times larger than in interstitial fluid.

**Conclusion.** The osmotic counterpressure hypothesis explains fluid homeostasis at high, mean and low capillary hydrostatic pressures. Plasma proteins and inorganic electrolytes contribute 0.4% and 94% to plasma osmolarity, respectively, so that plasma proteins have low osmotic (oncotic) pressure and despite high restriction of their passage across microvascular wall they contribute little to build up of osmotic counterpressure in comparison to electrolytes. However, absence or very low concentration of plasma proteins increases microvascular wall permeability to water and osmolytes compromising build up of osmotic counterpressure leading to development of interstitial oedema.

Marin Bulat and  
Marijan Klarica

Department of Pharmacology  
and Croatian Institute for Brain  
Research, University of Zagreb  
School of Medicine, Zagreb, Croatia

### Correspondence:


Marijan Klarica MD, PhD  
Department of Pharmacology  
University of Zagreb School of  
Medicine  
Šalata 11, HR - 10 000 Zagreb,  
Croatia.  
Tel.: + 385 - 1 - 45 - 66 - 839  
Fax: + 385 - 1 - 45 - 96 - 942  
E-mail: [mklarica@hiim.hr](mailto:mklarica@hiim.hr)

## INTRODUCTION

Microvascular vessels include arterial and venous capillaries and postcapillary venules where filtration and reabsorption of water volume and exchange of solutes take place. Microvascular walls are made of single layer of flattened endothelial cells and intercellular junctions which are covered intraluminally by a negatively charged coat called glycocalyx (1,2). In the development of concept of fluid filtration and reabsorption it was assumed that water and all plasma solutes except proteins pass freely across microvascular walls. This concept is known as Starling hypothesis of capillaries named in honour of E.H. Starling who first proposed it in 1896 (3). The hypothesis was elaborated in 1963 by Landis and Pappenheimer (4), and with small modifications it represents an important chapter in contemporary textbooks of physiology. The Starling hypothesis is used to explain maintenance of fluid homeostasis in body and development of fluid disbalance in various pathological conditions leading to interstitial oedema.

The Starling hypothesis of microvessels claims that rate of fluid filtration and reabsorption of fluid volume ( $J_v$ , volume/time) across microvascular walls is regulated by hydrostatic pressure in the capillary ( $HP_c$ ) and interstitium ( $HP_i$ ) and oncotic or colloid osmotic pressure of proteins in the capillary ( $COP_c$ ) and interstitium ( $COP_i$ ) according to the equation [4]:

$$J_v = L_p [(HP_c - HP_i) - (COP_c - COP_i)] \quad [1]$$



**FIGURE 1.** Schematic presentation of the Starling oncotic hypothesis of capillaries.  $HP_c$ , hydrostatic capillary pressure;  $COP_r$ , reabsorptive oncotic pressure. Fluid filtration takes place when  $HP_c > COP_r$ , while fluid reabsorption occurs when  $HP_c < COP_r$ . For explanation see text.

where  $L_p$  is hydraulic conductivity of the microvascular wall. Positive  $J_v$  means fluid movement out of capillary (filtration), negative into capillary (reabsorption). The effects of the Starling pressures on fluid filtration and reabsorption are shown in a simplified way in Fig. 1 along capillary of 1000  $\mu\text{m}$  length.  $HP_c$  is 30 mmHg at the arterial end but due to resistance to blood flow it falls to 10 mmHg at the venous end of the capillary.  $HP_c$  tends to force fluid outwards through the capillary wall.  $HP_i$  is omitted in Fig. 1 since it is close to zero, i.e. few mmHg positive or negative (subatmospheric) in most tissues.  $COP_c$  is 25 mmHg and it tends to cause reabsorption of fluid from interstitium which is opposed by  $COP_i$  of 8 mmHg. Thus, when  $COP_i$  (8 mmHg) is subtracted from  $COP_c$  (25 mmHg) the reabsorptive oncotic pressure ( $COP_r$ ) of 17 mmHg is obtained (Fig. 1). According to the Starling oncotic hypothesis the filtration of fluid takes place in arterial part of the capillary where  $HP_c > COP_r$ , whereas reabsorption of fluid occurs in venous capillary and postcapillary venules where  $COP_r > HP_c$ . Furthermore, it is assumed that  $COP_r$  does not change significantly since volumes of filtered and reabsorbed fluid are relatively small, and that a significant part of filtered fluid is absorbed in the lymphatic capillaries.

There are some data, however, which indicate that  $COP_r$  may not be a decisive factor in regulation of fluid filtration and reabsorption. In the development of the Starling oncotic hypothesis it was assumed that only plasma proteins show restricted passage across capillary walls, while all other plasma solutes pass relatively freely and cannot exert a significant osmotic pressure between plasma and interstitium. However, it is known that cerebral capillaries are negligibly permeable to inorganic ions (5,6) and that peripheral continuous capillaries restrict significantly passage of these solutes (7,8), what should be incorporated into any comprehensive hypothesis of capillaries. The values of  $HP_c$  in Fig. 1 are used arbitrary at heart level but these values may be much higher or lower so that  $COP_r$  might not be able to control fluid filtration and reabsorption. In humans in upright position the  $HP_c$  in feet nailfolds is above 90 mmHg so that  $COP_r$  by itself could not prevent fast development of feet oedema (2,9). In the lung capillaries the  $HP_c$  is about 7 mmHg, i.e. much lower than  $COP_r$  of 17 mmHg, and it is not clear how filtration of fluid can take place (10). When plasma proteins were decreased 65% by plasmapheresis in rabbits, no brain water increase was observed indicating that the Starling oncotic hypothesis is not operative in cerebral capillaries (11). In addition, in patients with genetic analbuminaemia the  $COP_c$  is decreased by 50% without development



of oedema what is difficult to explain by the Starling hypothesis (4,12).

In attempt to resolve these problems of the Starling oncotic hypothesis, we propose the osmotic counterpressure hypothesis of the capillaries which claims that not  $COP_p$  but plasma osmotic pressure changes in the capillaries ( $OP_c$ ) are instrumental in regulation of water filtration and reabsorption. However, plasma proteins are important for maintenance of normal permeability of microvascular walls to water and solutes (4,13,14), but they contribute little to plasma effective osmotic pressure (see below).

#### ASSUMPTIONS OF THE OSMOTIC COUNTERPRESSURE HYPOTHESIS

Osmolarity is defined as concentration of osmotically active particles (osmolytes), and is usually expressed in milliosmoles per litre of water (mosm/l). Plasma and interstitial fluid osmolarities are about 300 mosm/l (15), excluding kidney as special organ which is not considered here. Plasma Na and Cl constitute 142 and 108 mosm/l, respectively (15), what makes 83% of plasma osmolarity, while contribution of other inorganic ions ( $HCO_3^-$ , K, Ca, Mg,  $HPO_4^-$ ,  $H_2PO_4^-$ ,  $SO_4^-$ ) is 11%. Thus, while inorganic ions contribute 94% to plasma osmolarity, contribution of proteins and other organic substances (glucose, urea, aminoacids, lactate, creatine) is only 0.4% and 4.3 %, respectively. Since all mentioned plasma osmolytes are hydrophilic substances their passage across cell membrane and microvascular wall may be restricted, while plasma lipophilic substances (e.g.  $CO_2$  and  $O_2$ ) diffuse easily through membranes and do not contribute to effective osmotic pressure.

Difference of effective osmotic pressure ( $\Delta OP$ ) between two compartments separated by membrane or microvascular wall is calculated according to the modified van't Hoff equation (16):

$$\Delta OP \text{ (mmHg)} = \Delta C_{\text{mosm}} \sigma RT \quad [2]$$

where  $\Delta C_{\text{mosm}}$  is concentration difference of osmolytes (mosm/l),  $\sigma$  is reflection coefficient of osmolytes, T is absolute temperature (°K) and R is universal gas constant (0.06236 mmHg per mosm/l and degree °K). At normal body temperature of 37 °C (310 °K), RT (0.06236 × 310) equals 19.3 mmHg per mosm/l (16). Reflection coefficients ( $\sigma$ ) of osmolytes indicate how they are «reflected» from microvascular walls during water passage under hydrostatic or osmotic pressure, and theoretically they may range

from 1 representing complete impermeability (100% «reflection») down to 0, for a solute permeability equal to that of water ( $\sigma = 0$ ) (2). As discussed below, the inorganic ions such as Na and Cl have  $\sigma$  significantly higher than 0 (water), so that they should affect filtration and reabsorption of water. In consideration of transcapillary movement of water and osmolytes it is usually assumed that hydrostatic pressure drives water through specific water-only pathway, while water and small osmolytes are driven through small pores and proteins through a few large pores.

Equation [2] indicates that osmotic pressure (OP) of osmolytes is linearly related to their concentration, what is not the case for plasma proteins. For calculation of COP of plasma proteins Landis and Pappenheimer developed an empirical equation (4):

$$COP \text{ (mmHg)} = 2.1 c + 0.16 c^2 + 0.009 c^3 \quad [3]$$

where c is concentration of proteins expressed in grams per 100 ml of plasma. As fluid begins to be filtered through wall of arterial capillary, its composition is determined by the rates at which different plasma osmolytes can move by convection or diffusion in comparison to water. Concentration of a plasma osmolyte with  $\sigma > 0$  should increase during water ( $\sigma = 0$ ) filtration in arterial capillary since proportionally more water than osmolyte should pass across capillary wall (17). In such a way concentration of osmolyte should increase in arterial capillary creating an osmotic counterpressure ( $OcP_c$ ) which opposes the water filtration. Thus,  $OcP_c$  is osmotic pressure increase in arterial capillary above normal OP present in systemic blood circulation. According to Equation [2] this  $OcP_c$  depends on plasma concentration and  $\sigma$  of the osmolyte, as well as on water filtration rate. The estimation of water filtration rates in peripheral continuous capillaries are 1 - 4% of the plasma volume flow depending on  $HP_c$  (1,4), while this rate should be lower in cerebral capillaries (6). Taking a range of capillary water filtration rates and known  $\sigma$  of plasma osmolytes, the capillary osmotic counterpressure ( $OcP_c$ ) opposing water filtration can be calculated (see below).

#### OSMOTIC COUNTERPRESSURE IN CEREBRAL CAPILLARIES

Cerebral capillaries form the blood-brain barrier and are characterized by endothelial cells with tight intercellular junctions which encircle completely each endothelial cell. Water permeability of cerebral capillaries is relatively high (18), while the passage of proteins and



electrolytes is very limited (6), so that reflection coefficient ( $\sigma$ ) of proteins is 0.999 (1), and  $\sigma$  of Na and Cl about 0.98 could be estimated (6, 19, 20). If we take plasma concentration of Na 142 mosm/l (see above), at 0.2% water filtration rate this number of milliosmols would be contained in 0.998 l of plasma due to water loss. When concentration of Na is recalculated per l of plasma, we obtain (142 mosm/0.998 l) 142.285 mosm/l or an increase of 0.285 mosm/l. Similar calculation for Cl shows that its normal concentration of 108 mosm/l would rise to 108.216 mosm/l, or an increase of 0.216 mosm/l. Thus, total increase of Na and Cl osmolarity (0.285 + 0.216) is 0.501 mosm/l. For calculation of osmotic pressure this value should be multiplied by osmotic coefficient for NaCl which is 0.93 (16), so we obtain (0.501 × 0.93) 0.466 mosm/l. Taking  $\sigma = 0.98$  for NaCl (see above) calculated osmotic counterpressure in cerebral capillaries ( $OcP_c$ ) according to the Equation [2] is: 0.466 × 0.98 × 19.3 = 8.81 mmHg.

Thus, at water filtration rate of 0.2% an  $OcP_c$  of NaCl about 9 mmHg is generated in arterial capillaries. The rate of water filtration can increase or decrease depending on changes of  $HP_c$ . In Table 1. are shown some values of  $OcP_c$  of NaCl at different rates of water filtration. These values of  $OcP_c$  are not permitted to run down by diffusion and/or convection of NaCl across capillary wall since they are continuously maintained by plasma flow and fluid filtration.


It can be calculated that no significant oncotic counterpressure in plasma is generated during 0.2% water filtration rate. If concentration of plasma proteins is 7 g/100 ml, than at water filtration rate of 0.2%, this concentration would increase to 7.014 g/100 ml due to water loss. When these concentrations of proteins are included in Equation [3], the calculated  $COP_c$  are 25.627 and 25.706 mmHg, respectively. Thus, calculated oncotic counterpressure of plasma proteins (25.706 – 25.627) is 0.079 mmHg which does not change when multiplied by  $\sigma$  for proteins which

**TABLE 1.** Calculated osmotic counterpressures ( $OcP_c$ ) of NaCl in cerebral and skeletal muscle capillaries (mmHg) at different water filtration rates expressed in percentages (%) of plasma volume flow.

Cerebral capillaries		Skeletal muscle capillaries	
Filtration rate	$OcP_c$ of NaCl	Filtration rate	$OcP_c$ of NaCl
0.10%	4.40 mmHg	0.25%	5.63 mmHg
0.20%	8.81 mmHg	0.50%	11.28 mmHg
0.40%	17.66 mmHg	1.00%	22.66 mmHg
0.80%	35.46 mmHg	2.00%	45.79 mmHg
1.00%	44.41 mmHg	4.00%	93.49 mmHg

is 0.999 (see above). Such a small  $OcP_c$  of plasma proteins is not physiologically significant.

$HP_c$  in cerebral capillaries is not known, but hydrostatic pressure in pial arterioles (25  $\mu$ m d.) penetrating in brain parenchyma is 55 mmHg in cats (21), while pressure in pial venules (100 – 200  $\mu$ m d.) leaving parenchyma is 4 mmHg in rats (22). This suggests that a relatively high axial gradient of hydrostatic pressure along microvascular bed is present, where filtration and reabsorption of water occur. Since  $HP_c$  falls (Fig. 1) and  $OcP_c$  of NaCl rises along length of the capillary due to filtration of water and retention (sieving) of NaCl, at a point these pressures should become equal so that filtration equilibrium is reached, i.e. water filtration is brought to halt (Fig. 2). When such hypertonic plasma is delivered to venous capillaries and postcapillary venules, where  $HP_c$  is lower than  $OcP_c$ , osmotic reabsorption of water from interstitial fluid into these vessels takes place (Fig. 2). Due to water reabsorption the hypertonic plasma is diluted and finally normalized in postcapillary venules (not shown in Fig. 2), i.e.  $OcP_c$  is dissipated. Thus, according to our osmotic counterpressure hypothesis normal osmolarity in systemic blood circulation changes in microvessels: increase of osmolarity ( $OcP_c$ ) is generated in arterial capillaries due to water filtration and NaCl sieving (retention), while in venous capillaries and postcapillary venules this increased osmolarity ( $OcP_c$ ) is dissipated due to water reabsorption so that normal plasma osmolarity is delivered to veins. In the other words, when  $HP_c > OcP_c$  filtration of wa-



**FIGURE 2.** Schematic presentation of osmotic counterpressure hypothesis of capillaries.  $HP_c$ , hydrostatic capillary pressure;  $OcP_c$ , capillary osmotic counterpressure. Fluid filtration takes place when  $HP_c > OcP_c$ , while fluid reabsorption occurs when  $HP_c < OcP_c$ . For explanation see text.

ter takes place, whereas when  $OcP_c > HP_c$  reabsorption of water occurs leading to isosmolar plasma.

To keep this analysis simple we omit here the contribution of other plasma osmolytes except NaCl to development of  $OcP_c$  during water filtration. Namely, other plasma inorganic and organic osmolytes (see above) should contribute to development of  $OcP_c$  since they show limited capillary permeability (5,6). Various transport processes across microvascular walls such as facilitated influx of D-glucose (1) and aminoacids (23) or active efflux of organic acids (24,25,26), as well as transport of some inorganic ions (27), could contribute to a long-term maintenance of osmotic homeostasis in the brain. We assume that water filtration does not change significantly volume and osmolarity of interstitial fluid because cerebral capillaries form a dense and interconnected networks of vessels so that simultaneous filtration and reabsorption of water takes place everywhere between numerous adjacent capillary branches preventing significant changes of volume and osmolarity in interstitium (see below). It should be mentioned that no lymphatic system is present in brain so that only blood microvessels are instrumental in water and solutes reabsorption.

#### OSMOTIC COUNTERPRESSURE IN PERIPHERAL CAPILLARIES

The most abundant peripheral capillaries are those with continuous uninterrupted endothelial cells connected by intercellular junctions. These capillaries are found in skeletal, smooth and cardiac muscles, skin, lungs and connective tissues (1). Some studies gave somewhat different permeabilities of continuous capillaries for inorganic ions although it seems probable that these permeabilities in various organs are similar (28). In determination of  $\sigma$  for various osmolytes in microvessels two factors seems to be especially important: composition of perfusate and rate of perfusion. When capillaries are perfused with protein-free perfusate an increase in hydraulic conductivity and osmolytes permeability of microvascular walls is observed indicating that this structure lose its selectivity (4,13,14). In addition, if rate of perfusion is not sufficiently higher than permeability-surface product of microvessels for the solute studied, its reflection coefficient is underestimated (7,8).

In skeletal and heart muscle  $\sigma$  of NaCl is 0.50 or higher (7,8). If water filtration rate in skeletal muscles is 1% of plasma volume flow, normal Na plasma concentration of 142 mosm/l should increase to 143.434 mosm/l and Cl from 108 mosm/l to 109.091 mosm/l, respectively, due to wa-

ter loss. Thus, total increase of Na and Cl ( $1.434 + 1.091$ ) is 2.525 mosm/l. When this value is multiplied by osmotic coefficient for NaCl which is 0.93, we obtain 2.348 mosm/l. Taking  $\sigma$  for NaCl 0.50, the  $OcP_c$  for NaCl according Equation [2] is:  $2.348 \times 0.50 \times 19.3 = 22.66$  mmHg.

Thus, at 1% water filtration rate the  $OcP_c$  of NaCl of about 23 mmHg is generated. In Table 1 is shown  $OcP_c$  at different water filtration rates caused by different  $HP_c$ . When  $OcP_c$  of NaCl reaches the same value as  $HP_c$  along the length of arterial capillaries the water filtration should be halted, while in venous capillaries and postcapillary venules the  $OcP_c$  should assist in water reabsorption from interstitium as suggested above for cerebral capillaries (Fig. 2). When water filtration rate in skeletal muscle capillaries is 1%, concentration of plasma proteins 7 g/100 ml and  $\sigma$  of plasma proteins 0.90 (1), the calculated oncotic counterpressure (see above) is 0.4 mmHg, a very small contribution to  $OcP_c$  of NaCl.

As already mentioned  $HP_c$  in feet capillaries of humans in upright position is somewhat above 90 mmHg (9). Assuming 4% water filtration rate in such a case, an  $OcP_c$  of above 90 mmHg would be generated as can be seen in skeletal muscle capillaries in Table 1, what should prevent medically relevant development of feet oedema. On the contrary, the  $HP_c$  in pulmonary capillaries is low, about 7 mmHg (10), indicating that both rate of water filtration and generated  $OcP_c$  should be also low. However, pulmonary capillaries have great density so that rate of water filtration and reabsorption could be considerable per gram of tissue. Furthermore, we assume that intermittent changes of  $HP_c$  as occur in vasomotion and pulse pressure should contribute to fine tuning of fluid filtration and  $OcP_c$  according to metabolic needs of tissues. Since in peripheral tissues the lymphatic system is present a part of filtered fluid is absorbed in lymphatic capillaries and returned to bloodstream by lymph flow.

To get an insight in osmotic power of total plasma  $OP_c$  in comparison to  $COP_c$  of plasma proteins we can calculate their effective osmotic pressures which would develop across microvascular wall assuming that interstitial fluid is pure water. Taking plasma osmolarity 300 mosm/l and  $\sigma = 0.50$  for all plasma osmolytes, the effective  $OP_c$  would be according to Equation [2]:  $300 \times 0.50 \times 19.3 = 2895$  mmHg, or 116 times higher than  $COP_c$  of plasma proteins which is 25 mmHg. If we take  $\sigma = 0.10$  for all plasma osmolytes, the effective  $OP_c$  would be 579 mmHg, or 23 times greater than  $COP_c$ . This analysis indicates that

osmotic power of total plasma osmolarity is much greater than  $COP_c$ , and that  $OcP_c$ , but not  $COP_c$  or  $COP_r$ , should control fluid filtration and reabsorption across microvascular walls as elaborated above.

Plasma proteins and some other blood components are important for maintenance of integrity and normal permeability of microvascular walls. When microvessels are perfused with protein-free or blood-free perfusate the hydraulic conductivity and permeability to plasma solutes of microvascular walls increases several times (13,14). Under such conditions it is expected that  $\sigma$  and  $OcP_c$  of NaCl and other plasma osmolytes should decrease what would compromise normal filtration and reabsorption of fluid across microvascular wall and lead to development of interstitial oedema.

Our hypothesis of role of NaCl in regulation of water volume passage across microvascular walls is supported by some experimental and clinical observations. When hyperosmolar NaCl solution is applied intravascularly, the absorption of water from interstitium in skeletal muscles is greatly increased (29). In addition, hyperosmolar NaCl solution applied intravascularly in patients with increased intracranial pressure leads to augmented water reabsorption from brain parenchyma and fall of increased intracranial pressure (30,31). Restricted passage of NaCl in comparison to water across microvascular walls, as suggested by our osmotic counterpressure hypothesis, explains these observations.

Fenestrated capillaries are present in some tissues such as exocrine and endocrine glands, gastrointestinal mucosa and kidney (1). These vessels are characterized by circular fenestrae or pores that penetrate the endothelium which are usually closed by a very thin diaphragm. Fenestrated capillaries show high permeability to water and inorganic ions (32). Since we were not able to find published data of  $\sigma$  of inorganic ions in those vessels it is impossible to guess at this time whether high water filtration rate could lead to such increase of their osmolarity and  $OcP_c$  which would reach the filtration equilibrium. However, such a possibility should not be a priori excluded.

#### CAPILLARY OSMOTIC COUNTERPRESSURE AS NEGATIVE FEEDBACK CONTROL

Our osmotic hypothesis indicates that the  $OcP_c$  acts as negative feedback control which opposes the water filtration: higher  $HP_c$  and water filtration rate create

higher  $OcP_c$  (Table 1), which halts water filtration (Fig. 2). Furthermore, this  $OcP_c$  in venous capillaries and postcapillary venules is instrumental in water reabsorption what dissipates  $OcP_c$ . Thus, the water filtration and reabsorption rate ( $J_v$ ) can be expressed by following relation:

$$J_v = L_p (\Delta HP_{c_i} - \Sigma OcP_c) \quad [4]$$

where  $\Delta HP_{c_i}$  is difference of  $HP_c$  and  $HP_r$ , and  $\Sigma OcP_c$  is sum of counterpressures of all plasma osmolytes with  $\sigma > 0$ . As already discussed this  $\Sigma OcP_c$  is mostly due to NaCl and other inorganic ions which constitute 94% of total plasma osmolarity.

The question arises how a sudden increase of  $HP_c$  and water filtration rate from 1% to 4% would be reflected in interstitial fluid volume. In man volumes of blood and plasma are 5 l and 3 l, respectively, volume of blood in the capillaries is 4% of total blood volume (0.20 l of blood and 0.12 l of plasma) (10,33) while volume of interstitial fluid is 12 l. When 1% of capillary plasma volume (0.0012 l) is filtered into 12 l of interstitial fluid, the interstitial fluid volume would increase by 0.01%, while at 4% filtration rate (0.0048 l) this increase would be 0.04%. Thus, volume fluid changes in the arterial capillaries are proportionally 100 times or two order of magnitude larger than those in interstitial fluid. Since filtration and reabsorption of fluid are simultaneous processes, such minute increases of interstitial fluid volumes should be easily compensated by fluid absorption. Due to such minute changes of interstitial fluid volume we assume that osmolarity and pressure of interstitial fluid change very little in comparison to such changes in microvessels.

In conclusion, our osmotic counterpressure hypothesis of the capillaries suggests that osmotic counterpressure of plasma osmolytes is the main regulator of water filtration and reabsorption across microvascular walls and principal controller of interstitial fluid volume in physiological conditions. However, when permeability of microvascular walls is increased due to various pathological processes including significant hypoproteinaemia, the reflection coefficient of plasma osmolytes and their osmotic counterpressure should decrease while hydraulic conductivity of microvascular should increase leading to development of interstitial oedema.

**Acknowledgments** This work was supported by the Croatian Ministry of Science and Technology.

**Competing interests:** The authors have declared that no competing interest exists.

### Abbreviations

- $COP_c$  – capillary colloid osmotic (oncotic) pressure  
 $COP_i$  – interstitial oncotic pressure  
 $COP_r$  – reabsorptive oncotic pressure  
 $HP_c$  – capillary hydrostatic pressure  
 $HP_i$  – interstitial hydrostatic pressure  
 $J_v$  – rate of fluid filtration or reabsorption  
 $L_p$  – hydraulic conductivity of capillary wall  
 $OcP_c$  – capillary osmotic counterpressure  
 $OP_c$  – capillary osmotic pressure

### References

- 1 Renkin E M, Crone C. Microcirculation and capillary exchange. In: Greger R, Windhorst U, editors. *Comprehensive human physiology*. Berlin: Springer; 1996. p. 1965-79.
- 2 Michel CC. Exchange of fluid and solutes across microvascular walls. In: Seldini D W, Giebisch G, editors. *The kidney*. 3rd ed. Philadelphia: Lippincott Williams & Wilkins; 2000. p 61-84.
- 3 Starling EH. On the absorption of fluids from connective tissue spaces. *J Physiol (Lond)*. 1896; 19: 312-26.
- 4 Landis EM, Pappenheimer JR. Exchange of substances through the capillary walls. In: Hamilton W F, Dow P, editors. *Handbook of physiology, Circulation Vol. II*. Washington DC: American Physiological Society; 1963. p. 961-1034.
- 5 Katzman R, Pappius HM. *Brain electrolytes and fluid metabolism*. Baltimore: Williams & Wilkins; 1973.
- 6 Fenstermacher JD, Rapoport SI. Blood-brain barrier. In: Renkin EM, Michel CC, editors. *Handbook of physiology, Microcirculation, Vol. IV*. Bethesda MD: American Physiological Society; 1984. p. 969-1000.
- 7 Yudilevich DL, Alvarez OA. Water, sodium and thiourea transcapillary diffusion in the dog heart. *Am J Physiol*. 1967; 213: 308-314.
- 8 Wolf MB, Watson PD. Measurement of osmotic reflection coefficient for small molecules in cat hindlimbs. *Am J Physiol*. 1989; 256: H282 – H290.
- 9 Levick JR, Michel CC. The effects of position and skin temperature on the capillary pressures in the fingers and toes. *J Physiol (Lond)*. 1979; 274: 97-109.
- 10 Staub NC, Dawson CA. Pulmonary and bronchial circulation. In: Greger R, Windhorst U, editors. *Comprehensive human physiology*. Berlin: Springer; 1996. p. 2071-2078.
- 11 Zornow MH, Todd MM, More SS. The acute effects of changes in plasma osmolarity and oncotic pressure. *Anesthesiology* 1987; 67: 936-941.
- 12 Benhold H, Klaus D, Scheurlen PG. Volume regulation and renal function in analbuminaemia. *Lancet* 1960; 2: 1169 – 1170.
- 13 Watson PD. Effects of blood-free and protein-free perfusion on CFC in the isolated cat hindlimb. *Am J Physiol*. 1983; 245: H911-H919.
- 14 Huxley VH, Curry FE. Differential actions of albumin and plasma on capillary solute permeability. *Am J Physiol*. 1991; 260: H1645-H1654.
- 15 Guyton AC, Hall JE. *Textbook of medical physiology*. 9th ed. Philadelphia: WB Saunders; 1996. p. 301.
- 16 Baumgarten CM, Feher JJ. Osmosis and regulation of cell volume. In: Sperelakis N, editor. *Cell physiology sourcebook*. Academic Press, San Diego, 2001. p. 319-355.
- 17 Bulat M, Klarica M. Osmotic control of intracranial pressure. *Period Biol*. 2001; 103: 293-296.
- 18 Bolwing TG, Lassen NA. The diffusion permeability to water of the rat blood-brain barrier. *Acta Physiol Scand*. 1975; 93: 415-422.
- 19 Yudilevich DL, De Rose N. Blood-brain transfer of glucose and other molecules measured by rapid indicator dilution. *Am J Physiol*. 1971; 220: 841-846.
- 20 Fraser PA, Dallas AD. Measurement of filtration coefficient in single cerebral microvessels of the frog. *J Physiol (Lond)*. 1990; 423: 343-361.
- 21 Shapiro HM, Stomberg DD, Lee DR, Wiederhielm CA. Dynamic pressures in the pial arterial microcirculation. *Am J Physiol*. 1971; 221: 279-283.
- 22 Wiig H, Reed RK. Rat brain interstitial fluid pressure measured with micropipettes. *Am J Physiol*. 1983; 244: H239-H246.
- 23 Smith QR, Stoll J. Blood-brain barrier amino acid transport. In: Pardridge WM, editor. *Introduction to the blood-brain barrier*. Cambridge: Cambridge University Press; 1998. p. 188-197.
- 24 Bulat M, Živković B. Exchange of 5-hydroxyindoleacetic acid between spinal cord and lumbar cerebrospinal fluid. *J Physiol (Lond)*. 1978; 275: 191-197.
- 25 Vladić A, Strikić N, Jurčić D, Zmajević M, Klarica M, Bulat M. Homeostatic role of the active transport in elimination of 3H-benzylpenicillin out of the cerebrospinal fluid system. *Life Sci*. 2000; 67: 2375-2385.
- 26 Zmajević M, Klarica M, Varda R, Kudelić N, Bulat M. Elimination of phenolsulfonphthalein from the cerebrospinal fluid via capillaries in the central nervous system in cats by active transport. *Neurosci Lett*. 2002; 321: 123-125.
- 27 Keep RF, Ennis SR, Betz AL. Blood-brain barrier ion transport. In: Pardridge WM, editor. *Introduction to the blood-brain barrier*. Cambridge: Cambridge University Press; 1998. p. 207-213.
- 28 Crone C, Levitt DG. Capillary permeability to small solutes. In: Renkin EM, Michel CC, editors. *Handbook of physiology, Microcirculation Vol. IV*. Bethesda MD: American Physiological Society; 1984. P. 441-466.
- 29 Hamilton MT, Ward DS, Watson PD. Effect of plasma osmolarity on steady-state fluid shifts in perfused cat skeletal muscle. *Am J Physiol*. 1993; 265: R1318-R1323.
- 30 Saltarini M, Massarutti D, Baldassarre M, Nardi G, de Cole C, Fabris G. Determination of cerebral water content by magnetic resonance imaging after small volume infusion of 18% hypertonic saline solution in a patient with refractory intracranial hypertension. *Eur J*

- Emerg Med. 2002; 9: 262-265.
- 31 Doyle JA, Davis DP, Hoyt DB. The use of hypertonic saline in the treatment of traumatic brain injury. *J Trauma* 2001; 50: 367-383.
- 32 Mann GE, Smaje LH, Yudilevich DL. Permeability of the fenestrated capillaries in the cat submandibular gland to lipid-insoluble molecules. *J Physiol (Lond)*. 1979; 297: 335-354.
- 33 Holtz J. Peripheral circulation: fundamental concepts, comparative aspects of control in specific vascular sections, and lymph flow. In: Greger R, Windhorst U, editors. *Comprehensive human physiology*. Berlin: Springer; 1996. p. 1865-1915.