
Prejac, Juraj; Višnjević, Vjeran; Drmić, Stipe; Skalny, Andrey A.; Mimica, Ninoslav; Momčilović, Berislav

Source / Izvornik: **Journal of Trace Elements in Medicine and Biology, 2014, 28, 205 - 211**

Journal article, Accepted version

Rad u časopisu, Završna verzija rukopisa prihvaćena za objavljivanje (postprint)

<https://doi.org/10.1016/j.jtemb.2014.02.001>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:100867>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-05-16**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine
Digital Repository](#)





Središnja medicinska knjižnica

Prejac J., Višnjević V., Drmić S., Skalny A. A., Mimica N., Momčilović B.
(2014) *A novel concept to derive iodine status of human populations*
from frequency distribution properties of a hair iodine concentration.
Journal of Trace Elements in Medicine and Biology, 28 (2). pp. 205-11.
ISSN 0946-672X

<http://www.elsevier.com/locate/issn/0946672X>

<http://www.sciencedirect.com/science/journal/0946672X>

<http://dx.doi.org/10.1016/j.jtemb.2014.02.001>

<http://medlib.mef.hr/2771>

University of Zagreb Medical School Repository

<http://medlib.mef.hr/>

A NOVEL CONCEPT TO DERIVE IODINE STATUS OF HUMAN POPULATIONS
FROM FREQUENCY DISTRIBUTION PROPERTIES OF A HAIR IODINE
CONCENTRATION

¹Prejac J, ²Višnjević V, ³Drmić S, ⁴Skalny AA, ⁵Mimica N, ²Momčilović B.

¹Prejac Juraj, MD, Board Certified Specialist in Internal Medicine

University Hospital Centre Zagreb

Kišpatićeva 12, 10 000 Zagreb, CROATIA

Tel. (+38591) 202 7364, e-mail: juraj.prejac@gmail.com

²Višnjević Vjeran, Master of Physics – Geophysics

Institute for Research and Development of the Sustainable Eco Systems

Srebrnjak 59, 10 000 Zagreb, CROATIA

Tel. (+38591) 536 8295, e-mail: vjeranv@gmail.com

³Drmić Stipe, MD, Board Certified Specialist in Psychiatry

Neuropsychiatric Hospital “Dr. Ivan Barbot”

Jelengradska 1, 44317 Popovača, CROATIA

Tel. (+38591) 702 265, e-mail: stipe.drmic@gmail.com

⁴Skalny Anatolyevich Andrey, MD, Board Certified Specialist in Pediatrics

ANO Center for Biotic Medicine, Moscow, RUSSIA

Zemlyanoi val 46-48, 103604 Moscow, RUSSIA

Tel. (+7491) 681 27601, e-mail: andrey_sk@microelements.ru

⁵Mimica Ninoslav, Prof., MD

University Psychiatric Hospital Vrapče,

Bolnička cesta 32, 10 090 Zagreb, CROATIA

Tel. (+3851) 378 0678, e-mail: ninoslav.mimica@bolnica-vrapce.hr

²Momčilović Berislav, Prof., MD, Corresponding author

Institute for Research and Development of the Sustainable Eco Systems

Srebrnjak 59, 10 000 Zagreb, CROATIA

Tel (+3851) 243 0288, e-mail: berislav.momcilovic@gmail.com

SHORT TITLE: Human Iodine Status

SUMMARY

Today, human iodine deficiency is next to iron the most common nutritional deficiency in developed European and underdeveloped third world countries, respectively. A current biological indicator of iodine status is urinary iodine that reflects the very recent iodine exposure, whereas some long term indicator of iodine status remains to be identified. We analyzed hair iodine in a prospective, observational, cross-sectional, and exploratory study involving 870 apparently healthy Croatians (270 men and 600 women). Hair iodine was analyzed with the inductively coupled plasma mass spectrometry (ICP MS). Population (n_{870}) hair iodine (I_H) respective median was $0.499 \mu\text{g}\cdot\text{g}^{-1}$ (0.482 and $0.508 \mu\text{g}\cdot\text{g}^{-1}$) for men and women, respectively, suggesting no sex related difference. We studied the hair iodine uptake by the logistic sigmoid saturation curve of the median derivatives to assess iodine deficiency, adequacy and excess. We estimated the overt iodine deficiency to occur when hair iodine concentration is below $0.15 \mu\text{g}\cdot\text{g}^{-1}$. Then there was a saturation range interval of about 0.15 to $2.0 \mu\text{g}\cdot\text{g}^{-1}$ ($r^2=0.994$). Eventually, the sigmoid curve became saturated at about $2.0 \mu\text{g}\cdot\text{g}^{-1}$ and upward, suggesting excessive iodine exposure. Hair appears to be a valuable and robust long term biological indicator tissue for assessing the iodine body status. We propose adequate iodine status to correspond with the hair iodine (I_H) uptake saturation of $0.565 - 0.739 \mu\text{g}\cdot\text{g}^{-1}$ (55 – 65 %).

KEY WORDS: Iodine status, hair, sex, median derivatives, hair iodine saturation capacity

INTRODUCTION

Today's lack of iodine is one of the most common nutritional deficiencies in the world that is present in both the underdeveloped third world countries, as well as in developed European countries like the United Kingdom, Italy, and Germany [1, 2]. Thus far various methods have been suggested to assess the human iodine status and to detect the iodine deficiency and/or excess to combat the nutritional iodine deficiency [3]. Today, urinary iodine (UI) excretion is conventionally considered to be tolerable approximation of the very recent dietary iodine intake. However, the determination of UI provides little useful information on the long term iodine status of an individual [4]. Since even mild iodine deficiency should be avoided [5], there is a need for a robust and reliable diagnostic indicator for assessing the iodine body status [6].

The aim of this paper is to explore how much iodine may be found in the human hair of an apparently healthy population, to study the frequency distribution of the observed iodine concentrations relative to the sex, and to estimate the possible risk of iodine deficiency and overexposure [7]. We have previously demonstrated the high reliability of hair for the multi-element profile analysis [8]; the hair is easily accessible growing biological structure of keratin fiber protein [9], easy to store and transport, and it usually has concentrations of iodine well above the detection limits necessary for accurate chemical analysis.

SUBJECTS AND METHODS

This prospective, observational, cross-sectional, and exploratory study was approved by the Ethical Committee of the Institute for the Research and Development of the Sustainable Eco Systems (IRES) and conducted by strict adherence to the Declaration of Helsinki on Human Subject Research [10], and to the complementary Croatian national bylaws and regulations [11]. Every subject gave his/her written consent to participate in the

study and filled out a short questionnaire on his/her health status and medical history [12] (data not shown).

Hair iodine (I_H) was analyzed in a random sample of 870 apparently healthy adults (270 men, 600 women), 42.6 years old on average (SD 15.7, median 46), who were concerned with their health status. They came from a general population from across the country; most of them living in Zagreb, the capital city of Croatia. All the subjects were fed their usual home prepared mixed diet, and reported no adverse medical conditions. In Croatia dietary salt is regularly iodized and today Croatia is categorized as a country having an optimal urinary iodine (UI) excretion of 100-199 $\mu\text{g}\cdot\text{L}^{-1}$ [13, 14]. Hair analysis was performed by following the International Atomic Energy Agency recommendations [15] and other validated analytical methods and procedures [16]. Approximately 0.5 – 1.0 g of the hair was cut off from the occipital head region above the *protuberantia occipitalis externa*, and stored in numbered envelopes and kept refrigerated at 4 °C before they were randomly assigned for analysis. Individual hair samples were cut prior to chemical analysis to be less than one cm long, stirred 10 min in an ethyl ether/acetone (3:1 ww), rinsed three times with the redistilled H_2O , dried at 85 °C for one hour to constant weight, immersed one hour in 5% EDTA, rinsed again in the redistilled H_2O , dried at 85 °C for 12 hours, wet digested in $\text{HNO}_3/\text{H}_2\text{O}_2$ in a plastic tube, and sonicated. The samples were analyzed for iodine content (I_H) by the inductively coupled plasma mass spectrometry (ICP MS), (Elan-9000, Perkin-Elmer, USA) at the ANO Center for Biotic Medicine (CBM), Moscow, Russia; an ISO certified high tech analytical laboratory. All chemicals were pro analysis grade (Khimmed Sintez, Moscow, Russia). We used certified GBW0910b Human Hair Reference Material (Shanghai Institute of Nuclear Research, Academia Sinica, Shanghai 201849, China [CV (SD/Mean) 0.48][17].

Current CBM iodine reference values ($\mu\text{g}\cdot\text{g}^{-1}$) for I_H are 0.65 – 9.00 and 0.65 – 8.00 for men and women, respectively. Our detection limit for hair iodine was 0.01 $\mu\text{g}\cdot\text{g}^{-1}$, and the

coefficient of variation between the assays was 0.408 (SD/Mean)[8]. Iodine belongs to the pleiad of 208 elements sharing the same mass number (number of isotopes/elements): 1 Ag, 7 Cd, 12 In, 21 Sn, 27 Sb, 26 Te, 24 I, 25 Xe, 17 Cs, 17 Ba, 12 La, 11 Ce, 6 Pr, and 2 Nd [18].

To scrutinize the hair iodine concentration frequency distribution we used the median derivative model to fit the sigmoid logistic regression analysis function for men and women separately (Appendix) [7, 19]:

$$A_2 + (A_1 - A_2) / [1 + (x/x_0)^p]$$

Where A_1 is initial value (lower horizontal asymptote), A_2 is final value (upper horizontal asymptote), x_0 is center (point of inflection, in our case it is the median M_0), p is power (the parameter that affects the slope of the area about the inflection point). The Qtiplot Data Analysis and Scientific Visualization program was used for this analysis (www.soft.proindependent.com/qtiplot.html). The same program was used to assess the saturation exponential functions.

RESULTS

Iodine was detected in all the 870 hair samples and its concentration varied over a wide range from 0.01 to 114 $\mu\text{g} \cdot \text{g}^{-1}$, with a common median M_0 (n_{870}) = 0.499 $\mu\text{g} \cdot \text{g}^{-1}$ [men (n_{270}) = 0.482 $\mu\text{g} \cdot \text{g}^{-1}$, women (n_{600}) = 0.508 $\mu\text{g} \cdot \text{g}^{-1}$] (Fig.1). The mean, SD, and CV were 1.98 ± 8.68 (4.38) and 1.01 ± 1.62 (1.60) for men and women, respectively (Fig.1, Left). The box plot data were log transformed to correct for skewedness and kurtosis and the new distribution became Gaussian (Fig.1, right). There was no discernable difference between the number of men and women above and below the common median ($p < 0.5$) when a Chi square test used. We checked the health data from the interview records and contacted 10 subjects (subjects # 861 – 870) having the highest hair iodine; in 6 of them the diagnostic X-ray contrast medium

was used within the 6 month period preceding sampling, for another 4 subjects no data could be found to help identify the source of high iodine input/intake.

Median derivatives were pinpointed according to the procedure explained in full detail in the Appendix; they are shown in Table 1, and they were used to generate the logistic sigmoid saturation curve shown in Fig.2. Three distinct regions may be identified by simple visual examination of Fig.2 – (1) the lower horizontal asymptote, (2) the linear segment, and (3) the upper horizontal asymptote. By following the principles of bioassay analysis, we consider them to represent the iodine Deficiency (1), Adequacy (2), and Excess (3), respectively.

Based upon the logistic sigmoid curve of hair iodine median derivatives we suggest that iodine concentration below $0.15 \mu\text{g}\cdot\text{g}^{-1}$ for both men and women entails the overt iodine deficiency (Fig.2). Evidently, because of such low hair iodine concentrations body metabolism is in great need of iodine, so that little may be left for the hair follicle and hair growth, which may be used to explain the poor hair quality of iodine deficient persons; thyroxine advances the onset of anagen in resting hair follicles [20]. Above the lower horizontal asymptote hair iodine concentration of $0.15 \mu\text{g}\cdot\text{g}^{-1}$, there is progressive linear upward trend of iodine accumulation in the hair that is characteristic of a physiological saturation mechanism [18]. This distinct saturation curve begins to plateau somewhat below $2.0 \mu\text{g}\cdot\text{g}^{-1}$, such that the state of hair iodine oversaturation/overexposure has been reached (here the overexposure should not be confounded with the toxicity). Thus, the linear part of iodine physiological saturation dose-response curve covers the range of about $0.15 - 2.0$ iodine $\mu\text{g}\cdot\text{g}^{-1}$. The hair iodine linear saturation range is shown for both men and women combined (Fig 2, Box A) and separately for both sexes (Fig 2, Box B); apparently there was no discernible sex dependent difference in the hair iodine between men and women.

The linear segment $d_2D_2-u_2U_2$ ($0.15 - 1.835 \mu\text{g}\cdot\text{g}^{-1}$) of Fig.2 was further analyzed for the rate of iodine incorporation into the hair, i.e., the hair iodine saturation capacity (Fig.3). Thus, saturation capacity denotes the percentage sections between the lower and upper margins of the linear range of hair iodine distribution derived from Fig.2. Apparently, the rate of change was increasing proportionally in an exponential way. This exponential growth of the hair iodine over the linear range of hair iodine concentrations ($d_2D_2-u_2U_2$) can be further analyzed as if it were a three component kinetic model (Fig.3. Insert) [21, 22, 23]. This would allow us to tentatively differentiate (a) Sparse Adequate ($\Delta_1-\Delta_6$), (b) True Adequate ($\Delta_7-\Delta_9$), and (c) Ample Adequate ($\Delta_9-\Delta_{12}$) hair iodine status. That is quite a novel possibility to present data on iodine status in a range format and what may be of great help to provide for the personalized iodine supplementation if and when needed.

Thus, the I_H concentrations of $0.209 - 0.497 \mu\text{g}\cdot\text{g}^{-1}$ (saturation capacity 20 – 50%) may be regarded as iodine sparse (not deficient but low adequate), those from $0.565 - 0.739$ (saturation 55 – 65%) true iodine adequate, and concentrations of $0.857 - 1.222$ (saturation capacity 70 – 80%) may be regarded as iodine ample (high adequate but not excessive).

DISCUSSION AND CONCLUSION

Our results on hair iodine are in agreement with the other reported values. Iyengar V [24] has reported hair iodine range of $0.4-0.8, \mu\text{g}\cdot\text{g}^{-1}$, Caroli et al. [25] reported a range $0.03-4.2 \mu\text{g}\cdot\text{g}^{-1}$, and Wolowiec et al. [26] in their review of the results of control subjects in the six different studies, reported hair iodine means of $0.62, 0.59, 0.64, 0.61, 1.55, 1.50 \mu\text{g}\cdot\text{g}^{-1}$, respectively.

Today, after a lot of refinement, the prevailing consensus is that trace element hair analysis is a valuable method for assessing the nutritional metabolic status and assessing toxicity in a non-invasive way [27-28]. To our surprise the observed hair iodine median was

$I_{H,n=870} = 0.499 \mu\text{g}\cdot\text{g}^{-1}$ (0.482 and $0.508 \mu\text{g}\cdot\text{g}^{-1}$ for men and women, respectively), and what is below the current ANO CBM concentration standard for hair iodine of $0.65\text{-}9.00$ and $0.65\text{-}8.00 \mu\text{g}\cdot\text{g}^{-1}$ for men and women, respectively. That may suggest either an inadequately high ANO CBM standard, or an inadequate nutritional iodine intake, or both. We think that the observed discrepancy between this IRES study having lower estimates of adequate iodine nutritional status and those of higher ANO CBM standards stems, in part, from ANO CBM mechanical implementation of a preconceived percentile grid upon the untransformed (log) iodine analytical data. Instead of mechanically throwing the preconceived percentile grid upon the observed data, we inferred the median derivative grid out from the data set itself (Appendix). Contrary to the recently published data on how Croatian population is well supplied with iodine according to the WHO urinary iodine data criteria [3], apparently more than half of our population has an unsatisfactory low-adequate iodine status according to the data from this study. Indeed, our true/desired adequate iodine status would require a hair iodine saturation of $0.565 - 0.739 \mu\text{g}\cdot\text{g}^{-1}$ (55 - 65%) and that ample-adequate iodine would not exceed $2.0 \mu\text{g}\cdot\text{g}^{-1}$. What would be the hair iodine toxic level remains to be elucidated. This finding further emphasizes the importance of methodological challenges in the evaluation of a chosen and nutritionally influenced biomarker like hair and/or urine [29]. In our opinion, current indicator of iodine nutritional adequacy in Croatia (urinary iodine) provided too “optimistic” results in regard to the population iodine status, and a large sparse-adequate level iodine population segment may be lurking beneath the cover of a urinary iodine indicator. Furthermore, there may be a problem in supplying and/or usage of iodized salt to/by the public. We have been under constant pressure to reduce the daily salt intake for decades, and thus reduce available dietary iodine intake. Our method of analyzing trace element median derivatives [7, 19] offers for a new way to accurately analyze samples with a large inherent variability and apparently skewed and kurtous population frequency distribution when

presented in a standard linear fashion. This study presented a novel concept to derive iodine status in human population from frequency distribution properties of hair iodine concentration in a large cohort of subjects. Such a concept of deriving the “true adequate status” is completely different from the present concept of deriving recommendations in terms of avoiding iodine deficiency, including a safety margin [30].

Based upon the results of this study we suggest hair iodine can be used as a valuable and robust indicator of the long term dietary iodine exposure. Growing at a pace of about 0.3 - 0.4 mm/day, hair is the memory tissue where the elements are irretrievably accrued; hair is the memory log of the intermediary metabolic events in homeostatic control of all the essential elements. At the same time, blood iodine and/or urinary concentrations are indicative of a short time internal balancing of this element between the various tissue compartments before it is rapidly excreted from the body. This difference in time scale for different biological indicator tissue of hair iodine (weeks) vs. urinary iodine (days) readily explains why they are incommensurable for the meaningful comparison [31]. Hair is itself a dynamic tissue structure – some 90% of hair follicles are active (anagen phase), some 10% are dormant (telogen phase), and some degenerate only to rise anew some other time [20]. Moreover, the rate of cell division in the human hair follicle is second only to the bone marrow cells, and should accurately present the metabolic changes within the body [32]. Indeed, the initial diet is only a part of the gastrointestinal input into the “black box” of intermediary metabolism before its output end point is expressed in some relevant bio-indicator tissue [33, 34]; the relationship between the entry point of dietary iodine and its end point of hair concentrations is not of a “copy-paste” variety, since hair is an actively growing tissue of itself.

The primary goal of this paper is to draw attention of clinicians and other public health personnel that body iodine status is only partially assessed by measuring the thyroid gland hormones, instead of measuring iodine directly in some suitable biological matrix tissue like

hair [11]. Indeed, when iodine intake is abnormally low, adequate secretion of thyroid hormones may still be achieved by marked modification of thyroid activity. This adaptation to iodine deficiency is triggered and maintained by increased TSH stimulation [35]. It is pertinent to note here that low concentrations of iodide (I^-) stimulate thyroid hormone synthesis independently of TSH [36]. We think that the newly implemented Clinical Practice Guidelines for Hypothyroidism in Adults [37] would be supplemented with data on hair iodine. Indeed, evidence based iodine status assessment with hair iodine, would provide a reliable guide for proper iodine prophylaxis in preventing endemic goiter and would help personalized health protection in people under the increased metabolic energy demands [38]. In conclusion; We analyzed hair iodine in 870 apparently healthy subjects (270 men and 600 women) with the ICP MS. Hair appears to be a valuable and robust biological indicator tissue for assessing long term iodine body status. We propose that the true adequate iodine status would correspond with the hair iodine (I_H) concentration of $0.565 - 0.739 \mu\text{g} \cdot \text{g}^{-1}$, i.e., 55 - 65% of hair iodine uptake saturation capacity. The evidence presented qualifies hair iodine as a long- term personal bio-indicator of human iodine status. Therefore, we believe hair iodine analysis may help the WHO&UNICEF efforts to control IDD [39, 40]. This is one of the first uses of iodine assessment using a novel source (hair) and in a relatively large cohort, as a potentially more widely-applicable use of population iodine nutrition. The data need to be complemented in the future with data that permit to establish reference ranges indicating adequate iodine intake through the assessment of other parameters such as thyroid volume, thyroglobulin levels, thyroid function tests, and urinary iodine.

ACKNOWLEDGMENTS

This study was supported in part by the Croatian Ministry of Science, Education and Sport grant No. 292-0222412-2405 (formerly No. 022-0222412-2405). We would like to

thank Prof. Glenn I Lykken and Valene L Olivier, librarian, from the UND, Grand Forks, ND, USA for their help with English language and literature references, respectively.

The generous philanthropic support of the RCM, Isle of Man, UK to the corresponding author is greatly appreciated.

DISCLOSURE STATEMENT

All the authors claim no conflict of interest.

ETHICAL CONSIDERATIONS

This study was approved by the Ethical Committee of the Institute for the Research and Development of the Sustainable Eco Systems, Zagreb, CROATIA and the permission was granted by the Croatian Ministry of Science, Education and Sport.

Momčilović Berislav, Corresponding author

Institute for Research and Development of the Sustainable Eco Systems (IRES)

Srebrnjak 59, 10 000 Zagreb, CROATIA,

Tel (+3851) 243 0288, e-mail: berislav.momcilovic@gmail.com

REFERENCES (the corresponding author pdf articles are available upon the e-mail request)

- [1] Itaka M. Iodine deficiency disorders and clinical practice. JMAJ 2004;47:371-375.
- [2] Zimmermann M, Delange F. Iodine supplementation of pregnant women in Europe: a review and recommendations. European Journal of Clinical Nutrition 2004;58:979-984.
- [3] Andersson M, Karumbunathan V, Zimmermann MB. Global iodine status in 2011 and trends over the past decade. Journal of Nutrition 2012;142:744-750.

- [4] Soldin OP. Iodine status reflected by urinary concentrations: Comparison with the USA and other countries. In: Preedy VR, Burrow GN, Watson RR, editors. *Comprehensive Handbook of Iodine: Nutritional, Biochemical, Pathological and Therapeutic Aspects*. Burlington, MA, USA: Elsevier Inc., 2009, p. 1129-1137.
- [5] Visser JT. The elemental importance of sufficient iodine intake: A trace is not enough. *Endocrinology* 2006;147:2095-2097.
- [6] Zimmermann M, Trumbo PR. Iodine. *Advances in Nutrition* 2012;4:262-265.
- [7] Momčilović B, Prejac J, Višnjević V, Mimica N, Morović S, Čelebić A, Drmić S, Skalny AV. Environmental human silver exposure. *Toxicological and Environmental Chemistry* 2012;94:1238–1246.
- [8] Momčilović B, Prejac J, Ivičić N. A case report on analytical reproducibility of the hair multielement profile: A two years follow up. *Trace Elements in Medicine (Moscow)* 2009;10(1-2):33-38.
- [9] Robbins CR. *Chemical and physical behavior of human hair*. 5th ed. Heidelberg, Germany: Springer, 2012.
- [10] Brown M. Charter of rights is adopted in the UN. *The New York Times*, August 05, 2005. <http://www.nytimes.com/learnng/general/onthisday/big//0618.html>
- [11] Momčilović B, Prejac J, Brundić S, Skalny AV, Mimica N, Drmić S. An essay on human and elements, multielement profiles, and depression. *Translational Neuroscience* 2010;1:322-34.
- [12] Oppenheim AN. *Questionnaire design, interviewing and attitude measurement*. London: Continuum, 2004.
- [13] Kusić Z, Novosel SA, Dabelić N, Punda M, Rončević S, Labar Z, Lukinac Lj, Nöthig-Hus D, Stančić A, Kaić-Rak A, Mesáros-Kanjški E, Karner I, Smoje J, Milanović

- N, Katalenić M, Juresa V, Sarnavka V. Croatia has reached iodine sufficiency. *Journal of Endocrinological Investigation* 2003;26(8):738-42.
- [14] De Benoist B, McLean E, Andersson M. Iodine deficiency: The extent of the problem. In: Preedy VR, Burrow GN, Watson RR, editors. *The Comprehensive Handbook of Iodine*. Burlington, MA, USA: Elsevier Academic Press, 2009. p. 461-467.
- [15] International Atomic Energy Agency (IAEA). *Elemental analysis of biological materials*, IAEA-TEC DOC 197. Vienna, Austria, 1980.
- [16] Burges C. *Valid analytical methods and procedures*. Cambridge, UK: The Royal Society of Chemistry, 2000.
- [17] Momčilović B, Morović J, Ivičić N, Skalny AV. Hair and blood multielement profile for metabolic imaging of the major unipolar depression. Study rationale and design. *Trace Elements in Medicine (Moscow)* 2006;7(4):33-42.
- [18] Momčilović B, Prejac J, Momčilović R, Ivičić N, Veber D, Lykken GI. On the same element isotope mass number (pleiad) and the cluster of elements sharing the same mass numbers in the Periodic system – the “Cheshuya” (fish skin) model. *Trace Elements in Medicine (Moscow)* 2008;9(3-4):5-20.
- [19] Momčilović B, Prejac J, Višnjević V, Drmić S, Mimica N, Bukovec-Megla Ž, Brundić S, Skalny AV. The Muscle Immobility of Depression—The Weightlessness Within. *Psychology* 2012;3:825-833.
- [20] Rook A, Dawber R. *Diseases of hair and scalp*. Oxford, UK: Blackwell Scientific Publications, 1982.
- [21] Harpley FW, Stewart GA, Young PA. Principles of biological assay. In: Delanuois AL, section editor. *International Encyclopedia of Pharmacology and Therapeutics*, vol.1. Oxford, UK: Pergamon Press, 1973.

- [22] Campbell PN, Smith AD. Biochemistry illustrated. 3rd ed. Edinburgh, Scotland, UK: Churchill Livingstone, 1994.
- [23] Brody TM. Concentration-response relationships In: Human Pharmacology Molecular to Clinical (Brody TM, Larner J, Minneman KP, Neu HC, editors) 2nd ed Mosby, St. Louis, MO, USA 1994, 25-32
- [24] Iyengar V. Normal values for the elemental composition of human tissues and body fluids: a new look at an old problem. Trace elements in environmental health 1985;29:277-295
- [25] Caroli S, Alimonti A, Coni E, Petrucci F, Senofonte O, Violante N. The assessment of reference values for elements in human biological tissues and fluids: a systematic review. Critical Reviews in Analytical Chemistry 1994;24:363-398
- [26] Wolowiec P, Michalak I, Chojnacka K, Mikulewicz M. Hair analysis in health assessment. Clinica Chimica Acta 2013;419:139-171
- [27] Cuttler AH. Hair test interpretation: finding hidden toxicities, 2004.
<http://www.nonmalgm.com/buythbooks.html>
- [28] Wilson L. Nutritional balancing and hair mineral analysis, 4th ed.
<http://www.drwilson.com>
- [29] Glynn RJ. Complex relations of genetic polymorphism with nutritionally influenced biomarkers. American Journal of Clinical Nutrition 2012;95:1001-1002.
- [30] Food and Nutrition Board, Institute of Medicine. Dietary reference intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. Washington, DC, USA: National Academy Press, 2002

- [31] Momčilović B, Prejac J, Višnjević V, Skalnaya MG, Mimica N, Drmić S, Skalny AV. Incommensurability of human hair and whole blood iodine. *Trace Elements in Medicine* (Moscow) 2012;13(1):20–24.
- [32] Hordinsky MK. Alopecia. In: Bologna JL, Jorizzo JL, Rapini RP, editors. *Dermatologia*. Edinburgh, UK: Mosby, 2003.
- [33] Ashby RW. *An introduction to cybernetics*. New York, NY, USA: John Willey and Sons, Inc., 1965.
- [34] Momčilović B. The epistemology of trace element balance and interaction [Plenary Lecture]. In: Hurley Ls, Keen CL, Lonerdal B, Rucker RB, editors. *Trace Elements in Man and Animals 6*. New York, NY, USA: Plenum Press, 1988, p. 173-177.
- [35] Braverman LE, Utiger RD, editors. *Werner & Ingbar's the thyroid. A fundamental and clinical text*, 9th ed. Philadelphia, PA, USA: Lippincott Williams and Wilkins, 2005.
- [36] Biesalski HK, Grimm P. *Pocket atlas of nutrition*. Stuttgart, Deutschland: Georg Thieme Verlag, 2005.
- [37] Garber JG, Cobin RH, Gharib H, Hennessey Jv, Klein I, Mechanick JI, Pessah-Pollack R, Singer PA, Woeber KA. Clinical practice guidelines for Hypothyroidism in adults. *Endocrine Practice* 2012;18:988-1038.
- [38] Lingappa VR, Farey K, editors. *Physiological medicine – clinical approach to basic medical physiology*. New York, USA: McGraw Hill Co. Inc., 2000.
- [39] World Health Organization (WHO). *Assessment of iodine deficiency disorders and monitoring their elimination. A guide for programme managers*. 3rd ed. Geneva, Switzerland: WHO, 2007.
- [40] Andersson M, De Benoist B, Darnton Hill I, Delange F, editors. *World Health Organization (WHO) and UNICEF. Iodine deficiency in Europe. A continuing public health problem*. Geneva, Switzerland: WHO, 2007.

- [41] Smylevich I, Dougherty ER. Probabilistic Boolean network. Philadelphia, PA, USA:
Society for Industrial and Applied Mathematics, 2012.

Table 1. Hair Iodine median derivative concentrations (MDC) for Men (D₁-D₆ downward MDC, U₁-U₆ upward MDC) and Women (d₁-d₆ downward MDC, u₁-u₆ upward MDC)($\mu\text{g}\cdot\text{g}^{-1}$).

MEN Median(M ₀) _{n270} = 0.482 $\mu\text{g}\cdot\text{g}^{-1}$ Iodine						WOMEN Median(M ₀) _{n600} = 0,508 $\mu\text{g}\cdot\text{g}^{-1}$ Iodine					
MDC	n	Iodine	MDC	n	Iodine	MDC	N	Iodine	MDC	n	Iodine
D ₁	135	0.200	U ₁	135	0.960	d ₁	300	0.200	u ₁	300	1.055
D ₂	68	0.150	U ₂	68	1.765	d ₂	150	0.150	u ₂	150	1.960
D ₃	34	0.150	U ₃	34	4.320	d ₃	75	0.150	u ₃	75	3.250
D ₄	17	0.086	U ₄	17	9.600	d ₄	38	0.083	u ₄	38	5.380
D ₅	9	0.041	U ₅	9	19.450	d ₅	19	0.031	u ₅	19	6.890
D ₆	5	0.024	U ₆	5	34.890	d ₆	10	0.015	u ₆	10	8.780
Common Median (M ₀) _{n870} = 0.499 $\mu\text{g}\cdot\text{g}^{-1}$ Iodine											

FIGURE LEGENDS

Figure 1. Left side. Box&whisker plot of the hair iodine log concentrations. – Min/Max, X 1% /99% percentile, \square Mean, \blacklozenge outliers, top “whiskers” = maximum, greatest value excluding outliers, bottom “whiskers” = minimum, least value excluding outliers, box: bottom line = lower quartile, 25% of data less than this value, box: top line = upper quartile, 25% of data greater than this value, box: middle line = median.

Figure 1. Right side. After the log transformation the skewed and kurtous hair iodine data follow the Gaussian frequency distribution.




Figure2. The difference between the hair Iodine median derivatives of Men_{n=270} (□) and Women_{n=600} (○) combined. D, U Men downward (D) and upward (U) median derivatives, d, u Women downward (d) and upward (u) median derivatives.

--- Logistic function: $A_2 + (A_1 - A_2)/(1 + (X/X_0)^p)$,

--- 0.95 confidence limit, ··· 0.95 prediction limit

Men: $Y = 0.978 + (-0.015 - 0.978)/[1 + (X/0.456)^{1.624}]$

Women: $Y = 0.995 + (-0.011 - 0.995)/[1 + (X/0.499)^{1.532}]$

Box A. Iodine linear saturation range for (♂+♀)(log conc).

Box B. Iodine linear saturation range separate for ♂ and ♀ (log conc).

(see Appendix for model and Table 1. for input values)




Figure 3. Hair iodine linear range saturation ($I=0.150 - 1.835 \mu\text{g}\cdot\text{g}^{-1}$, $n_{\text{♀}}+n_{\text{♂}}=870$)

--- Exponential fit $Y = 0.0397 + 0.0004e^{0.0751 \cdot X}$ ($r^2=0.998$), - - - 0.95 confidence limit, ...


0.95 prediction limit

Increment $\Delta(\text{delta})=\text{Following}(\%) - \text{Preceding}(\%)$


$$\Delta_1 - \Delta_6 = 0.0007 \cdot X + 0.2014; (R^2 = 0.958)$$

$$\Delta_7 - \Delta_9 = 0.0027 \cdot X - 0.0813; (R^2 = 0.987)$$

$$\Delta_{10} - \Delta_{12} = 0.0094 \cdot X - 0.554; (R^2 = 0.979)$$



Appendix. The median derivatives model (Population Size, PS = 1.000)



<--■-->

<--■-->

$$D_5 = D_0/32 \quad 0.016$$

$$U_5 = U_4 + U_0/32$$

0.983

<■>

<■>

$$D_6 = D_0/64 \quad 0.008$$

$$U_6 = U_5 + U_0/64$$

0.992

■

■

We studied the frequency distribution of hair iodine (H·I) median and its derivatives to assess the iodine deficiency, adequacy and excess. First we assess the median (M_0) hair iodine concentration of our subject population. By definition, one half of the studied population was above the median (upward median branch, U_0), and the other half was below the median (downward median branch, D_0). Hence, the population size (PS) for M_0 is the sum of the respective upward and downward median branches around the central inflection "hinge" M_0 , i.e., $PS = U_0 + D_0 = 0.5 + 0.5 = 1.0$. Both the respective upward and downward median branches can be further divided in the same "median of median" way into a series of sequential median derivatives ($U_{0,1,2,3 \dots n-1, n}$ and $D_{0,1,2,3, \dots n-1, n}$). For every median derivative of the population, the actual hair iodine concentration can be identified. Thus, instead of mechanically throwing the preconceived percentile grid upon the observed data, we inferred the median derivative grid out from the data set itself [41].