

The association of semaphorins 3C, 5A and 6D with liver fibrosis stage in chronic hepatitis C

Papić, Neven; Židovec Lepej, Snježana; Gorenec, Lana; Grgić, Ivana; Gašparov, Slavko; Filipec Kanižaj, Tajana; Vince, Adriana

Source / Izvornik: **PLOS ONE, 2018, 13**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/doi:10.1371/journal.pone.0209481>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:105:061232>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-08-04**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine
Digital Repository](#)



RESEARCH ARTICLE

The association of semaphorins 3C, 5A and 6D with liver fibrosis stage in chronic hepatitis C

Neven Papic^{1,2*}, Snjezana Zidovec Lepej³, Lana Gorenc³, Ivana Grgic³, Slavko Gasparov^{2,4}, Tajana Filipec Kanizaj^{2,5}, Adriana Vince^{1,2}

1 Department of Viral Hepatitis, University Hospital for Infectious Diseases Zagreb, Zagreb, Croatia, **2** University of Zagreb, School of Medicine, Zagreb, Croatia, **3** Department of Immunological and Molecular Diagnostics, University Hospital for Infectious Diseases, Zagreb, Croatia, **4** Department of Pathology and Cytology, University Hospital Merkur, Zagreb, Croatia, **5** Department of Gastroenterology, University Hospital Merkur, Zagreb, Croatia

* npapic@bfm.hr



OPEN ACCESS

Citation: Papic N, Zidovec Lepej S, Gorenc L, Grgic I, Gasparov S, Filipec Kanizaj T, et al. (2018) The association of semaphorins 3C, 5A and 6D with liver fibrosis stage in chronic hepatitis C. PLoS ONE 13(12): e0209481. <https://doi.org/10.1371/journal.pone.0209481>

Editor: Jason Blackard, University of Cincinnati College of Medicine, UNITED STATES

Received: June 3, 2018

Accepted: December 6, 2018

Published: December 28, 2018

Copyright: © 2018 Papic et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This publication was in part supported by the Croatian Science Foundation project titled "Infectomics Study of Human Liver Non-parenchymal Cells in Chronic hepatitis C" (principal investigator Professor Adriana Vince, project number IP-11-2013) as well as by the grant "Strengthening the capacity of CerVirVac for research in virus immunology and vaccinology", KK.01.1.1.01.0006, awarded to the Scientific

Abstract

Semaphorins are a diverse family of immunoregulators recently recognized to play a major role in various phases of immune responses. Their role in chronic viral hepatitis C (CHC) and contribution to the progression of liver disease is unknown. The aim of this study was to analyse the association of secreted semaphorins with the severity of liver disease in patients with CHC. Serum concentrations of semaphorins were measured in 114 treatment-naive CHC patients and 36 healthy controls. Serum concentrations of SEMA3A, SEMA3C, SEMA5A, SEMA6B and SEMA6D were significantly increased in patients with CHC compared to controls. While serum concentrations of SEMA3C and SEMA6D significantly increased with fibrosis stage in both HCV-g1 and HCV-g3 infections, the concentration of SEMA5A inversely correlated with fibrosis stage in both HCV genotypes. ROC analysis showed that serum concentrations of SEMA3C (>4.0ng/mL, AUC 0.88) and SEMA6D (>4.5, AUC 0.82) had higher AUC than widely used APRI (AUC 0.71) and FIB-4 (AUC 0.74) scores. Serum concentrations of SEMA3C and SEMA6D significantly decreased after DAA and PEG IFN- α /ribavirin therapy, while the serum concentration of SEMA5A significantly increased after DAAs therapy. Immunohistochemistry confirmed the expression of SEMA3C and SEMA5A in hepatocytes, endothelial cells and lymphocytes of cirrhotic livers from CHC patients but not in controls. In conclusion, we provide the first evidence that SEMA3C, SEMA5A and SEMA6D can be considered as markers of liver injury in CHC.

Introduction

Since its discovery 30 years ago, chronic hepatitis C (CHC) has evolved from an incurable disease to the prototype of chronic viral infection that can be easily and safely cured. Still, chronic hepatitis C remains the leading cause of cirrhosis, hepatocellular carcinoma and liver transplantation [1, 2]. Thus, the scientific focus in CHC research is moving rapidly from antiviral treatment towards complications of chronic infection and its early identification. Now we are

Centre of Excellence for Virus Immunology and Vaccines and co-financed by the European Regional Development Fund.

Competing interests: The authors have declared that no competing interests exist.

able to eliminate the virus, but significant gaps in the knowledge of biological and immunological mechanisms of chronic infection, fibrogenesis and carcinogenesis remain.

System biology approaches and whole genome sequencing analyses have enabled us to identify molecular signatures that might be associated with the lack of effective immune responses, development of tolerance and low level inflammation associated with persistent HCV infection [3–9]. In transcriptome studies of *in vitro* HCV infected liver non-parenchymal cells, the semaphorin pathway emerged as a novel consistent finding associated with HCV infection [5–7, 10].

Semaphorins are a large family of secreted and membrane-bound biological response modifiers present in many organ systems, including the nervous, epithelial, and immune system that mediate the development, angiogenesis and oncogenic transformation [11–13]. A recent research describes the role of selected semaphorins in cellular immunity and subsequent levels of inflammatory response [12, 13]. While some suppress the immune cell activation, proliferation and reduce the production of pro-inflammatory cytokines, others stimulate the immune-mediated responses [12, 13]. The importance of plexins and semaphorins has been further emphasized in allergic and immune-mediated disorders, such as systemic lupus erythematosus (SLE), asthma, psoriasis, and multiple sclerosis [14]. So far, semaphorins have not been studied in HCV infected patients.

Based on the potential correlation between semaphorins' expression and disease progression, we hypothesized that semaphorins are associated with liver fibrosis resulting from chronic viral hepatitis and therefore may be a novel biomarker of liver cirrhosis. In the present study we investigated the expression profiles of selected semaphorins in the sera and liver samples from HCV infected patients, their correlations with clinicopathological data and treatment outcomes. The potential usefulness as biomarkers for early diagnosis of liver cirrhosis was also assessed.

Materials and methods

Patients and samples

The prospective study included 114 treatment-naive adult patients with chronic hepatitis C receiving clinical care at the Croatian Reference Center for Viral Hepatitis, Zagreb, Croatia between January 2016 and September 2018. All patients tested negative for HIV and HBV infection. Additionally, paired samples before and 12 weeks after antiviral therapy were analyzed in 23 patients (14 treated with direct acting antivirals without interferon, and 9 patients treated with pegylated interferon alpha) that achieved sustained virological response (SVR, defined as undetectable HCV RNA at 12 weeks after the end of antiviral therapy). 36 voluntary blood donors without relevant medical history and without obesity, the use of medication and/or alcohol abuse were included as healthy controls (HC).

The patients were enrolled consecutively upon signing the informed consent. Ethical approval was obtained from the Ethics Committee at the University Hospital for Infectious Diseases Zagreb, Croatia, in accordance with the declaration of Helsinki.

Whole blood samples were collected by venipuncture into sterile tubes without anticoagulant and the blood was allowed to clot for 20 minutes at room temperature. Serum was separated by centrifugation at 2000xg for 10 min. The samples were divided into aliquots to avoid repeated freeze and thaw cycles and stored at -80°C until testing.

Additionally, immunohistochemistry was performed on specimens obtained from 6 patients at the Liver Transplant Center (University Hospital Merkur, Zagreb, Croatia). Three specimens were obtained from healthy liver donors immediately after the explantation and three from the explanted cirrhotic CHC livers.

Methods

The routine workup included complete blood count, liver function tests, HCV and HBV markers and HIV Ag/Ab, which were performed using commercially available assays. HCV RNA was quantified by using the standardized real-time PCR assay Abbott Real Time HCV test (Abbott Molecular, Illinois, USA) with a lower limit of detection of 12 IU/mL.

HCV genotypes were determined by using the Versant HCV genotype 2.0 Assay (Siemens Healthcare Diagnostics, Erlangen, Germany). The stage of liver fibrosis was assessed by transient elastography (Fibroscan, Echosens, Paris, France) [15]. APRI and FIB-4 score were calculated for all patients and were used as a surrogate marker of disease severity [16, 17]. The quantification of semaphorins was performed by using the standardized ELISA assays (Human Semaphorin -3A, -4D, -3C, -5A, -6B and -6D ELISA Kit, MyBioSource, San Diego, CA, USA), as recommended by the manufacturer. Quantification ranges of semaphorin ELISAs were as follows: SEMA-3A (31,25–2000 pg/ml), SEMA-4D (0,625–20 ng/ml), SEMA-3C 0–10 ng/ml, SEMA-5A (0,78–50 ng/ml), SEMA-6B (0,625–20 ng/ml) and SEMA-6D (0,625–20 ng/ml).“

Immunohistochemical staining: unstained slides (4 μ m thick) were generated from the whole tissue blocks (2 blocks per case) for immunohistochemical staining using antibodies SEMA3C and SEMA5A. Immunohistochemical staining was performed in the automated immunostainer (Dako Autostainer, LINK 48, Denmark) using Envision (DAKO Denmark) for SEMA3C and ImmPRESS (Vector laboratories, USA) polymer detection systems for SEMA5A according to the manufacturer's instructions. Rabbit polyclonal anti-semaphorin 3C antibody (Abcam, Cambridge, UK) was used at 1:50 dilution. Goat polyclonal anti-semaphorin 5A antibody (Abcam, Cambridge, UK) was used at 1:80 dilution. Human pancreas tissue was used as a positive control for SEMA3C expression and human colon tissue as a positive control for SEMA5A expression (both cytoplasmic staining). For the negative control, adjacent tissue sections were stained without the primary antibody.

Statistical analysis

The demographic, clinical characteristics and laboratory data were evaluated and presented descriptively. Categorical data were expressed in frequencies and relevant percentage. The significance of the observed differences in frequencies between relevant subgroups was tested with the chi-square test or the Fisher's exact test when appropriate. For continuous variables, we calculated mean values with standard deviations or median values with 25th and 75th percentile depending on distribution. Comparative statistics between two or more groups was generated using the Mann–Whitney U or the Kruskal–Wallis test, as appropriate. Correlations were analyzed using the Spearman's rank correlation coefficient. The diagnostic performances of the laboratory variables considered were compared using a receiver operating characteristic (ROC) analysis. Sensitivity, specificity, positive and negative predictive values and likelihood ratios (LRs) were calculated. All tests were two-tailed; a $P < 0.05$ was considered statistically significant. Statistical analyses were performed using prism (ver 5.0) statistical software (Graphpad Software, San Diego, CA) and MedCalc for Windows, ver 11.5.1.0 (MedCalc Software, Mariakerke, Belgium).

Results

Patient characteristics

The characteristics of the patients are shown in [Table 1](#). The majority of CHC patients were infected with HCV genotype 1 ($n = 70$) and 44 patients were infected with genotype 3. CHC

Table 1. Baseline patient characteristics.

	Fibrosis stage			
	1	2	3	4
Liver stiffness, kPa	5.5 (4.6–6.1)	7.9 (7.3–8.8)	10.2 (9.6–11)	19.2 (14.0–30.7)
Age, years	51.5 (43.7–65.5)	57 (51.75–65.25)	59.5 (41.5–66.7)	59 (48.7–63)
Male sex	10 (8.8%)	12 (10.5%)	12 (10.5%)	22 (19.3%)
HCV genotype 1	12 (10.5%)	14 (12.3%)	18 (15.8%)	26 (22.8%)
HCV genotype 3	10 (8.8%)	8 (7.0%)	10 (8.8%)	16 (14.0%)
HCV RNA, log 10	5.1 (4.9–5.4)	6.3 (5.2–6.7)	5.5 (4.5–6.1)	5.8 (5.2–6.1)
Aspartate transaminase, IU/L	45 (34.5–87.5)	61.5 (47–129)	79 (61.2–89)	82 (60.5–101.3)
Alanine transaminase, IU/L	67.5 (39.2–124.5)	77 (58.7–148)	97 (68–124.8)	92 (57–127.5)
Platelets, $\times 10^9/L$	230 (184–300)	202 (147–231)	131.5 (98.2–194.3)	122.5 (78–220.5)
Albumins, g/L	43 (38.3–45.3)	42 (37.5–44.7)	38.5 (35–42.0)	34.4 (26.7–40.8)
International normalized ratio	1 (0.99–1.12)	1.0 (0.9–1.1)	1.13 (1.0–1.3)	1.3 (1.1–1.8)
APRI score	0.53 (0.37–0.68)	0.93 (0.58–1.72)	1.69 (0.85–2.5)	1.57 (0.82–3.01)
FIB4 score	1.21 (0.89–2.14)	2.3 (1.5–4.2)	3.08 (2.08–5.19)	3.31 (2.05–6.35)

Presented are baseline patient clinical, laboratory and virological characteristics according to fibrosis stage as measured by transient elastography. Data are presented as frequencies (%) or medians with interquartile ranges.

<https://doi.org/10.1371/journal.pone.0209481.t001>

patients were classified into four groups according to the transient elastography (F1 n = 22; F2 n = 22, F3 n = 28, F4 n = 42). The study also included 36 healthy controls (19 males, median age of 51 years (IQR 39–59.7)).

Elevated serum concentrations of semaphorins in patients with chronic hepatitis C

We examined serum semaphorin concentrations in CHC group compared to healthy controls. In comparison to healthy controls, serum levels of SEMA3A, SEMA3C, SEMA5A, SEMA6B and SEMA6D were significantly higher in CHC patients, whereas only SEMA4D was not significantly altered, as shown in Fig 1A. There was no difference in semaphorin expression in healthy controls regarding sex and age.

Serum concentrations of SEMA3C, SEMA5A and SEMA6D correlate with fibrosis stage

Next, we analyzed serum concentrations of semaphorins in different fibrosis stages and HCV genotype infections (Fig 1B–1D). While serum concentrations of SEMA3C and SEMA6D significantly increased with fibrosis stage in both HCV-g1 and HCV-g3 infection, serum concentration of SEMA5A inversely correlated with fibrosis stage in both HCV genotypes. SEMA3A and SEMA6B serum concentrations showed no correlation with fibrosis stage.

In CHC patients, both SEMA3C and SEMA6D showed correlation with the APRI ($r = 0.38$, $p < 0.0001$ and $r = 0.36$, $p < 0.0001$, respectively) and FIB4 score ($r = 0.39$, $p < 0.0001$ and $r = 0.37$, $p < 0.0001$). SEMA3C and SEMA6D showed positive correlation with AST ($r = 0.25$, $p = 0.0052$ and $r = 0.25$, $p = 0.0047$, respectively), and negative correlation with platelet count ($r = -0.35$, $p < 0.0001$ and $r = -0.33$, $p = 0.0003$). SEMA5A inversely correlated with AST ($r = -0.23$, $p = 0.0125$) and APRI ($r = -0.36$, $p < 0.0001$) and FIB4 ($r = -0.37$, $p < 0.0001$), and positively with platelet count ($r = 0.35$, $p = 0.0001$). Neither SEMA3C, SEMA6D nor SEMA5A correlated with age, viraemia and ALT.




Fig 1. Serum concentrations of semaphorins in patients with chronic hepatitis C (CHC) and healthy controls. (A) Patients with CHC have elevated serum levels of semaphorins compared to healthy controls. Serum concentration of SEMA5A decreases with fibrosis progression (B), while serum concentrations of SEMA3C (C) and SEMA6D (D) increase with fibrosis stage. Data are presented as median with interquartile range and analyzed by Mann-Whitney *U*-test or Kruskal-Wallis test, as appropriate.

<https://doi.org/10.1371/journal.pone.0209481.g001>

Serum SEMA3C and SEMA6D are superior to APRI and FIB-4 in predicting the presence of cirrhosis and significant fibrosis

Based on the above-mentioned correlations, the diagnostic performances of the selected semaphorins were compared to routinely used APRI [16] and FIB-4 [17] scores using a receiver operating characteristic (ROC) analysis.

Firstly, serum concentrations of SEMA3C and SEMA6D showed better sensitivity and specificity for detection of liver cirrhosis (>4.0, sensitivity 87.5%, specificity 78.8%, AUC 0.88 and >4.5, sensitivity 72.9%, specificity 85.7%, AUC 0.82, respectively) as compared to widely used APRI score (>1.4 sensitivity 68.7%, specificity 69.1%, AUC 0.71) or FIB-4 score (>3.25




Fig 2. The ROC curve of laboratory parameters for discrimination of liver cirrhosis (F4) (A) and significant liver fibrosis (F2-4) (B). Shown are calculated cut-off values (Youden index), accuracy as measured by the area under the ROC curve (AUC) and probability as calculated p-values.

<https://doi.org/10.1371/journal.pone.0209481.g002>

sensitivity 64.5%, specificity 70.4%, AUC 0.74), as shown in Fig 2A. Serum concentration of SEMA5A <12ng/mL showed sensitivity of 85.0% and specificity of 63.3% (AUC 0.79) in the detection of cirrhosis.

Secondly, we examined the diagnostic accuracy in detection of significant fibrosis (F2-F4). Again, SEMA3C >3.5 (sensitivity 78.3%, specificity 89.4%, AUC 0.88), SEMA6D >3.0 (sensitivity 77.0%, specificity 95.4%, AUC 0.90) and SEMA5A <15 (sensitivity 85.3%, specificity 72.3%, AUC 0.83) showed comparable discrimination of significant fibrosis as did APRI (AUC 0.84) and FIB4 (AUC 0.85) scores, as shown in Fig 2B.

Serum levels of SEMA3C and SEMA6D decrease after DAA and PegIFN α treatment

Next, we examined serum concentrations of semaphorins in 14 patients treated with DAAs and 9 patients treated with PegIFN α before and 12 weeks after completion of treatment. At baseline, liver fibrosis was classified as F1/2 in 39% and F3/4 in 61% of patients. Antiviral therapy significantly improved fibrosis stage, as measured by fibroscan 12 weeks after treatment completion, with 60% of cases classified as F1/2 after therapy (Fig 3A). After treatment all patients had AST and ALT values within reference range.

Importantly, serum concentrations of SEMA3C and SEMA6D significantly decreased after DAA (4.35, IQR 3.62–5.03 to 0.64, IQR 0.28–0.96, $p < 0.0001$, and 4.13, IQR 1.2–5.85 to 1.69, IQR 0.84–2.67, $p = 0.0340$, respectively) and PegIFN α therapy (3.49, IQR 2.00–5.03 to 1.46, IQR 0.96–2.1, $p = 0.0083$ and 3.17, IQR 1.81–4.87 to 1.23, IQR 0.59–2.19, $p = 0.0244$, respectively). The serum concentration of SEMA5A significantly increased after DAAs therapy (11.05, IQR 8.67–15.75 to 15.75, IQR 13.65–22.88, $p = 0.0241$), but there were no significant changes in patients treated with PegIFN α (Fig 3B–3D). There were no significant changes in serum concentrations of SEMA6B.




Fig 3. Serum semaphorins levels and liver fibrosis in patients before and after antiviral therapy. Changes in the proportion of fibrosis (F1-4) (A), serum SEMA3C (B), SEMA6D (C) and SEMA5A (D) in patients before and after antiviral therapy are shown. Paired patient concentrations are shown (n = 23). Data were analyzed by Mann-Whitney U-test.

<https://doi.org/10.1371/journal.pone.0209481.g003>

SEMA3C and SEMA5A expression in liver tissue samples

To gain further insight into tissue expression of SEMA3C and SEMA5A, we performed immunohistochemistry in cirrhotic CHC human livers in comparison to healthy donors. In healthy livers SEMA3C and SEMA5A expression was not detected by immunohistochemistry, neither in hepatocytes, nor in endothelial cells (Fig 4A and Fig 5A). By contrast, in CHC cirrhotic livers SEMA3C and SEMA5A expression was detected in hepatocytes as well in endothelial cells and lymphocytes. In HCV cirrhosis, SEMA3C showed moderate positivity in endothelial cells and lymphocytes (Fig 4C) and weak positivity in hepatocytes (Fig 4D). SEMA5A was expressed mostly in hepatocytes (Fig 5B and 5C). In addition, SEMA3C and SEMA5A were expressed in HCV HCC samples.

Discussion

During chronic HCV infection, a sustained cytokine production drives persistent low-grade inflammation in the absence of immunocompetent Th1 responses thus allowing the constant




Fig 4. Immunohistochemical staining of SEMA3C in healthy and cirrhotic HCV liver. While there is no expression of SEMA3C markers in healthy liver tissue (A, magnification 40x), in cirrhotic HCV liver (B, magnification 10x), there is a moderate expression in endothelial cells and lymphocytes (C, magnification 40x) and weak in hepatocytes (D, magnification 40x).

<https://doi.org/10.1371/journal.pone.0209481.g004>

viral replication [18]. However, the understanding of these phenomena remains unclear. The impairment of balance in the cytokine milieu modulates local T helper and Treg responses, which are unable to eliminate the infection, but drive hepatic damage and fibrogenesis [19]. Indeed, many cytokines have shown to be good predictors of interferon treatment responses [19, 20], however they failed to show the clinically relevant diagnostic accuracy in identifying patients with severe fibrosis or therapeutic application.

There is growing evidence that the signaling pathways triggered by semaphorin interactions with their receptors plexins and neuropilins influence the outcome of the immune responses; while some suppress immune cell activation and proliferation, others stimulate immune-mediated responses [21]. Here we provide the first evidence that semaphorins might play a role in HCV infection as well.

Firstly, we demonstrated that circulating levels of SEMA3A, SEMA3C, SEMA5A, SEMA6B and SEMA6D are detectable and elevated in the serum of patients with CHC. Class 3




Fig 5. Immunohistochemical staining of SEMA5A in healthy and cirrhotic HCV liver. While there is no expression of SEMA5A markers in healthy liver tissue (A, magnification 40x), there is a strong expression in HCV cirrhotic livers (B, magnification 10x), in endothelial cells and lymphocytes and in hepatocytes (C, magnification 40x).

<https://doi.org/10.1371/journal.pone.0209481.g005>

semaphorins, specifically SEMA3A, previously shown to inhibit the T-cell activation, act as a positive regulator in the innate immune system by promoting proinflammatory cytokines production [12, 13, 22]. The importance of SEMA3A in autoimmune disorders is highlighted by the finding of reduced expression of SEMA3A in patients with rheumatoid arthritis and SLE that correlates with T-cell mediated inflammation and disease severity [21]. Less is known about the immunoregulatory role of SEMA3C, but it was shown that SEMA3C promotes the migration of dendritic cells [12, 13]. SEMA4D, activator of T- and B- cells, was shown to induce inflammation [12, 13]. SEMA5A promotes T-cell and NK cell proliferation and induces the secretion of Th17 proinflammatory cytokines that are implicated in SLE and RA disease activity [12, 14, 21]. SEMA6D is involved in the regulation of late phases of T-cell activation and generation of T-cell memory [12, 21].

Secondly, we found that semaphorin expression correlates with fibrosis stage. The role of semaphorins in fibrogenesis has not been studied, so the pathophysiological explanation of our findings is to be determined. Our data show that SEMA3C and SEMA6D significantly increase with fibrosis stage. Interestingly, pro-fibrogenic role of SEMA3C was shown in adipose tissue where SEMA3C increased the production of extracellular matrix and growth factors that are associated with the pathogenesis of fibrosis [23]. Recently, class III semaphorins were shown to induce the migration and invasive capacity of fibroblast-like synoviocytes in the model of rheumatoid arthritis [24]. SEMA6D activates vascular endothelial growth factor-receptor 2 implicated in liver fibrinogenesis [25]. We found that serum concentration of SEMA5A inversely correlates with fibrosis. So far it was shown that SEMA5A regulates angiogenesis by increasing endothelial cell proliferation, decreasing apoptosis and extracellular matrix degradation through matrix metalloproteinase [26].

Next, we have shown that SEMA3C and SEMA6D have superior accuracy to APRI and FIB-4 in predicting the presence of cirrhosis. Early and accurate assessment of the severity of liver fibrosis is essential in the management and prognosis of patients with CHC and CHB. Two most common non-invasive scores, APRI and FIB-4, that don't require extra laboratory testing, are limited in prediction of significant fibrosis with relatively low sensitivity (60% and 75%, respectively) and specificity (65% and 60%, respectively) [27].

Additionally, the concentrations of SEMA3C and SEMA6D decrease following treatment with antiviral drugs (DAA) or a combination of immunomodulatory and antiviral drugs (PegIFN α and ribavirin). These results show that successful treatment resulting in viral eradication reduces the intensity of semaphorin synthesis. Additionally, they suggest that HCV infection itself, particularly in liver hepatocytes as shown by immunohistochemistry, might represent a key stimulus for semaphorin synthesis in chronic infection.

Although our results are limited by the relatively small number of patients, we provide the first evidence of the role of semaphorins in HCV infection. Due to ethical issues we assessed the stage of liver fibrosis by non-invasive technics only, as recommended and described in current international guidelines.

In conclusion, we demonstrated that circulating levels of SEMA3A, SEMA3C, SEMA5A, SEMA6B and SEMA6D are detectable and elevated in the serum of patients with CHC. Based on the correlation with the biochemical and fibroelastography degree of the liver damage, SEMA3C, SEMA5A and SEMA6D can be considered markers of liver injury in CHC and are superior to APRI and FIB-4 in predicting the presence of cirrhosis. The regulation of

semaphorins and their receptors could be a potential future therapeutic target to decrease liver inflammation and fibrogenesis in chronic viral hepatitis.

Acknowledgments

The authors acknowledge professor Arijana Pavelic for editing and proofreading the manuscript.

Author Contributions

Conceptualization: Neven Papic, Snjezana Zidovec Lepej, Adriana Vince.

Data curation: Lana Gorenec, Ivana Grgic, Tajana Filipec Kanizaj.

Formal analysis: Neven Papic, Ivana Grgic, Slavko Gasparov.

Funding acquisition: Adriana Vince.

Investigation: Snjezana Zidovec Lepej, Lana Gorenec, Slavko Gasparov, Tajana Filipec Kanizaj, Adriana Vince.

Methodology: Neven Papic, Snjezana Zidovec Lepej, Lana Gorenec, Ivana Grgic, Slavko Gasparov.

Project administration: Neven Papic, Adriana Vince.

Supervision: Snjezana Zidovec Lepej, Adriana Vince.

Validation: Tajana Filipec Kanizaj.

Writing – original draft: Neven Papic.

Writing – review & editing: Snjezana Zidovec Lepej, Lana Gorenec, Ivana Grgic, Slavko Gasparov, Tajana Filipec Kanizaj, Adriana Vince.

References

1. Maaroufi A, Vince A, Himatt SM, Mohamed R, Fung J, Opare-Sem O, et al. Historical epidemiology of hepatitis C virus in select countries-volume 4. *J Viral Hepat.* 2017; 24 Suppl 2:8–24. <https://doi.org/10.1111/jvh.12762> PMID: 29105285
2. Stanaway JD, Flaxman AD, Naghavi M, Fitzmaurice C, Vos T, Abubakar I, et al. The global burden of viral hepatitis from 1990 to 2013: findings from the Global Burden of Disease Study 2013. *Lancet.* 2016; 388(10049):1081–8. [https://doi.org/10.1016/S0140-6736\(16\)30579-7](https://doi.org/10.1016/S0140-6736(16)30579-7) PMID: 27394647
3. Blackham S, Baillie A, Al-Hababi F, Remlinger K, You S, Hamatake R, et al. Gene expression profiling indicates the roles of host oxidative stress, apoptosis, lipid metabolism, and intracellular transport genes in the replication of hepatitis C virus. *Journal of virology.* 2010; 84(10):5404–14. <https://doi.org/10.1128/JVI.02529-09> PMID: 20200238
4. Folkers ME, Delker DA, Maxwell CI, Nelson CA, Schwartz JJ, Nix DA, et al. ENCODE tiling array analysis identifies differentially expressed annotated and novel 5' capped RNAs in hepatitis C infected liver. *PLoS one.* 2011; 6(2):e14697. <https://doi.org/10.1371/journal.pone.0014697> PMID: 21359205
5. Negash AA, Ramos HJ, Crochet N, Lau DT, Doehle B, Papic N, et al. IL-1beta production through the NLRP3 inflammasome by hepatic macrophages links hepatitis C virus infection with liver inflammation and disease. *PLoS pathogens.* 2013; 9(4):e1003330. <https://doi.org/10.1371/journal.ppat.1003330> PMID: 23633957
6. Papic N, Delker D, Liu S, Rong X, Vince A, Hagedorn C. RNA SEQUENCING ANALYSIS OF HUMAN LIVER SINUSOIDAL ENDOTHELIAL CELLS REVEALS EVIDENCE FOR AN ANTI-INFLAMMATORY ROLE DURING HCV INFECTION. *Journal of hepatology.* 2013; 58:S146–S7. [https://doi.org/10.1016/S0168-8278\(13\)60355-4](https://doi.org/10.1016/S0168-8278(13)60355-4)
7. Papic N, Maxwell CI, Delker DA, Liu S, Heale BS, Hagedorn CH. RNA-sequencing analysis of 5' capped RNAs identifies many new differentially expressed genes in acute hepatitis C virus infection. *Viruses.* 2012; 4(4):581–612. <https://doi.org/10.3390/v4040581> PMID: 22590687

8. Walters KA, Syder AJ, Lederer SL, Diamond DL, Paeper B, Rice CM, et al. Genomic analysis reveals a potential role for cell cycle perturbation in HCV-mediated apoptosis of cultured hepatocytes. *PLoS pathogens*. 2009; 5(1):e1000269. <https://doi.org/10.1371/journal.ppat.1000269> PMID: 19148281
9. Woodhouse SD, Narayan R, Latham S, Lee S, Antrobus R, Gangadharan B, et al. Transcriptome sequencing, microarray, and proteomic analyses reveal cellular and metabolic impact of hepatitis C virus infection in vitro. *Hepatology*. 2010; 52(2):443–53. <https://doi.org/10.1002/hep.23733> PMID: 20683944
10. Vince AP, N. Unresolved issue in hepatitis C: The role of liver non-parenchymal cells and semaphorins. *Infect Dis Rep*. 2018; 10(1). Epub 29 March 2018. <https://doi.org/doi.org/10.4081/idr.2018.7651>
11. Capparuccia L, Tamagnone L. Semaphorin signaling in cancer cells and in cells of the tumor microenvironment—two sides of a coin. *J Cell Sci*. 2009; 122(Pt 11):1723–36. <https://doi.org/10.1242/jcs.030197> PMID: 19461072
12. Kumanogoh A, Kikutani H. Immunological functions of the neuropilins and plexins as receptors for semaphorins. *Nature reviews Immunology*. 2013; 13(11):802–14. PMID: 24319778
13. Roney K, Holl E, Ting J. Immune plexins and semaphorins: old proteins, new immune functions. *Protein & cell*. 2013; 4(1):17–26. <https://doi.org/10.1007/s13238-012-2108-4> PMID: 23307780
14. Vadasz Z, Toubi E. Semaphorins: their dual role in regulating immune-mediated diseases. *Clin Rev Allergy Immunol*. 2014; 47(1):17–25. <https://doi.org/10.1007/s12016-013-8360-4> PMID: 23397481
15. Castera L, Fornis X, Alberti A. Non-invasive evaluation of liver fibrosis using transient elastography. *Journal of hepatology*. 2008; 48(5):835–47. <https://doi.org/10.1016/j.jhep.2008.02.008> PMID: 18334275
16. Lin ZH, Xin YN, Dong QJ, Wang Q, Jiang XJ, Zhan SH, et al. Performance of the aspartate aminotransferase-to-platelet ratio index for the staging of hepatitis C-related fibrosis: an updated meta-analysis. *Hepatology*. 2011; 53(3):726–36. <https://doi.org/10.1002/hep.24105> PMID: 21319189
17. Sterling RK, Lissen E, Clumeck N, Sola R, Correa MC, Montaner J, et al. Development of a simple non-invasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology*. 2006; 43(6):1317–25. <https://doi.org/10.1002/hep.21178> PMID: 16729309
18. Thimme R, Bukh J, Spangenberg HC, Wieland S, Pemberton J, Steiger C, et al. Viral and immunological determinants of hepatitis C virus clearance, persistence, and disease. *Proceedings of the National Academy of Sciences of the United States of America*. 2002; 99(24):15661–8. <https://doi.org/10.1073/pnas.202608299> PMID: 12441397
19. Larrubia JR, Moreno-Cubero E, Lokhande MU, Garcia-Garzon S, Lazaro A, Miquel J, et al. Adaptive immune response during hepatitis C virus infection. *World journal of gastroenterology: WJG*. 2014; 20(13):3418–30. <https://doi.org/10.3748/wjg.v20.i13.3418> PMID: 24707125
20. Kurelac I, Lepej SZ, Grljic I, Gorenec L, Papic N, Dusek D, et al. Chemokine CXCL10 at week 4 of treatment predicts sustained virological response in patients with chronic hepatitis C. *J Interferon Cytokine Res*. 2012; 32(8):386–91. <https://doi.org/10.1089/jir.2012.0006> PMID: 22799464
21. Nishide M, Kumanogoh A. The role of semaphorins in immune responses and autoimmune rheumatic diseases. *Nat Rev Rheumatol*. 2018; 14(1):19–31. <https://doi.org/10.1038/nrrheum.2017.201> PMID: 29213125
22. Wen H, Lei Y, Eun SY, Ting JP. Plexin-A4-semaphorin 3A signaling is required for Toll-like receptor- and sepsis-induced cytokine storm. *The Journal of experimental medicine*. 2010; 207(13):2943–57. <https://doi.org/10.1084/jem.20101138> PMID: 21098092
23. Mejhert N, Wilfling F, Esteve D, Galitzky J, Pellegrinelli V, Kolditz CI, et al. Semaphorin 3C is a novel adipokine linked to extracellular matrix composition. *Diabetologia*. 2013; 56(8):1792–801. <https://doi.org/10.1007/s00125-013-2931-z> PMID: 23666167
24. Tang MW, Malvar Fernandez B, Newsom SP, van Buul JD, Radstake T, Baeten DL, et al. Class 3 semaphorins modulate the invasive capacity of rheumatoid arthritis fibroblast-like synoviocytes. *Rheumatology (Oxford)*. 2018. <https://doi.org/10.1093/rheumatology/kex511> PMID: 29471421
25. Lu Y, Xu Q, Chen L, Zuo Y, Liu S, Hu Y, et al. Expression of semaphorin 6D and its receptor plexin-A1 in gastric cancer and their association with tumor angiogenesis. *Oncol Lett*. 2016; 12(5):3967–74. <https://doi.org/10.3892/ol.2016.5208> PMID: 27895757
26. Sadanandam A, Rosenbaugh EG, Singh S, Varney M, Singh RK. Semaphorin 5A promotes angiogenesis by increasing endothelial cell proliferation, migration, and decreasing apoptosis. *Microvasc Res*. 2010; 79(1):1–9. <https://doi.org/10.1016/j.mvr.2009.10.005> PMID: 19850054
27. Gamil M, Alboraie M, El-Sayed M, Elsharkawy A, Asem N, Elbaz T, et al. Novel scores combining AFP with non-invasive markers for prediction of liver fibrosis in chronic hepatitis C patients. *Journal of medical virology*. 2018. <https://doi.org/10.1002/jmv.25026> PMID: 29315641