

Single nucleotide polymorphisms and health behaviours related to obesity - trawling the evidence in the prospect of personalised prevention

Fišter, Kristina; Polašek, Ozren; Vuletić, Silvije; Kern, Josipa

Source / Izvornik: **Studies in Health Technology and Informatics, 2009, 150, 762 - 766**

Journal article, Accepted version

Rad u časopisu, Završna verzija rukopisa prihvaćena za objavljivanje (postprint)

<https://doi.org/10.3233/978-1-60750-044-5-762>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:380084>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-07-10**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine
Digital Repository](#)





Središnja medicinska knjižnica

Fišter K., Polašek O., Vuletić S., Kern J. (2009) *Single nucleotide polymorphisms and health behaviours related to obesity - trawling the evidence in the prospect of personalised prevention. Studies in health technology and informatics, 150. pp. 762-766. ISSN 0926-9630*

<http://booksonline.iospress.nl/Content/View.aspx?piid=64>

<http://dx.doi.org/10.3233/978-1-60750-044-5-762>

<http://medlib.mef.hr/677>

University of Zagreb Medical School Repository

<http://medlib.mef.hr/>

Single nucleotide polymorphisms and health behaviours related to obesity—trawling the evidence in the prospect of personalised prevention

Kristina FIŠTER^{a,1}, Ozren POLAŠEK^b, Silvije VULETIĆ^b and Josipa KERN^b

^a*Department of medical sociology and health economics, Andrija Štampar School of Public Health, Zagreb, Croatia*

^b*Department of medical statistics, epidemiology and medical informatics, Andrija Štampar School of Public Health, Zagreb, Croatia*

Abstract. Efforts aimed at primary and secondary prevention of cardiovascular diseases, the major killer of contemporary adult populations, largely rely on modification of risk behaviours related to smoking, physical activity, dietary intake, and alcohol consumption, and also control of obesity and hypertension, the interim risk states between health and disease. We propose that the extent to which the gene x ‘obesogenic’ environment interaction depends on associations between particular single nucleotide polymorphisms (SNPs) and behavioural risk factors for overweight or obesity determines opportunities for novel, personalised preventive interventions. We systematically searched for SNPs that might be of interest for this postulate and we present various SNPs that have been shown to be associated with overweight or obesity and behavioural risk factors for developing

¹ Corresponding Author: Kristina Fišter, Andrija Štampar School of public health, Rockefellerova 4, HR-10000 Zagreb, Croatia; E-mail: kfister@snz.hr

these traits, and thus hold promise for future design of personalised preventive interventions.

Keywords. cardiovascular diseases, obesity, genomics, single nucleotide polymorphism, health behaviour

Introduction

Efforts aimed at primary and secondary prevention of cardiovascular diseases, the major killer of contemporary adult populations [1-3], largely rely on modification of risk behaviours related to smoking, physical activity, dietary intake, and alcohol consumption [4, 5]. Control of obesity and hypertension, the interim risk states between health and disease, constitute another large part of preventive ventures. In the public health that increasingly turns its attention toward genomics, a major challenge is to understand the role of genetic variants in susceptibility to chronic diseases and associated risk factors [6]. This has required characterising the nature of gene variation, assembling an extensive catalogue of single nucleotide polymorphisms (single base-pair mutations that occur at a specific site in the DNA sequence, SNPs) in candidate genes, and performing association and other gene mapping studies.

A step further, we need to incorporate findings from genomics in real life interventions. Beyond associations studied in classical epidemiology—those of behavioural risk factors and obesity phenotype [7]—or even beyond the major genes that play a deciding role in monogenic obesity, such as is leptin deficiency for example, we were interested in SNPs—primary genetic information and variants where the

genetic predisposition could be discovered—and their role in developing common obesity [8]. No single SNP will cause a complex trait; however, in a gene x environment interaction, a combination of variants exposed to what is often called 'obesogenic' environment will increase the relative risk that an individual develops the trait.

We propose that the extent to which this process is mediated by associations between particular SNPs and behavioural risk factors for overweight or obesity determines opportunities for novel, personalised preventive interventions. In the informational abundance of over 10 million human SNPs that are currently listed in publicly accessible data bases [9], we aimed to identify SNPs that might be of interest for this postulate—that is which are linked with both obesity and behavioural risk factors.

1. Material and methods

We searched CDSR, MEDLINE, INSPEC, CC, and CCTR for surveys in any language that examined the associations between any SNPs and behaviours implicated in the aetiology of human obesity, namely physical activity, smoking, diet, and alcohol consumption. Two researchers checked all abstracts for eligibility and we included in this report only those articles that found significant associations between identified SNPs and one or more behaviours of interest. Animal studies were excluded.

2. Results

Our initial search returned 77 abstracts of which 18 were deemed eligible for inclusion (Tables 1 and 2). In one article, where 26 SNPs on the fat mass and obesity associated (FTO) gene were found to be associated with body mass index (BMI), two variants—rs1477196 and rs1861868—were only associated with obesity in people with low levels of physical activity [10]. No association between these two variants and BMI was found among people with above-average physical activity scores.

Another article indicated that alcohol consumption may play a protective mediating role in one variant's impact on glucose metabolism: in men, carriers of 14672C>G in promoter region of hormone-sensitive lipase locus (LIPE) who don't drink alcohol had higher glucose levels than non-carriers, but there were no differences among people who do drink alcohol [11]. In the Oxodeoxyguanosine (OGG1) gene, variant Ser(326)Cys was found to be associated with the risk for breast cancer, but only among moderate alcohol drinkers, while another variant in the same gene—11657A/G—was associated with increased body weight [12].

Variants in the myotubularin-related protein 9 (MTMR9) gene, SLC6A14 gene, and SH2-B gene showed the potential to affect control of appetite [13-15].

A number of articles implicated various SNPs, located on several genes, in changing carriers' response to diet [16-24]. In the TUB gene, for example, AG heterozygote and AA homozygote of the rs2272382 derived less energy from fat, and both were associated with increased energy intake from carbohydrates [16]. Both rs22728133 and rs1528133 were also associated with higher glycaemic load in the diet,

Table 1. SNPs incriminated in the pathophysiology of obesity and linked with risk behaviours (physical activity, alcohol consumption, and control of appetite)

| People | SNP | Phenotype associations |
|--|---|---|
| 704 healthy Old order Amish people [10] | rs1477196 and rs1861868 on fat mass and obesity associated (FTO) gene | Associated with body mass index in people with low physical activity scores (adjusted for age and sex) |
| Population of mostly overweight and obese 373 men and 361 woman [11] | 14672C>G in promoter region of hormone-sensitive lipase locus (LIPE) gene | In women, LIPE 14672G was associated with significantly higher total cholesterol, LDL-cholesterol and apoE; in men, carriers who don't drink alcohol have higher glucose levels than non-carriers |
| 1,058 cases and 1,102 controls [12] | Ser(326)Cys and 11657A/G in Oxodeoxyguanosine (OGG1) gene | Ser(326)Cys associated with breast cancer risk among moderate alcohol drinkers 11657A/G associated with BMI>25 |
| First: 93 cases 469 controls; Second: 564 cases 562 controls; Third: 394 cases 958 controls [13] | rs2293855 in myotubularin-related protein 9 (MTMR9) gene | MTMR9 mRNA levels increased after fasting and decreased after high-fat diet – regulation of hypothalamic neuropeptides and thus possibly control of |

| | | appetite |
|---|---|---|
| Sample of 218 obese Finnish sibling pairs; independent samples of 837 cases and 968 controls [14] | SNP haplotype of the SLC6A14 gene | Evidence of linkage emerged mainly from the obese male sib pairs, suggesting a gender-specific effect for the underlying gene |
| 2455 white female twins [15] | A tagging SNP/tSNP, Ala484Thr (rs7498665) in the region encompassing the human SH2-B gene | Ala484Thr (minor allele frequency 0.38) was associated with serum leptin, total fat, waist circumference, and body weight |

which was higher than glycaemic load among the wild types. Concerning the APO gene [18], among people with APOA5-1131T (major allele) the BMI increased with higher fat intake; however, in APOA5-1131C (minor allele) no increase was seen in BMI with increased fat consumption. Carriers of APOA5-1131C minor allele had a lower risk for overweight and obesity, but not when fat intake was low.

UCP-3 was exposed as an anti-thrifty gene that dissipates energy as heat and prevents obesity [19], while variants in the adiponectin gene had an impact on insulin resistance [21]. In the initial report of the RIVAGE study, some SNPs showed interactions with the metabolic response to diet (through ApoE and LDL-cholesterol and triacylglycerols, apoA-IV and LDL cholesterol, MTP and LDL-cholesterol, intestinal fatty acid-binding protein, and triacylglycerols) [22].

Lastly, ethnic specific and region specific responses, possibly related to diet, were shown to be mediated by several SNPs in the human integrin beta 2 subunit (ITGB2) gene, the diacylglycerol acyltransferase (DGAT) gene, as well as thrifty genes FABP", MTP, CAL10, beta 3AR, apo-E, UCP2, UCP3-p, PPARgama2, and LEPR [26-28].

3. Conclusion

We have identified in the literature a number of SNPs that are associated with increased risk for overweight or obesity and also with behavioural risk factors for these traits. These and most probably many other SNPs hold promise for future design of personalised interventions for prevention of cardiovascular diseases.

Acknowledgements. This research was supported by the European Commission (Project Reference: 224176) and by the Croatian Ministry of Science, Education and Sport (Project No: 108-1080135-0264).

Table 2. SNPs incriminated in the pathophysiology of obesity and linked with risk behaviours (diet)

| People | SNP | Phenotype associations |
|--|---|--|
| 1680 middle-aged Dutch women [16] | rs2272382, rs227283, and rs1528133 in the TUB gene | Eating behaviour associated with body composition and macronutrient intake |
| 451 obese participants [17] | P129T polymorphism in fatty acid amide hydrolase (member of the endocannabinoid (ECS) system) | After six weeks of low fat diet, carriers had a significantly greater decrease in total cholesterol and triglycerides, compared with wild type |
| 1,073 men and 1,207 women in the Framingham offspring study [18] | APOA5-1131T>C polymorphism (present in 13% of the studied population) | Modulates the effect of fat intake on BMI and risk for overweight or obesity |
| 214 overweight women from Korea [19] | Haplotype 1 (ht1) (CGTACC) on the uncoupling protein 3 (UCP-3) gene | After one month of low-energy diet, associated with greater reduction in body weight, BMI, body fat mass; but not with body fat free mass |
| 453 overweight women from Korea [20] | A-3826G, A-1766G, and Ala64Thr (G+1068A) on UCP-1 gene | After one month of very low calorie diet, ht3[GAG] associated with faster reduction in waist-to-hip ratio and |

| | | |
|---|---|---|
| | | body fat mass |
| 249 non-diabetic overweight or obese people from Korea[21] | 276G>T at adiponectin (ADIPOQ) gene | Modifies response to low calorie diet |
| 300 patients randomised to two diet groups over 3 to 12 months [22] | Various SNPs on several genes (see text) | Interactions with metabolic response to Mediterranean/low fat diet or Western type diet |
| 30 men and 29 women [23] | -11377 C>G at the adiponectin gene | C/C homozygous men had a greater decrease in the steady-state plasma glucose concentrations when changing from SFA-rich to MUFA-rich diet |
| 458 overweight women [24] | 10 polymorphisms in uncoupling protein UCP-2 and UCP-3 | Modified response to a one-month very-low calorie diet regimen |
| 651 people of Japanese ethnicity (274 Hawaiian Americans and 377 native Japanese people) [25] | rs235326 in the gene encoding human integrin beta 2 subunit (ITGB2) | In Hawaiian Americans (whose diet has become “westernised”): compared with C carriers, TT homozygotes were 3.29-times more likely to be obese; no such association was found among people living in Japan |
| 1,357 obese adults | 79-bp T-to-C on the | Not associated with |

| | | |
|---|--|---|
| and children from France [26] | 3' region of the diacylglycerol acyltransferase (DGAT) encoding gene | obesity-related phenotypes in this study, although a positive association has been reported in Turkish women |
| People living in affluent societies in several parts of Asia and Pacific islands [27] | Thrifty SNPs encoding FABP", MTP, CAL10, beta 3AR, apo-E, UCP2, UCP3-p, PPARgama2 and LEPR | Differences in these SNPs between Mongoloids and Caucasoids may have been caused by natural selection depending on the types of agricultures practised in different regions and consequently diet |

References

- [1] AHA. *The Americas*. Available from: <http://www.americanheart.org/presenter.jhtml?identifier=2575>.
- [2] *Total CVD mortality in Europe*. Available from: <http://www.heartstats.org/datapage.asp?id=754>.
- [3] WHO. Available from: <http://www.who.int/dietphysicalactivity/publications/facts/cvd/en/>.
- [4] AHA guidelines for primary prevention of cardiovascular disease and stroke: 2002 update, *Circulation* **106** (2002), 388–91.
- [5] AHA/ACC guidelines for secondary prevention for patients with coronary and other atherosclerotic vascular disease: 2006 update, *Circulation* **113** (2006), 2363–72.
- [6] M. Cargill, D. Altshuler, J. Ireland, P. Sklar, K. Ardlie, N. Patil, et al, Characterization of single-nucleotide polymorphisms in coding regions of human genes, *Nat genet* **22** (1999), 231–8.
- [7] S. Musić Milanović, A. Ivičević Uhernik, K. Fišter, Health behaviour factors associated with obesity in adult population in Croatia, *Coll Antropol* 2009 in press.
- [8] K. Clement, T.I. Sorensen, *Obesity: genomics and postgenomics*, Informa HealthCare, 2007.
- [9] D.L. Wheeler, T. Barrett, D.A. Benson, S.H. Bryant, K. Canese, D.M. Church, et al, Database resources of the National Center for Biotechnology Information, *Nucleic Acids Res* **33** (2005) D39–D45.
- [10] E. Rampersaud, B.D. Mitchell, T.I. Pollin, M. Fu, H. Shen, J.R. O'Connell, et al, Physical activity and the association of common FTO gene variants with body mass index and obesity, *Arch Intern Med* **168** (2008), 1791–7.
- [11] L. Qi, H. Shen, I. Larson, J.R. Barnard, E.J. Schaefer, J.M. Ordovas, Genetic variation at the hormone sensitive lipase: gender-

specific association with plasma lipid and glucose concentrations, *Clin Genet* **65** (2004), 93–100.

- [12] P. Rossner, M.B. Terry, M.D. Gammon, F.F. Zhang, S.L. Teitelbaum, S.M. Eng, et al, OGG1 polymorphisms and breast cancer risk, *Cancer Epidemiol Biomarkers Prev* **15** (2006), 811–5.
- [13] T. Yanagiya, A. Tanabe, A. Iida, S. Saito, A. Sekine, A. Takahashi, et al, Association of single-nucleotide polymorphisms in MTMR9 gene with obesity, *Human Molecular Genetics* **16** (2007), 3017–26.
- [14] E. Suviolahti, L.J. Oksanen, M. Ohman, R.M. Cantor, M. Ridderstrale, T. Tuomi, et al, The SLC6A14 gene shows evidence of association with obesity, *J Clin Invest* **112** (2003), 1762–72.
- [15] Y. Jamshidi, H. Snieder, D. Ge, T.D. Spector, S.D. O'Dell, The SH2B gene is associated with serum leptin and body fat in normal female twins, *Obesity (Silver Spring)* **15** (2007), 5–9.
- [16] J.V. van Vliet-Ostaptchouk, N.C. Onland-Moret, R. Shiri-Sverdlov, P.J. van Gorp, A. Custers, P.H. Peeters, et al, Polymorphisms of the *TUB* gene are associated with body composition and eating behavior in middle-aged women, *Plos ONE* **3** (2008), e1405.
- [17] J. Aberle, I. Fedderwitz, N. Klages, E. George, F.U. Beil, Genetic variation in two proteins of the endocannabinoid system and their influence on body mass index and metabolism under low fat diet, *Horm Metab Res* **39** (2007), 395–7.
- [18] D. Corella, C.Q. Lai, S. Demissie, L.A. Cupples, A.K. Manning, K.L. Tucker, et al, APOA5 gene variation modulates the effects of dietary fat intake on body mass index and obesity risk in the Framingham Heart Study, *J Mol Med* **82** (2007), 119–28.
- [19] M.H. Cha, H.D. Shin, K.S. Kim, B.H. Lee, Y. Yoon, The effects of uncoupling protein 3 haplotypes on obesity phenotypes and very

low-energy diet-induced changes among overweight Korean female subjects, *Metabolism* **55** (2006), 578–86.

- [20] H.D. Shin, K.S. Kim, M.H. Cha, Y. Yoon, The effects of UCP-1 polymorphisms on obesity phenotypes among Korean female subjects, *Biochem Biophys Res Commun* **335** (2005), 624–30.
- [21] M.J. Shin, Y. Jang, S.J. Koh, J.S. Chae, O.Y. Kim, J.E. Lee, et al, The association of SNP276G>T at adiponectin gene with circulating adiponectin and insulin resistance in response to mild weight loss, *Int J Obes (Lond)* **30** (2006), 1702–8.
- [22] S. Vincent, R. Planells, C. Defoort, M.C. Bernard, M. Gerber, J. Prudhomme, P. Vague, et al, Genetic polymorphisms and lipoprotein responses to diets, *Proc Nutr Soc* **61** (2002), 427–34.
- [23] P. Pérez-Martínez, J. López-Miranda, C. Cruz-Teno, J. Delgado-Lista, Y. Jiménez-Gómez, J.M. Fernández, et al, Adiponectin gene variants are associated with insulin sensitivity in response to dietary fat consumption in Caucasian men, *J Nutr* **138** (2008), 1609–14.
- [24] Y. Yoon, B.L. Park, M.H. Cha, K.S. Kim, H.S. Cheong, Y.H. Choi, et al, Effects of genetic polymorphisms of UCP2 and UCP3 on very low calorie diet-induced body fat reduction in Korean female subjects, *Biochem Biophys Res Commun* **359** (2007), 451–6.
- [25] T. Awaya, Y. Yokosaki, K. Yamane, H. Usui, N. Kohno, A. Eboshida, Gene-environment association of an ITGB2 sequence variant with obesity in ethnic Japanese, *Obesity (Silver Spring)* **16** (2008), 1463–6.
- [26] S.K. Coudreau, P. Tounian, G. Bonhomme, P. Froguel, J.P. Girardet, B. Guy-Grand, et al, Role of the DGAT gene C79T single-nucleotide polymorphism in French obese subjects, *Obes Res* **11** (2003), 1163–7.
- [27] Y. Kagawa, Y. Yanagisawa, K. Hasegawa, H. Suzuki, K. Yasuda, H. Kudo, et al, Single nucleotide polymorphisms of thrifty genes for energy metabolism: evolutionary origins and prospects for

intervention to prevent obesity-related diseases, *Biochem Biophys Res Commun* **295** (2002), 207–22.