

Prediction of stature based on radiographic measurements of cadaver long bones: a study of the Croatian population

Petrovečki, Vedrana; Mayer, Davor; Šlaus, Mario; Strinović, Davor; Škavić, Josip

Source / Izvornik: **Journal of Forensic Sciences, 2007, 52, 547 - 52**

Journal article, Accepted version

Rad u časopisu, Završna verzija rukopisa prihvaćena za objavljivanje (postprint)

<https://doi.org/10.1111/j.1556-4029.2007.00419.x>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:455860>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-06**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)





Središnja medicinska knjižnica

Petrovečki V., Mayer D., Šlaus M., Strinović D., Škavić J. (2007)
***Prediction of stature based on radiographic measurements of cadaver long bones: a study of the Croatian population.* Journal of Forensic Sciences, 52 (3). pp. 547-52. ISSN 0022-1198**

<http://www.wiley.com/bw/journal.asp?ref=0022-1198>

<http://www3.interscience.wiley.com/journal/118519059/home>

<http://dx.doi.org/10.1111/j.1556-4029.2007.00419.x>

<http://medlib.mef.hr/726>

University of Zagreb Medical School Repository

<http://medlib.mef.hr/>

Prediction of Stature Based on Radiographic Measurements of Cadaver Long Bones: A Study of Croatian Population

Vedrana Petrovečki¹, M.D., Ph.D.; Davor Mayer¹, M.D., M.Sc.; Mario Šlaus¹, Ph.D.; Davor Strinović¹, M.D., Ph.D; and Josip Škavić¹, M.D., Ph.D.

¹ Department of Forensic Medicine and Criminology, Zagreb University School of Medicine, Zagreb, Croatia.

Corresponding author:

Vedrana Petrovečki, M.D., Ph.D.

Department of Forensic Medicine and Criminology

Šalata 11

HR-10000 Zagreb, Croatia

Fax: +385 1 4590 221

E-mail: vedranap@mef.hr

This work has been supported in part by Research Grant No. 0108249 from the Ministry of Science, Education and Sports of the Republic of Croatia.

* A part of the study was presented in poster form at the 16th Meeting of the International Association of Forensic Sciences in September 2002, Montpellier, France.

Running title: Prediction of Human Stature

ABSTRACT: We tested a new approach to the stature prediction that could be used in the identification process of human skeletal remains of unknown identity. The stature of 19 female and 21 male adult cadavers was measured within 24 hours after death and considered equal to the living stature. The anteroposterior radiographs of all limbs were taken and the maximum length of the six long bones was measured from radiographs. There was a significant difference in the stature and maximum length of long bones between female and male cadavers ($p < 0.001$ for all). The correlation between the stature and long bone length was best for the humerus in females ($r = 0.792$) and the tibia in males ($r = 0.891$). Regression equations specific to Croatian population were computed separately for each long bone in males and females and proven reliable in predicting the living stature of the individual.

KEYWORDS: forensic science, forensic radiology, body height, long bone length, Croatian population

Stature estimation is an indispensable part of the identification process of human skeletal remains or body parts (1–3). The accuracy of stature estimation depends on the completeness and state of preservation of the remains, which are often poor. It is, therefore, important to have reliable methods for calculation of stature that can be applied to different bones. In this article, we introduce a new approach to stature prediction based on the measurements of long bones by anteroposterior radiographic imaging.

Anthropometric investigations have shown that ratios between the stature, limb length, and long bone dimensions in an individual are constant during the period of growth and remain the same in adulthood, but they vary depending on stature and ancestry (3–5). Furthermore, recent studies have shown that long bone length might be calculated, and consequently used to estimate the stature, even in cases where only a part of the body (6) or part of the bone (7–9) are available for analysis.

The first comprehensive investigation in this field on European skeletons was done by Telkkä (10) in 1950. Trotter and Gleser (2) performed their research on a sample of North American skeletons. In their first study, they measured long bones of European-American and African-American skeletons from the Robert J. Terry Anatomical Skeletal Collection and long bones of the skeletons of military personnel killed during World War II in the Pacific war zone. In the next study, Trotter and Gleser (11) measured long bones from the skeletons of military personnel killed in the Korean War and found clear differences in the body proportions that depended on the ancestry (European-American, African-American or Asian). Their study also showed differences in body proportions of European-Americans of different ancestries (Mexican population versus other European-Americans) as well as differences in body proportions of European-Americans of the same ancestry in the period from World War II to the Korean War. Their conclusions are considered fundamental in this field and still serve as guidelines for further studies.

Recent research in European population has been performed by De Mendonça (12) on a sample of Portuguese population and by Rodinova et al. (13) on a sample of Bulgarian population. Both investigations were conducted on cadavers and measured the length of long bones in “fresh” condition, i.e. immediately after or before autopsy, according to the standard anthropological techniques (14). De Mendonça (12) measured disarticulated right humerus and right femur after having removed the soft tissue, whereas Rodinova et al. (13) measured the lengths of right and left humeri, tibiae, and fibulae without disarticulating the joints or removing the soft tissues.

Muñoz et al. (15) carried out an investigation on a sample of Spanish population. They measured the stature of healthy adults and then took anteroposterior teleradiographs of the right upper and right lower limb. The length of long bones was determined on the basis teleradiographic findings. However, the authors did not follow the standard anthropological techniques (14).

Similar methodology to measure the length of long bones by use of x-ray imaging was applied by Sarajlić (16), who developed regression equations for stature calculation based on the lengths of femur, tibia, and fibula of 50 Bosnian male cadavers. Ross and Konigsberg (17) offered new formulae for estimating the stature of Eastern-European male population based on the known lengths of the humerus, femur, and tibia.

So far, predictive regression equations for the stature have not been established for Croatian population. It was a considerable problem for the identification team assembled by the Croatian Government to perform identification of human remains from the 1991-1995 war in Croatia.

Since it is ethically unacceptable to cause any unnecessary mutilation to the cadaver or irradiate living patients without a valid medical indication, we decided to use non-invasive methods to determine the stature of cadavers. The aim of our study was to introduce a new

approach to stature estimation based on the cadaver body length and radiographically determined length of long bones. On the basis of these measurements, regression equations were created for the purpose of the identification of unknown skeletal remains recovered in Croatia. The accuracy of these equations was compared to the equations used in previous forensic and anthropological investigations in Croatia and equations derived from the data on populations of neighboring geographical regions (10,12,16–19).

Subjects and Methods

Subjects

The study of long bones in relation to stature was carried out at the Department of Forensic Medicine and Criminology, Zagreb University School of Medicine, between November 2000 and August 2003. Measurements had been made on cadavers before they underwent autopsy. Forty-three consecutive cadavers, 20 female and 23 male, were examined during the study period. There was a similar number of cadavers of each sex and each female cadaver was matched by age with a male cadaver (± 5 years). Inclusion criteria were age ≥ 24 years; absence of physical injuries; absence of visible shortening of any of the limbs; absence of bone pathology, bone surgery, and any other pathological process in medical history that might have influenced the length of long bones. The individuals included in the study either died of natural causes at home (n=3) or in Zagreb University Hospital Center (n=32) or committed suicide by hanging (n=8).

After radiography, 3 cadavers (1 female and 2 male) were excluded from the investigation because of the previously existing fracture of at least one of the long bones. The final group consisted of 40 cadavers, 19 female (median age 67 years, range 35-82) and 21 male (median age 60 years, range 31-86). There was no statistically significant difference in age between female and male cadavers ($t = -1.74$, $p = 0.089$).

Control group consisted of 3 male and 2 female skeletons recovered during the identification process of victims of the 1991-1995 war in Croatia and 1 female who died of natural causes in Zagreb University Hospital Center. Data on statures of 3 male war victims were obtained from the Croatian Army medical records. Data on the stature of two female war victims were obtained from their families, whereas the stature of the female cadaver was measured directly. All individuals were aged over 24 years at the time of death. In the control group, postcranial measurements were taken from the right side of the body according to anthropological rules

(14). The length of skeletonized bone was increased by 2 millimeters to achieve comparability with the bone length on radiograph. This value, 2 mm, corresponds to the thickness of joint cartilage, which gradually disappears during the postmortem changes (2).

Methods

All the procedures used in the study were in accordance with the ethical standards of the Helsinki Declaration.

Cadaver stature

In this study, we used a special tool for measuring the stature of cadavers. The tool, which was composed of a measuring panel and two boards, was installed on the x-ray table (Fig. 1). A metal gauge with a scale graduated in millimeters was fixed along the panel. The cadaver was placed in supine position on the panel and the top of the head (bregma) was brought into contact with the fixed board on the cranial end of the panel. The sliding board on the caudal end of the panel was positioned against the soles of cadaver's feet. Cadaver's stature was measured in millimeters by the metal gauge and considered to be the same as the height of the living subject. For females older than 48 years and males older than 46 years, stature value was corrected according to Giles' tables (20) to compensate for the decline in stature due to aging.

Radiography

The cadaver was secured to the panel by self-adhesive tape to ensure immobility during radiography of the large joints of the limbs (21,22). Under control of a fluoroscope, an injection needle was positioned to indicate the center of the joint cavity. The skin under the needle in correct position was marked at the point where the central, nondivergent x-rays passed through the center of the joint cavity. This method allowed us to minimize the magnification due to the conical divergence of x-rays (22). A standard cassette (Ortho Gradual Gevamic Cassette, AGFA, Leverkusen, Germany) with 30x90 cm x-ray film was used for radiography. Both upper limbs with their large joints fitted on a single radiograph

and were displayed together on the same x-ray film (Fig. 2). Joints of lower limbs were x-rayed independently, each leg on a separate radiograph. The distance between the focus and x-ray film during radiography was 100 cm. Anteroposterior radiographs showed the ends of long bones in the position corresponding to the position of long bones on the osteometric board.

Measurement of Long Bone Length from X-ray Images

The maximum lengths of all six long bones were measured according to the standard anthropological techniques (14) directly from the anteroposterior radiographs by use of a ruler with a measurement accuracy of 1 mm. The length of long bones measured from anteroposterior radiographs equals the length of “fresh” long bones with joint cartilage (22).

Statistical Analysis

Normality of numerical data distribution was tested with the Kolmogorov-Smirnov test. Data were presented as means with standard deviation or 10-90% confidence intervals showing the data range. Two-way ANOVA was used to test the effect of two independent variables on numerical data.

Association of numerical parameters was assessed by Pearson's correlation coefficient r . Whenever r was significant, i.e. $r > 0.6$ (23), univariate regression equations were computed according to the equation $y = \beta_0 + \beta_1 x$, where y was a dependent variable, β_0 and β_1 were regression coefficients, i.e. intercept and slope, and x was the independent variable. Values of regression coefficients were presented with standard errors. Regression line on scattergrams was always shown with the 95% confidence interval limits.

Multiple regression equations were computed according to the equation $y = \beta_0 + \beta_i x_i$, where y was a dependent variable, β_0 and β_i were regression coefficients, and x_i were independent variables.

Statistical analysis was performed with MedCalc for Windows (version 7.5, Frank Schoonjans Inc., Mariakerke, Belgium) and SPSS for Windows (version 7.0, SPSS Inc., Chicago, IL, USA).

Only p values <0.05 were considered significant.

Results

The difference in stature between female and male cadavers was significant ($p < 0.001$; Fig. 3). A significant difference between females and males was also found in length of all six long bones ($p < 0.001$ for all bones; Table 1). When the length of right and left long bones of the same cadaver were compared, no difference was found ($p > 0.05$ for all, Table 1). Naturally, the length of each long bone used in further calculations was the average value of the length of the left and right long bone.

Correlation and univariate regression analyses showed that the association between the stature and length of long bones was substantial, given the high and statistically significant values of Pearson's correlation coefficients, ranging from 0.649 (radius) to 0.792 (humerus) in females and from 0.815 (ulna) to 0.891 (tibia) in males (Table 2).

All regression coefficients were statistically significant (Table 2). Examples of regression equations with graphical presentation are given in Figure 3 for calculating female stature from the humerus length, chosen according to the highest correlation coefficient in females (Fig. 4A), and for calculating male stature from the tibia length, chosen according to the highest correlation coefficient in males (Fig. 4B).

Additionally, we compared correlation coefficients for each long bone of female and male cadavers (Table 2), but no significant difference was found (data not shown).

When a multiple stepwise regression model was applied, only the humerus in females and the tibia in males were found to be significant and independent estimators of human stature. The data obtained were similar to those from univariate regression analysis, with values of regression coefficients close to those presented in Table 2.

The actual stature of six control subjects was compared with statures calculated according to regression equations in Table 2 and with statures calculated according to regression equations

published by other seven authors (Fig. 5). The statures according to other studies were calculated as an average of quoted regression factors, used by Rollet (18), or quoted regression equations from all other studies (10,12,16-19) for each long bone. Stature calculated according to the equations from the present study is close to the real stature of control subjects, with the greatest difference being 4 cm, found in one male subject (Fig. 5).

Discussion

The mean statures of the subjects included in our study were in accordance with thereported data for the mean stature of the Croatian population (24). This is of particular importance for this study considering how well the studied sample represents the entire population. The values of body stature obtained in this study are almost identical to the results of Muñoz et al. (15) obtained on a Spanish population sample in 2001, with the mean of 161.2 ± 6.2 cm for females and 175.3 ± 6.8 cm for males. Interestingly, measuring stature in a Portuguese population sample in 2000, De Mendonça estimated the mean stature of females and males at 157.7 cm (range 145-175) and 167.9 cm (range 153-185), respectively (12), suggesting that the Portuguese may be shorter than Spaniards.

For females older than 48 years ($n=17$) and males older than 45 years ($n=17$) in our study, stature was corrected according to Giles' tables (20) to compensate for the decline in stature due to aging. In his work, Giles (20) showed that the process of height decline in men starts after the age of 45, somewhat earlier than in women in whom the decline in height begins after the age of 47, so these ages are considered a turning point for the correction of living stature. The similar practice was used by Sarajlić (16).

The literature also quotes another approach of living stature correction with respect to the postmortem stature. Some studies (10,12) report the difference between the living stature and stature measured after death and that the stature of the cadaver is on average 2 cm taller than the stature of the living person. Trotter and Gleser (2) found that cadavers included in their study were on average 2.5 cm taller compared with their living stature. However, they concluded that a unique correction value could not be applied since the correction depended on the time interval between death and measurement, environmental conditions, and measurement method. Terry (25) described in detail the measurements of stature performed

on donated corpses. Since several days, sometimes even weeks, had elapsed between the time of death and the measurements he performed, Terry concluded that the increase in the stature of cadaver models in such circumstances resulted from the loss of water and muscle tonicity (25). In our study, we did not include this type of correction of body height. The measurements of body height and x-ray imaging of all cadavers in our study were performed within 24 hours of death. As rigor mortis begins in this early postmortem interval and lasts on average 57 ± 14 hours (26), neither the loss of muscle tonicity nor a significant water loss could have had a direct effect on the body height. This is also confirmed by the fact that the measurements of body height of subjects in this study match those of contemporary Croatian population (24). Other authors also consider cadaver length to be the same as the living stature (16).

In our sample, no statistically significant difference was found in the bone length between the left and right limbs in either sex. However, further estimations were made using the average left/right bone length. Other authors also used the average length of paired bones in their studies (2,10,16). Choi et al. (27) observed no discrepancy in the length of bones between the left and right side and performed regression analysis with right bones length only. Some other authors adopted a similar practice, measuring only long bones of the right limbs (12,15).

The present study confirmed a very distinct and statistically significant correlation between the length of all long bones and the stature, which is consistent with findings of the previous studies. Some authors also reported a sex difference in strength of this correlation. Trotter and Gleser (11), for instance, have demonstrated that long bones of European-American female population correlate with stature better than long bones of the European-American male population. On the other hand, long bones of African-American males showed greater correlation with stature than long bones of African-American females (2). Quite similar observation was reported by Muñoz et al. (15).

In our study, correlation coefficients were higher in male than female cadavers for all six long bones, suggesting that long bone length correlates with stature better in male than female population. However, statistical analysis failed to confirm this observation.

In the present study, separate analysis of correlation of each long bone with the stature showed that the stature correlated best with the humerus in females ($r=0.792$) and the tibia in males ($r=0.891$). Telkkä (10) also found the greatest correlation between the stature and the humerus in females and the stature and the fibula in males. Most authors showed that the long bones of the leg correlated with body height better than the long bones of the arm in both sexes (2,3,15,27).

High and statistically significant correlation coefficients for six long bones allowed us to calculate the regression equations separately for females and males. Accordingly, the known maximum length of any of the six long bones allowed for the calculation of body stature. In identification practice, it means that the individual's living stature may be derived from postmortem remains if there is at least one complete long bone (humerus, radius, ulna, femur, tibia or fibula). Thus, the best results are achieved with long bones that correlate best with stature. For our Croatian population sample, these bones were the humerus in females and the tibia in males.

Multiple regression analysis showed that the best results in calculating stature were obtained by including a single bone, the one that was considered to be in best correlation with the body stature. Adding values of any other long bone did not contribute to the accuracy of body height calculation. On the other hand, results of multiple regression analysis from other studies showed that the most reliable estimates of stature were achieved by a combination of either two (2,27) or three long bones (15).

It is a common practice to check on a control group the applicability of equations derived from a population sample for which regression equations have not yet been computed and to

compare them with already existing equations reported in other studies (12,27–29). In addition to the equations shown in Table 2, we also used equations derived from other European population samples (10,12,16–18), which are commonly used in investigations by forensic and anthropological experts in Croatia, as well as the best known and most frequently used equations for predicting stature derived from the series of European-American population samples (19). Because the maximum lengths of long bones were measured according to anthropological techniques (14), the obtained values could be inserted into equations from the above-mentioned studies. According to Brogdon (22), the length of long bones measured from anteroposterior radiographs equals the length of “fresh” long bones with joint cartilage. Therefore, the long bone lengths measured from anteroposterior radiographs of the female cadaver in our control group were directly inserted into both the equations from this study and equations from the De Mendonça’s study (12). To use these measurements with other equations derived from skeletons (10,17–19) it was necessary to subtract 2 mm, i.e. the thickness of joint cartilage lost during postmortem decomposition (2). Other control subjects in our study were skeletons, so it was necessary to add 2 mm before inserting the lengths of their long bones in our equations and equations from the studies of De Mendonça (12) and Sarajlić (16).

We showed that living statures of control subjects, estimated by equations from this study, had the maximum deviations of ± 4 cm. Similar deviations were reported by Trotter (19). When the equations of other authors were used, the deviations were even larger, with a maximum of ± 7 cm.

It must be pointed out that some of the studies used for comparison of the present results were published more than 100 years ago and that the correlation coefficients between the long bone lengths and the stature are characteristic for particular population. Secular growth of the human population, together with the differences in the correlation between the stature and

long bone length between ancestries, requires continuous updating and development of population-specific methods of stature estimation. The equations from our study are a useful tool for estimating the living stature. We are aware that our relatively small sample size could be a limitation of our study. However, taking into consideration the above factors, these equations could be used for more accurate stature estimation of unknown human remains recovered in Croatia. Future research could enhance the relevance of the present results.

These new regression equations for estimation of the living stature were derived from the contemporary Croatian population and they substantially contribute to the identification process of the victims of 1991-1995 war in Croatia – the task that has become everyday forensic practice in Croatia.

Acknowledgement

We thank Professor Mladen Petrovečki for statistical analysis of the study data and useful advice during the preparation of this manuscript.

References

1. Mall G, Hubig M, Büttner A, Kuznik J, Penning R, Graw M. Sex determination and estimation of stature from the long bones of the arm. *Forensic Sci Int* 2001;117: 23-30.
2. Trotter M, Gleser GC. Estimation of stature from long bones of American whites and negroes. *Am J Phys Anthropol* 1952;10:463-514.
3. Duyar I, Pelin C. Body height estimation based on tibia length in different stature groups. *Am J Phys Anthropol* 2003;122:23-7.
4. Liharžik FP. *La loi de la croissance et la structure de l'homme*. Vienne: De l'imprimerie impériale royale de la cour et de l'état, 1862;7-12.
5. Aldegheri R, Agostini S. A chart of anthropometric values. *J Bone Joint Surg Br* 1993;75:86-8.
6. Ozaslan A, Iscan MY, Ozaslan I, Tugcu H, Koc S. Estimation of stature from body parts. *Forensic Sci Int* 2003;132:40-5.
7. Badkur P, Nath S. Use of regression analysis in reconstruction of maximum bone length and living stature from fragmentary measures of the ulna. *Forensic Sci Int* 1990;45:15-25.
8. Holland TD. Estimation of adult stature from fragmentary tibias. *J Forensic Sci* 1992;37:1223-9.
9. Prasad R, Vettivel S, Jeyaseelan L, Isaac B, Chandi G. Reconstruction of femur length from markers of its proximal end. *Clin Anat* 1996;9:28-33.
10. Telkkä A. On the prediction of human stature from the long bones. *Acta Anatomica* 1950;9:103-17.
11. Trotter M, Gleser GC. A re-evaluation of estimation of stature based on measurements of stature taken during life and of long bones after death. *Am J Phys Anthropol* 1958;16:79-123.

12. De Mendonça MC. Estimation of height from the length of long bones in Portuguese adult population. *Am J Phys Anthropol* 2000;112:39-48.
13. Radoinova D, Tenekedjiev K, Yordanov Y. Stature estimation from long bone lengths in Bulgarians. *Homo* 2002;52:221-32.
14. Bass WM. *Human osteology: a laboratory and field manual*. 4th ed. Columbia: Missouri Archaeological Society; 1995.
15. Muñoz JI, Liñares-Iglesias M, Suárez-Peñaranda JM, Mayo M, Miguéns X, Rodríguez-Calvo MS, et al. Stature estimation from radiographically determined long bone length in a Spanish population sample. *J Forensic Sci* 2001;46:363-6.
16. Sarajlić N, Cihlarž Z, Klonowski EE, Selak I. Stature estimation for Bosnian male population. *Bosn J Basic Med Sci* 2006;6:62-7.
17. Ross AH, Konigsberg LW. New formulae for estimating stature in the Balkans. *J Forensic Sci* 2002;47:165-7.
18. Krogman WM, Iscan MY. *The human skeleton in forensic medicine*. 2nd ed. Springfield: Charles C Thomas, 1986.
19. Trotter M. Estimation of stature from intact long limb bones. In: Stewart TD, editor. *Personal identification in mass disasters*. Washington: National Museum of Natural History, Smithsonian Institution, 1970;71-83.
20. Giles E. Corrections for age in estimating older adults' stature from long bones. *J Forensic Sci* 1991;36:898-901.
21. Bešenski N, Škegro N. Radiografska tehnika skeleta [Radiographic technique of the skeleton, in Croatian]. 2nd ed. Zagreb: Školska knjiga, 1990;9-141.
22. Brogdon BG. *Forensic radiology*. Boca Raton: CRC Press, 1998.
23. Dawson-Saunders B, Trapp RG. *Basic and clinical biostatistics*. London – Singapore: Prentice-Hall Int. Inc., 2001.

24. Mustajbegović J. Vrijednosti ventilacijskih funkcija pluća poljoprivrednih radnika. [Ventilatory capacity values in farmers, in Croatian]. Dissertation. Zagreb: Zagreb University School of Medicine, 1992;48-50.
25. Terry RJ. On measuring and photographing the cadaver. *Am J Phys Anthropol* 1940;26:433-47.
26. Madea B, Krompecher T, Knight B. Muscle and tissue changes after death. In: Knight B, editor. *The estimation of the time since death in early postmortem period*. London: Edward Arnold, 1995;138-67.
27. Choi BY, Chae YM, Chung IH, Kang HS. Correlation between the postmortem stature and the dried limb-bone lengths of Korean adult males. *Yonsei Medical Journal* 1997;38:79-85.
28. Kate BR, Mujumdar RD. Stature estimation from femur and humerus by regression and autometry. *Acta Anatomica* 1976;94:311-20.
29. Pelin IC, Duyar I. Estimating stature from tibia length: a comparison of methods. *J Forensic Sci* 2003;48:708-12.

Additional information and reprint requests:

Vedrana Petrovečki, M.D., Ph.D.

Department of Forensic Medicine and Criminology

School of Medicine University of Zagreb

Šalata 11, HR-10000 Zagreb

Croatia

E-mail: vedranap@mef.hr

TABLE 1–Long bone lengths (mean±S.D., in cm) in 40 cadavers according to sex and body side.

Bone	Females (n=19)		Males (n=21)	
	left	right	left	right
Humerus	30.8±1.8	30.9±1.7	33.3±1.4	33.5±1.3
Ulna	24.8±1.1	25.0±1.2	26.9±1.2	27.0±1.2
Radius	22.7±1.1	22.9±1.2	24.8±1.1	25.0±1.1
Femur	43.3±2.3	43.3±2.4	46.7±2.0	46.6±2.1
Tibia	35.3±2.2	35.4±2.1	37.9±2.0	37.9±2.1
Fibula	34.9±1.7	34.8±1.7	37.8±1.7	37.7±1.8

Two-way ANOVA results used to calculate effect of sex ($p < 0.001$ for all bones), side of the body ($p > 0.05$ for all bones) and sex and side interaction ($p > 0.05$ for all) to the length of bones.

TABLE 2–Correlation and univariate regression equations comparing stature of subjects (dependent variable, y) with length of long bones

(independent variable, x). Each regression equation is computed as $y = \beta_0 + \beta_1 x$, separately for males and females.*

Bone	Females (n=19)				Males (n=21)			
	r	$\beta_0 \pm \text{S.E.}(\beta_0)$	$\beta_1 \pm \text{S.E.}(\beta_1)$	S.E.(y)	r	$\beta_0 \pm \text{S.E.}(\beta_0)$	$\beta_1 \pm \text{S.E.}(\beta_1)$	S.E.(y)
Humerus	0.792	69.3±17.4	3.0±0.6	4.15	0.823	82.1±14.5	2.7±0.4	2.52
Ulna	0.700	56.6±26.2	4.2±1.0	4.92	0.815	94.7±12.9	2.9±0.5	2.56
Radius	0.649	76.6±24.4	3.8±1.1	5.21	0.821	93.3±12.9	3.2±0.5	2.52
Femur	0.764	68.3±19.3	2.2±0.4	4.42	0.870	86.9±11.3	1.9±0.2	2.17
Tibia	0.701	83.3±19.6	2.2±0.6	4.91	0.891	102.2±8.4	1.9±0.2	2.00
Fibula	0.747	57.5±22.7	3.0±0.7	4.59	0.890	90.7±9.8	2.2±0.3	2.03

*Statistical note: r – Pearson’s correlation coefficient; β_0 – first regression coefficient (intercept) with standard error S.E.(β_0); β_1 – second regression coefficient (slope) with standard error S.E.(β_1); S.E.(y) – standard error of estimation. All correlation coefficients were found significant at $p \leq 0.001$ except $p=0.003$ for the radius in females. All regression coefficients were found significant at $p < 0.05$. (No comparison between females and males was performed.)

Figure legend

FIG. 1—Tool for measuring the length of cadavers placed on the x-ray table. Measuring panel from transparent perspex with attached (a) metal gauge with scale graduated in millimeters; (b) fixed board on the cranial end; and (c) sliding board on the caudal end.



FIG. 2—Hand radiograph (subject from the study, female, age 31). Both hands, left (L) and right (R), are on the same radiograph. The ends of long bones are visible on anteroposterior radiographs of each large joint in order to enable the measurement of long bone lengths. Injection needles determine the centers of the joint cavities. Original film size: 30x90 cm.



FIG. 3–Box-and-whisker plot with a scattergram of stature in females (n=19, mean±S.D. = 162.4±6.7 cm, 10-90% = 153.1-170.4 cm) and males (n=21, mean±S.D. = 173.6±4.3 cm, 10-90% = 168.2-178.2 cm). The intersexual difference in stature was found significant ($t = -6.35$, $p < 0.001$).




FIG. 4–Regression lines with the 95% confidence level comparing stature of the subjects (dependent variable, y) with the length of their humerus (A, for females) or tibia (B, for males) as an independent variable (x). Regression equations are copied from Table 2.





FIG. 5—Comparison of body stature with stature calculated according to methods from six different, previously published studies. Control subjects are designated by letters A – F.

