

# Chronic respiratory symptoms in Croatian Adriatic island metapopulations

---

Žuškin, Eugenija; Smolej-Narančić, Nina; Škarić-Jurić, Tatjana; Barbalić, Maja; Rudan, Pavao; Kujundžić-Tiljak, Mirjana; Pucarín-Cvetković, Jasna; Schachter, E. Neil

Source / Izvornik: **Croatian Medical Journal, 2006, 47, 627 - 634**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:751335>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-08-17**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



## Chronic Respiratory Symptoms in Croatian Adriatic Island Metapopulations

Eugenija Žuškin<sup>1</sup>, Nina Smolej Narančić<sup>2</sup>, Tatjana Škarić-Jurić<sup>2</sup>, Maja Barbalić<sup>2</sup>, Pavao Rudan<sup>2</sup>, Mirjana Kujundžić-Tiljak<sup>3</sup>, Jasna Pucarín-Cvetković<sup>1</sup>, E. Neil Schachter<sup>4</sup>

<sup>1</sup>Department of Environmental and Occupational Health, Andrija Štampar School of Public Health, Zagreb University School of Medicine, Zagreb, Croatia

<sup>2</sup>Institute for Anthropological Research, Zagreb, Croatia

<sup>3</sup>Department of Medical Statistics, Epidemiology, and Medical Informatics, Andrija Štampar School of Public Health, Zagreb University School of Medicine, Zagreb, Croatia

<sup>4</sup>The Mount Sinai School of Medicine, New York, NY, USA

> **Correspondence to:**

Eugenija Žuškin  
Andrija Štampar School of  
Public Health  
Zagreb University School of  
Medicine  
Rockefellerova 4  
10000 Zagreb, Croatia  
ezuskin@snz.hr

> **Received:** January 25, 2006

> **Accepted:** April 21, 2006

> **Croat Med J. 2006;47:627-34**

**Aim** To investigate the prevalence of chronic respiratory symptoms in 9 metapopulations on Adriatic islands in Croatia, and the relationship between respiratory symptoms and individual genetic background.

**Methods** We obtained random sample of 1001 adult inhabitants of 9 Adriatic island villages in Croatia, that also included immigrants to these villages. European Union respiratory health questionnaire and World Health Organization non-communicable diseases questionnaire were used. Personal genetic histories were reconstructed, based on the two-generation ancestral pedigrees. Bivariate and multivariate methods were used in the analysis.

**Results** Women reported the occurrence of acute dyspnea ( $P=0.017$ ), cough ( $P=0.002$ ), and asthma ( $P=0.002$ ) more often than men. Gender was the strongest predictor for acute and/or chronic cough (odds ratio [OR], 1.69; 95% confidence interval [CI], 1.23-2.33) and asthma (OR, 2.00; 95% CI, 1.00-4.01), whereas smoking was the strongest risk factor for acute and chronic dyspnea (OR, 1.90; 95% CI, 1.21-2.99) and airway narrowing (OR, 1.84; 95% CI, 1.18-2.87). Residence on the northern islands increased the odds of allergy, whereas the highest odds ratio of 3.20 was associated with the interaction of northern residence and immigrant background. Genetic background was a significant predictor only for the occurrence of allergy symptoms.

**Conclusion** Differences in respiratory findings among the island inhabitants were often associated with smoking prevalence. Interaction of residence on northern Adriatic islands and immigrant background proved to be the strongest predictor for the occurrence of allergy symptoms. This study indicated that environmental factors played a very important role in the occurrence of respiratory symptoms.

Nonspecific respiratory symptoms in general population have recently received increased attention, as respiratory diseases are associated with a significant proportion of chronic morbidity (1,2). A complex interaction between genetic background and exposure to multiple environmental stimuli is widely recognized in the etiology of many lung diseases (3). The role of genetics is being intensively investigated, using predominately genome screens and association studies. These studies have identified regions of the genome which are linked with the phenotypes of asthma and atopy (4,5), or with the occurrence of chronic obstructive pulmonary disease (COPD) (6). The most important environmental risk factor for the development of respiratory diseases is tobacco smoke inhalation, either among smokers, or non-smokers through passive smoking (7,8). Air pollution, such as exposure to organic or inorganic dusts, fumes, or gasses, is another highly prevalent environmental risk factor.

The most common chronic respiratory diseases are asthma and chronic obstructive pulmonary disease, two distinct inflammatory disorders, with different pathogenesis, clinical courses, and distinct treatment strategies (9,10). Asthma is a very common disorder, affecting people of all ages. Chronic obstructive pulmonary disease is a progressive, slow onset debilitating disease of the airways, characterized by a gradual loss of lung function and reduction in life quality (11-13). The term COPD includes chronic bronchitis, emphysema, or a combination of these conditions. It is a highly prevalent smoking-related condition, thus representing a significant economic burden to the health care system (14).

The aim of this study was to analyze the self-reported prevalence of asthma and other chronic respiratory symptoms in the population of 9 villages on Adriatic islands in Croatia, that also included immigrants to these villages. We also examined geographical variation among the studied villages in the prevalence of these symptoms and the possible relation of respiratory symptoms to

the genetic background inferred on the basis of two-generation pedigrees.


## Subjects and methods

### Setting

The “1001 Dalmatians” research program was launched with an aim to study multiple small isolated populations of Croatian islands that share a similar environment and lifestyle (15). The data were collected during 2002 and 2003. The details of this program are described elsewhere (16).

### Subjects

This study included subjects from the “1001 Dalmatians” research program. Random samples of adult inhabitants were collected from 9 villages on the Adriatic islands of Rab (villages Banjol, Barbat, Lopar, Rab, and Supetarska Draga), Vis (Vis and Komiža), Lastovo, and Mljet, in Croatia (Figure 1). Sampling was based on computerized randomization of the most complete and accessible population registries in the villages. The total sample consisted of 454 men and 547 women. Apart from the village of residence, the sample was divided into sub-samples according to age, gender, and smoking habits. According to age, subjects were divided into three groups:  $\leq 35$  years, 36 to 64 years, and  $\geq 65$  years, and into



**Figure 1.** Geographic location of the investigated islands of Rab, Vis, Lastovo and Mljet. Immigrants in the islands originate from the mainland Croatia.

two categories according to smoking habits: non-smokers (including former smokers) and smokers. Subjects were also divided into two regional subgroups, according to geographical position of the studied island in the Adriatic Sea. Northern subgroup included subjects from the villages on the island of Rab, whereas southern subgroup included subjects from the villages on the islands of Vis, Lastovo, and Mljet.

Subjects were then classified into categories according to personal genetic history, based on the two-generation ancestral pedigrees. We classified each subject into one of the 3 categories based on their presumed individual genome-wide heterozygosity. Category "Inbred" included examinees with all four grandparents born in the studied village. Category "Outbred" included examinees with 1 to 4 grandparents born in different small villages on each island. Category "Immigrants" included examinees who had between 1 and 4 grandparents born on the mainland and (majority of examinees were also born on the mainland). Although personal genetic history is not, in strict terms, an ordered categorical variable, we could expect differences among the categories in mean individual genome-wide heterozygosity. The lowest mean value of individual genome-wide heterozygosity would be expected in the category "Inbred", which included individuals who were recently inbred, followed by the categories "Outbred" and "Immigrants."

The ethical approval for this research was obtained from appropriate research ethics committees in Croatia and Scotland. Informed written consent was obtained from all participants in the study.

### **Respiratory symptoms**

Data on respiratory symptoms and diseases were obtained from the European Union respiratory health questionnaire (17), and World Health Organization (WHO) non-communicable diseases questionnaire (18). All subjects were asked about acute dyspnea, chronic dyspnea, airway

narrowing, acute and/or chronic cough, asthma, asthma therapy, and allergy symptoms. Data on smoking habits included the number of cigarettes smoked per day and the number of smoking years, expressed as pack-years. Pack years were counted as the number of cigarettes smoked daily, multiplied by years of smoking and divided by 20. Acute dyspnea was defined as occasional dyspnea during the last 12 months, whereas chronic dyspnea was defined as regular dyspnea in the last 12 months. Airway narrowing was recorded if the subject had the feeling of airway narrowing in the morning during the last 12 months. Acute cough was defined as occasional cough during the last 12 months and chronic cough as daily cough in the morning or during the day over the last two years. Subjects were considered to have asthma if they had difficulties in breathing and/or asthma attack with reduction in lung function. Asthma therapy was defined as taking of asthma medications on a regular basis. Allergic symptoms related to environmental (atmospheric or food) pollutants were also recorded.

### **Statistical analysis**

Differences in frequencies of respiratory symptoms and diseases between subgroups were tested using  $\chi^2$  test. Age differences were tested by univariate analysis of variance (ANOVA). Association of respiratory symptoms and diseases with multiple risk factor variables was assessed by logistic regression analysis. Analyses were performed by SPSS 10.0 statistical package for Windows (SPSS Inc., Chicago, IL, USA), with statistical significance set at  $P < 0.05$ .

### **Results**

In the initial analysis, based on descriptive methods, we investigated gender differences in respiratory symptoms. Age and prevalence of smoking were not significantly different between genders, but significant differences in smoking habits were recorded. Men smoked longer and more

**Table 1.** Basic characteristics (mean  $\pm$  standard deviation, range) of 1001 subjects from Croatian Adriatic villages in 2002-2003

Characteristics	Gender		P*
	men (n=454)	women (n=547)	
Age (y)	55.0 $\pm$ 15.5 (18-87)	55.0 $\pm$ 14.1 (19-88)	0.945
Current tobacco smokers prevalence (n; %)	103 (22.7)	121 (22.1)	0.445†
age (y)	47.9 $\pm$ 15.2 (18-79)	46.2 $\pm$ 11.7 (20-73)	0.343
smoking intensity (cigarettes/d)	21.8 $\pm$ 12.8 (1-50)	16.9 $\pm$ 10.8 (1-60)	0.002
smoking duration (y)	25.0 $\pm$ 13.9 (1-62)	21.3 $\pm$ 8.9 (3-45)	0.016
pack-years‡	27.4 $\pm$ 21.8 (0.10-90.00)	18.8 $\pm$ 16.3 (0.25-90.00)	0.001

\*ANOVA.

†Fischer exact test.

‡Pack years were counted as the number of cigarettes smoked daily multiplied by years of smoking and divided by 20.

**Table 2.** Respiratory symptoms and diseases by gender in 1001 subjects from Croatian Adriatic villages in 2002-2003

Respiratory symptoms	No. (%) of subjects		P*
	men (n=454)	women (n=547)	
Acute dyspnea	108 (23.8)	164 (30.0)	0.032
Chronic dyspnea	56 (12.3)	70 (12.8)	0.849
Airway narrowing	54 (11.9)	79 (14.4)	0.262
Acute and/or chronic cough	75 (16.5)	133 (24.3)	0.003
Asthma	5 (1.1)	23 (4.2)	0.003
Asthma therapy	11 (2.4)	25 (4.6)	0.087
Allergy symptoms	66 (14.5)	98 (17.9)	0.170

\*Fischer exact test.

intensively and had higher pack-years count (Table 1). The prevalence of respiratory symptoms also exhibited a certain difference between genders (Table 2). Women had higher prevalence of all respiratory symptoms, with significantly higher findings for acute dyspnea, acute and/or chronic cough, and asthma. Among age groups, there were no significant differences in respiratory symptoms (Table 3).

Smoking was significantly associated with the occurrence of acute and chronic dyspnea in men smokers, whereas other respiratory symptoms and diseases did not show significant association with smoking (Table 4).

In the northern group of islands there was significantly higher prevalence of all respiratory symptoms and diseases, except for asthma and asthma therapy (Table 5). We recorded significant differences in personal genetic histories background for the total and regional samples. Slightly more than a half of the total sample was inbred, with higher percentage of inbred subjects in the southern group. Immigrants were equally represented in both groups (Table 6).

Multivariate logistic regression analysis was initially performed, using a full model which included gender, age, smoking status, region of residence, and genetic background (Table 7). Gender was the strongest predictor for acute and/or chronic cough and asthma, and smoking was the strongest risk factor for acute and chronic dyspnea and airway narrowing. Region was a significant risk factor for all the symptoms, except for asthma, whereas genetic background was a significant predictor only for allergy symptoms with odds ratio of 1.97 associated with being immigrant.

In the extended analysis, personal genetic histories-by-region interaction was included with

**Table 3.** Respiratory symptoms and diseases by gender and age groups in 1001 subjects from Croatian Adriatic villages in 2002-2003

Respiratory symptoms	No. (%) of subjects per age group (y)							
	men (n=454)				women (n=547)			
	<35 (n=61)	36-64 (n=240)	>65 (n=153)	P*	<35 (n=53)	36-64 (n=331)	>65 (n=163)	P*
Acute dyspnea	8 (13.1)	59 (24.6)	41 (26.8)	0.096	15 (28.3)	91 (27.5)	58 (35.6)	0.175
Chronic dyspnea	5 (8.2)	31 (12.9)	20 (13.1)	0.572	5 (9.4)	43 (13.0)	22 (13.5)	0.733
Airway narrowing	7 (11.5)	31 (12.9)	16 (10.5)	0.759	5 (9.4)	45 (13.6)	29 (17.8)	0.253
Acute and/or chronic cough	9 (14.8)	37 (15.4)	29 (19.0)	0.604	9 (17.0)	85 (25.7)	39 (23.9)	0.387
Asthma	0	4 (1.7)	1 (0.7)	0.435	1 (1.9)	14 (4.2)	8 (4.9)	0.635
Asthma therapy	0	7 (2.9)	4 (2.6)	0.409	1 (1.9)	16 (4.8)	8 (4.9)	0.616
Allergy symptoms	11 (18.0)	34 (14.2)	21 (13.7)	0.702	8 (15.1)	68 (20.5)	22 (13.5)	0.135

\* $\chi^2$  test.

**Table 4.** Respiratory symptoms and diseases by gender and smoking habits in 1001 subjects from Croatian Adriatic villages in 2002-2003

Respiratory symptoms	No. (%) of subjects					
	men (n = 454)			women (n = 547)		
	non-smokers (n = 351)	smokers (n = 103)	P*	non-smokers (n = 426)	smokers (n = 121)	P*
Acute dyspnea	74 (21.1)	34 (33.0)	0.017	124 (29.1)	40 (33.1)	0.432
Chronic dyspnea	37 (10.5)	19 (18.4)	0.040	53 (12.4)	17 (14.0)	0.645
Airway narrowing	37 (10.5)	17 (16.5)	0.119	58 (13.6)	21 (17.4)	0.307
Acute and/or chronic cough	55 (15.7)	20 (19.4%)	0.368	106 (24.9)	27 (22.3)	0.632
Asthma	4 (1.1)	1 (1.0)	0.999	21 (4.9)	2 (1.7)	0.130
Asthma therapy	10 (2.8)	1 (1.0)	0.469	22 (5.2)	3 (2.5)	0.322
Allergy symptoms	55 (15.7)	11 (10.7)	0.265	76 (17.8)	22 (18.2)	0.999

\*Fisher exact test.

**Table 5.** Respiratory symptoms and diseases by region in 1001 subjects from Croatian Adriatic villages in 2002-2003

Respiratory symptoms	No. (%) of subjects per region		
	South (n = 467)	North (n = 534)	P*
Acute dyspnea	110 (23.6)	162 (30.3)	0.019
Chronic dyspnea	46 (9.9)	80 (15.0)	0.017
Airway narrowing	51 (10.9)	82 (15.4)	0.040
Acute and/or chronic cough	79 (16.9)	129 (24.2)	0.005
Asthma and asthma therapy	18 (3.9)	24 (4.5)	0.639
Allergy symptoms	55 (11.8)	109 (20.4)	0.001

\*Fisher exact test.

gender, age, and smoking as predictor variables. Among the studied respiratory symptoms and diseases, the interaction was significant only for allergy symptoms (OR, 1.32; 95% CI, 1.17-1.48). Stratification to four interaction items (region-by-genetic background) indicated that the highest odds ratios were recorded in immigrant group from the northern islands (OR, 3.20), followed by outbred group from the northern islands (OR, 1.73) and immigrant groups from the southern islands (OR, 1.12). Outbred group from the southern islands exhibited the lowest odds ratio (OR, 0.74), relative to all inbred examinees.

## Discussion

This study describes respiratory findings in the population of selected Croatian island villages. Interestingly, men and women exhibited the same prevalence of smoking, although men smoked longer and more frequently than women, which resulted in significantly higher pack-years count.

However, women exhibited higher prevalence of acute dyspnea, acute/chronic cough,

**Table 6.** Genetic background in the total sample and by region of residence in 1001 subjects from Croatian Adriatic villages in 2002-2003

Samples	No. (%) of subjects		
	inbred	outbred	immigrants
Total sample (n = 1001)	529 (52.8)	382 (38.2)	90 (9.0)
Regional samples:			
South (n = 467)	267 (57.2)	153 (32.8)	47 (10.0)
North (n = 534)	262 (49.1)	229 (42.9)	43 (8.0)

\*P=0.004,  $\chi^2$ test.

and asthma. Watson et al (19) demonstrated that gender plays an important role in the prevalence of COPD. In their study, despite significantly lower number of pack-years of smoking, women were more likely to report severe dyspnea than men. Similarly, French et al (20) demonstrated that women with chronic cough were more inclined to seek medical help than men, because their health-related quality of life was more adversely affected and because they were more apt to suffer from physical complaints. Hardie et al (21), who studied respiratory symptoms in men and women, showed that these symptoms occurred half as likely in men as in women, indicating that female gender was a predictor of dyspnea, current asthma, and chronic bronchitis.

Higher prevalence of respiratory symptoms was found in the villages on northern islands. Benner et al (22) investigated the prevalence of asthma and wheezing in the population of two different climate areas of Saudi Arabia and reported major differences between the coastal and inland areas. Similar study was carried out in Germany (23), reporting the significant differences among elderly inhabitants in the preva-



**Table 7.** Multivariate screening for predictor variables for respiratory symptoms by logistic regression analysis

Independent variable	Predictors (OR, 95% CI)*					
	acute dyspnea	chronic dyspnea	airway narrowing	acute and/or chronic cough	asthma and its therapy	allergy symptoms
Gender:						
male (referent)	1.00	1.00	1.00	1.00	1.00	1.00
female	1.42 (1.06-1.90) P=0.018	1.04 (0.71-1.53) P=0.846	1.31 (0.90-1.91) P=0.161	1.69 (1.23-2.33) P=0.001	2.00 (1.00-4.01) P=0.049	1.24 (0.87-1.76) P=0.231
age (y)	1.03 (1.01-1.04) P<0.001	1.02 (1.01-1.04) P=0.004	1.02 (1.00-1.03) P=0.026	1.01 (1.00-1.02) P=0.052	1.01 (0.99-1.04) P=0.269	0.99 (0.98-1.01) P=0.287
Smoking:						
nonsmokers (referent)	1.00	1.00	1.00	1.00	1.00	1.00
smokers	1.98 (1.40-2.83) P<0.001	1.90 (1.21-2.99) P=0.006	1.84 (1.18-2.87) P=0.007	1.23 (0.83-1.81) P=0.310	0.51 (0.19-1.36) P=0.180	0.82 (0.53-1.27) P=0.370
Region:						
South (referent)	1.00	1.00	1.00	1.00	1.00	1.00
North	1.62 (1.20-2.17) P=0.001	1.80 (1.21-2.68) P=0.004	1.63 (1.11-2.40) P=0.012	1.69 (1.22-2.32) P=0.001	1.18 (0.62-2.22) P=0.616	1.89 (1.32-2.71) P<0.001
Genetic background:						
inbred (referent)	1.00	1.00	1.00	1.00	1.00	1.00
outbred	0.92 (0.68-1.25) P=0.606	0.95 (0.63-1.42) P=0.785	0.93 (0.63-1.38) P=0.723	0.86 (0.62-1.20) P=0.374	1.37 (0.69-2.72) P=0.364	1.23 (0.86-1.77) P=0.264
immigrants	1.02 (0.60-1.73) P=0.940	1.48 (0.77-2.85) P=0.238	0.80 (0.39-1.65) P=0.546	0.80 (0.44-1.43) P=0.446	2.41 (0.94-6.18) P=0.067	1.97 (1.13-3.43) P=0.017

\*Abbreviations: OR – odds ratio; CI – confidence intervals.

lence of allergic symptoms. The authors suggested that non-specific causes related to different lifestyles played an important role in the pathogenesis of respiratory symptoms.

A possible explanation to the North-South difference would be indoor pollution. Various risk factors from the indoor environment, such as heating and cooking installations, use of a humidifier, treatment air system or ventilation systems, wooden floor, oil fume contamination, smoke, pets, and wall-to-wall carpets have been associated with the occurrence of respiratory findings (24,25). Women exposed to bio-fuel smoke and passive cigarette smoke suffered more from respiratory illnesses and had decreased pulmonary functions (26-28). Some of these factors might be variably expressed on southern and northern islands, thus causing higher prevalence of respiratory findings among inhabitants of the northern islands.

Multivariable analysis revealed that gender was significantly associated with the prevalence of acute dyspnea, and the prevalence of acute and chronic cough. Borderline significance was recorded in case of asthma and asthma therapy. Age as a predictor variable was significantly as-

sociated with acute and chronic dyspnea, and airway narrowing, whereas borderline significance was recorded for acute and chronic cough. Smoking was associated only with acute and chronic dyspnea and airway narrowing. However, the highest number of significant findings was obtained with region as a predictor variable: all findings except asthma yielded significant results. Finally, personal genetic histories were significantly associated only with the occurrence of allergy symptoms, and borderline significance was recorded with the occurrence of asthma. This is in line with the familiar occurrence of various allergy and asthma symptoms, often occurring in the subtle interaction between genetic and environmental factors (3,4).

A combination of residence on the northern islands and the immigrant background in a stepwise regression model was associated with the highest risk of respiratory allergy. This was an unexpected finding, as inbreeding is expected to act as a depressor of various physiological functions (29). A possible explanation could be related to previous environmental exposure of the immigrant group to polluted air (30). Unfortunately, it is well beyond the scope of this

study to investigate previous environmental exposure.

The main shortcoming of the present study was the use of self-reported data on the respiratory symptoms, as there might have been a various degree of recall bias. Personal genetic histories might not have been correct in all cases, as genealogical data are prone to various levels of uncertainty.

Nevertheless, this study suggests an association between gender and the respiratory symptoms and pathology. There are geographical differences in the occurrence of respiratory findings between northern and southern Adriatic islands. Various dependent variables were significantly associated with the occurrence of respiratory symptoms and diseases, but genetic effects could be only linked to the presence of allergy symptoms. Personal genetic histories were not significantly associated with occurrence of other respiratory symptoms.

#### Acknowledgments

The study was partially supported by the grants from the Ministry of Science, Education, and Sports of the Republic of Croatia (grants No. 0108330, 0196001, and 0196005), and the grants from the British Council, the Wellcome Trust, the Royal Society, and Medical Research Council. The authors are grateful to a large number of individuals (medical students of the Zagreb University School of Medicine, Croatia; local general practitioners and nurses in study populations; employees of several other Croatian institutions, including the University of Rijeka and Split, Croatia; Croatian Institute of Public Health; Institutes of Public Health in Split and Dubrovnik, Croatia; and the Institute for Anthropological Research in Zagreb, Croatia) for their individual help in planning and carrying out the field work related to the project. There are no conflicts of interest related to this manuscript.

#### References

- 1 Lebowitz MD, Bronnimann S, Camilli AE. Asthmatic risk factors and bronchial reactivity in non-diagnosed asthmatic adults. *Eur J Epidemiol.* 1995;11:541-8. [Medline:8549728](#)
- 2 Smith MH, Anderson RT, Bradham DD, Longino CF Jr. Rural and urban differences in mortality among Americans 55 years and older: analysis of the National Longitudinal Mortality Study. *J Rural Health.* 1995;11:274-86. [Medline:10153687](#)
- 3 Kleeberger SR, Peden D. Gene-environment interactions in asthma and other respiratory diseases. *Annu Rev Med.* 2005;56:383-400. [Medline:15660518](#)
- 4 Blumenthal MN. The role of genetics in the development of asthma and atopy. *Curr Opin Allergy Clin Immunol.* 2005;5:141-5. [Medline:15764904](#)
- 5 Malerba G, Pignatti PF. A review of asthma genetics: gene expression studies and recent candidates. *J Appl Genet.* 2005;46:93-104. [Medline:15741670](#)
- 6 Ning W, Li CJ, Kaminski N, Feghali-Bostwick CA, Alber SM, Di YP, et al. Comprehensive gene expression profiles reveal pathways related to the pathogenesis of chronic obstructive pulmonary disease. *Proc Natl Acad Sci U S A.* 2004;101:14895-900. [Medline:15469929](#)
- 7 Anderson HR. Smoking habits and their relationship to chronic lung disease in a tropical environment in Papua New Guinea. *Bull Physiopathol Respir (Nancy).* 1974;10:619-33. [Medline:4441755](#)
- 8 Dautzenberg B. Tobacco related diseases [in French]. *Rev Prat.* 2004;54:1877-82. [Medline:15655911](#)
- 9 Mugusi F, Edwards R, Hayes L, Unwin N, Mbanja JC, Whiting D, et al. Prevalence of wheeze and self-reported asthma and asthma care in an urban and rural area of Tanzania and Cameroon. *Trop Doct.* 2004;34:209-14. [Medline:15510944](#)
- 10 Booker R. Chronic obstructive pulmonary disease: importance of diagnosis. *Br J Nurs.* 2004;13:871-5. [Medline:15284652](#)
- 11 Booker R. Chronic obstructive pulmonary disease: non-pharmacological approaches. *Br J Nurs.* 2005;14:14-8. [Medline:15750483](#)
- 12 Doherty DE. The pathophysiology of airway dysfunction. *Am J Med.* 2004;117 Suppl 12A:11S-23S. [Medline:15693639](#)
- 13 James AL, Palmer LJ, Kicic E, Maxwell PS, Lagan SE, Ryan GF, et al. Decline in lung function in the Busselton Health Study: the effects of asthma and cigarette smoking. *Am J Respir Crit Care Med.* 2005;171:109-14. [Medline:15486340](#)
- 14 Ramsey SD, Sullivan SD. Chronic obstructive pulmonary disease: is there a case for early intervention? *Am J Med.* 2004;117 Suppl 12A:3S-10S. [Medline:15693638](#)
- 15 Rudan I, Campbell H, Rudan P. Genetic epidemiological studies of eastern Adriatic "island isolates, Croatia: objective and strategies. *Coll Antropol.* 1999;23:531-46. [Medline:10646227](#)
- 16 Rudan I, Biloglav Z, Vorko-Jovic A, Kujundzic-Tiljak M, Stevanovic R, Ropac D, et al. Effects of inbreeding, endogamy, genetic admixture, and outbreeding on human health: a "1001 Dalmatians" study. *Croat Med J.* 2006;47:601-10. [Medline:16909458](#)
- 17 Bellia V, Pistelli F, Giannini D, Scichilone N, Catalano F, Spatafora M, et al. Questionnaires, spirometry and PEF monitoring in epidemiological studies on elderly respiratory patients. *Eur Respir J Suppl.* 2003;40:21s-7s. [Medline:12762570](#)
- 18 Pardell H, Roure E, Drygas W, Morava E, Nussel E, Puska P, et al. East-west differences in reported preventive practices. A comparative study of six European areas of the WHO-CINDI programme. *Eur J Public Health.* 2001;11:393-6. [Medline:11766479](#)
- 19 Watson L, Vestbo J, Postma DS, Decramer M, Rennard S, Kiri VA, et al. Gender differences in the management and experience of Chronic Obstructive Pulmonary Disease. *Respir Med.* 2004;98:1207-13. [Medline:15588042](#)
- 20 French CT, Fletcher KE, Irwin RS. A comparison of



- gender differences in health-related quality of life in acute and chronic coughers. *Chest*. 2005;127:1991-8. [Medline:15947311](#)
- 21 Hardie JA, Vollmer WM, Buist AS, Bakke P, Morkve O. Respiratory symptoms and obstructive pulmonary disease in a population aged over 70 years. *Respir Med*. 2005;99:186-95. [Medline:15715185](#)
- 22 Bener A, al-Jawadi TQ, Ozkaragoz F, Anderson JA. Prevalence of asthma and wheeze in two different climatic areas of Saudi Arabia. *Indian J Chest Dis Allied Sci*. 1993;35:9-15. [Medline:8225434](#)
- 23 Filipiak B, Heinrich J, Nowak D, Wichmann HE. The distribution in specific IgE and the prevalence of allergic symptoms in 25-64-years old inhabitants of an eastern and a western German city – results from Augsburg and Erfurt. *Eur J Epidemiol*. 2001;17:77-84. [Medline:11523581](#)
- 24 Clarisse B, Seta N, El Hasnaoui A, Terrioux P, Momas I. How to describe indoor environment in epidemiological studies dealing with respiratory diseases [in French]. *Rev Epidemiol Sante Publique*. 2002;50:179-200. [Medline:12011735](#)
- 25 Yue W, Pan XC, Ding J. Risk factors for allergic asthma in a case-control study among adults [in Chinese]. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2004;25:706-9. [Medline:15553398](#)
- 26 Dutt D, Srinivasa DK, Rotti SB, Sahai A, Konar D. Effect of indoor air pollution on the respiratory system of women using different fuels for cooking in an urban slum of Pondicherry. *Natl Med J India*. 1996;9:113-7. [Medline:8664820](#)
- 27 Pope CA III, Xu X. Passive cigarette smoke, coal heating, and respiratory symptoms of nonsmoking women in China. *Environ Health Perspect*. 1993;101:314-6. [Medline:8275988](#)
- 28 Ng TP, Hui KP, Tan WC. Respiratory symptoms and lung function effects of domestic exposure to tobacco smoke and cooking by gas in non-smoking women in Singapore. *J Epidemiol Community Health*. 1993;47:454-8. [Medline:8120499](#)
- 29 Rudan I, Rudan D, Campbell H, Carothers A, Wright A, Smolej-Narancic N, et al. Inbreeding and risk of late onset complex disease. *J Med Genet*. 2003;40:925-32. [Medline:14684692](#)
- 30 Voisin C. Urban air pollution and respiratory health: the stakes involved and methods of approach [in French]. *Bull Acad Natl Med*. 1989;173:437-45. [Medline:2790531](#)

ANNOUNCEMENT

**Symposium: Recent Advances in Endemic Nephropathy  
The Role of Toxins in an Environmental Disease**

October 20-22, 2006, Zagreb, Croatia

Organized by:

University of Zagreb, School of Medicine, Zagreb, Croatia;  
State University of New York at Stony Brook, NY, USA;  
Croatian Center for Endemic Nephropathy, Slavonski Brod, Croatia

Chairs:

Nada Čikeš, Arthur P. Grollman, Bojan Jelaković

Main topics:

The etiology of endemic nephropathy and its associated urothelial cancer,  
with an emphasis on connecting basic and clinical science with public health.  
Recent studies on the role of nephrotoxic and carcinogenic agents (ochratoxin A and aristolochic acid).

Official language:

English

Contact:

Dr. Živka Dika  
Department for Nephrology and Arterial Hypertension  
University Hospital Center Zagreb  
Kišpatičeva 12, Zagreb, Croatia  
[zivkadika@yahoo.co.uk](mailto:zivkadika@yahoo.co.uk)  
[www.endemic-nephropathy.mef.hr](http://www.endemic-nephropathy.mef.hr)