

Venous thromboembolism in Croatia - Croatian Cooperative Group for Hematologic Diseases (CROHEM) study

Pulanić, Dražen; Gverić-Krečak, Velka; Nemet-Lojan, Zlatka; Holik, Hrvoje; Coha, Božena; Babok-Flegarić, Renata; Komljenović, Mili; Knežević, Dijana; Petrovečki, Mladen; Zupančić Šalek, Silva; ...

Source / Izvornik: **Croatian Medical Journal, 2015, 56, 550 - 557**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.3325/cmj.2015.56.550>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:299883>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-02-28**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



Croat Med J. 2015;56:550-7
doi: 10.3325/cmj.2015.56.550

Venous thromboembolism in Croatia – Croatian Cooperative Group for Hematologic Diseases (CROHEM) study

Aim To analyze the incidence and characteristics of venous thromboembolism (VTE) in Croatia.

Methods The Croatian Cooperative Group for Hematologic Diseases conducted an observational non-interventional study in 2011. Medical records of patients with newly diagnosed VTE hospitalized in general hospitals in 4 Croatian counties (Šibenik-Knin, Koprivnica-Križevci, Brod-Posavina, and Varaždin County) were reviewed. According to 2011 Census, the population of these counties comprises 13.1% of the Croatian population.

Results There were 663 patients with VTE; 408 (61.54%) had deep vein thrombosis, 219 (33.03%) had pulmonary embolism, and 36 (5.43%) had both conditions. Median age was 71 years, 290 (43.7%) were men and 373 (56.3%) women. Secondary VTE was found in 57.3% of participants, idiopathic VTE in 42.7%, and recurrent VTE in 11.9%. There were no differences between patients with secondary VTE and patients with idiopathic VTE in disease recurrence and sex. The most frequent causes of secondary VTE were cancer (40.8%), and trauma, surgery, and immobilization (38.2%), while 42.9% patients with secondary VTE had ≥ 2 causes. There were 8.9% patients ≤ 45 years; 3.3% with idiopathic or recurrent VTE. Seventy patients (10.6%) died, more of whom had secondary (81.4%) than idiopathic (18.6%) VTE ($P < 0.001$), and in 50.0% VTE was the main cause of death. Estimated incidence of VTE in Croatia was 1.185 per 1000 people.

Conclusion Characteristics of VTE in Croatia are similar to those reported in large international studies. Improved thromboprophylaxis during the presence of risk factors for secondary VTE might substantially lower the VTE burden.

Dražen Pulanić^{1,2,3}, Velka Gverić-Krečak⁴, Zlatka Nemet-Lojan⁵, Hrvoje Holik⁶, Božena Coha⁶, Renata Babok-Flegarić⁷, Mili Komljenović⁸, Dijana Knežević⁶, Mladen Petrovečki^{9,10}, Silva Zupančić Šalek^{1,2,3}, Boris Labar^{1,2}, Damir Nemet^{1,2,3}

¹Division of Hematology, Department of Internal Medicine, University Hospital Center Zagreb, Zagreb, Croatia

²Medical School University of Zagreb, Zagreb, Croatia

³Faculty of Medicine Osijek, J.J. Strossmayer University of Osijek, Osijek, Croatia

⁴Division of Internal Medicine, General Hospital Šibenik, Šibenik, Croatia

⁵Division of Internal Medicine, General Hospital "Dr. Tomislav Bardek," Koprivnica, Croatia

⁶Division of Internal Medicine, General Hospital "Dr. Josip Benčević," Slavonski Brod, Croatia

⁷Division of Internal Medicine, General Hospital Varaždin, Varaždin, Croatia

⁸Division of Internal Medicine, General Hospital "Hrvatski Ponos," Knin, Croatia

⁹Department of Clinical Laboratory Diagnosis, Dubrava University Hospital, Zagreb, Croatia

¹⁰Department of Medical Informatics, Rijeka University School of Medicine, Rijeka, Croatia

Received: October 17, 2015

Accepted: December 21, 2015

Correspondence to:

Dražen Pulanić
Kišpatičeva 12
10000 Zagreb, Croatia
dpulanic@yahoo.com

Venous thromboembolism (VTE), including deep venous thrombosis (DVT) and pulmonary embolism (PE), is a major health problem in the world, associated with significant morbidity and mortality (1-9). Incidence rates for VTE mostly vary from 1 to 2 in 1000 individuals per year (1-5,7,9). PE, the most serious manifestation of VTE, has a mortality rate of more than 15% in the first 3 months after diagnosis, with short-term survival of less than 60% (10,11). Cohen et al estimated that the number of VTE-related deaths across the European Union (EU) was 543 454 per year, which was more than double the number of combined deaths in EU due to AIDS, breast and prostate cancer, and traffic accidents (8).

VTE in survivors is associated with several chronic consequences of the disease that can severely impair the patients' quality of life, including post-thrombotic syndrome (PTS) and pulmonary hypertension (PH), as well as recurrent VTE. PTS affects at least one-third of patients after DVT (8,12-15) and PH affects 4%-5% of patients after PE (8,16,17). VTE has significant incidence of recurrence: 10.1% at 6 months, 12.9% after 1 year, and 30.4% after 10 years (18).

Total VTE-related costs to health care system are enormous. For example, the total cost of VTE to the UK National Health Service in 1993 was £235-£257 million (€349-€382 million), and the combined direct and indirect costs in 2004/2005 were approximately £640 million (€950 million), and are even higher when PTS is taken into account (8,19,20).

VTE is a multifactorial disease, resulting from a complex interaction of genetic and acquired factors. Although some studies estimated that genetics was related to up to 60% of the risk of VTE (including FV Leiden and prothrombin G20210A mutations, deficiencies of protein C, S and antithrombin, and elevations of some procoagulant factors) (21), there is also a large number of acquired risk factors for VTE such as immobilization, surgery, trauma, cancer, pregnancy and puerperium, oral contraceptives, autoimmune diseases, and other disorders (1-8,21).

In spite of the importance of VTE, there is not enough data on its incidence and characteristics in transitional countries. Also, although several studies analyzed the epidemiology of VTE in different study settings (1-9), there is still not much information on conditions present at the diagnosis of thrombosis, comparing idiopathic and secondary (provoked) VTE. Therefore, the Croatian Cooperative Group for Hematologic Diseases (CROHEM) analyzed the incidence and characteristics of idiopathic and secondary newly di-

agnosed VTE in Croatia in 2011, the year of the most recent national population census.

MATERIAL AND METHODS

CROHEM conducted an observational non interventional study of patients with newly diagnosed VTE hospitalized in major general hospitals in four Croatian counties: Šibenik-Knin (General Hospital Šibenik, Šibenik, and General Hospital "Hrvatski Ponos," Knin), Koprivnica-Križevci (General Hospital "Dr. Tomislav Bardek," Koprivnica), Brod-Posavina (General Hospital "Dr. Josip Benčević" in Slavonski Brod and Nova Gradiška), and Varaždin County (General Hospital Varaždin) from January 1 until December 31 2011. The study was approved by the medical ethics committee of each hospital.

From patients' hospital records we collected information on age, sex, date of incident, category of VTE (DVT, PE, or both); characteristics of VTE (idiopathic or secondary, first or recurrent); and comorbidities. Diagnosis and treatment of VTE was performed in each hospital according to standard clinical practice and local policy and was not part of this study analysis.

VTE was classified as idiopathic or secondary. It was classified as secondary VTE (DVT, PE, or both) if it was associated with major trauma, surgery, or marked immobility; sepsis; cancer active within 6 months of VTE event; implanted venous catheter; pregnancy or puerperium at the time of VTE; usage of some prothrombotic medicaments at the time of VTE (ie, oral contraceptives, hormonal replacement therapy); autoimmune or other prothrombotic diseases (ie, antiphospholipid syndrome, systemic lupus erythematosus, inflammatory bowel disease, neurologic diseases with paralysis); or if occurred after >6 h long distance airplane flight. VTE was classified as idiopathic when none of these precipitating factors were registered. VTE was also classified as the first or recurrent. VTE was considered recurrent if prior VTE events were registered in the medical record. Patients who died during hospitalization when VTE was diagnosed were recorded and the main cause of death was categorized as VTE-related or related to some other main cause.

According to the 2011 Population Census, the Republic of Croatia had 4 284 889 inhabitants (48.2% men), and the population was on average 41.7 years old (men 39.9, women 43.4 years), which places it among the oldest nations in Europe (22). Of the counties included in this study, one was from the southern (Dalmatian) part

of Croatia (Šibenik-Knin County, with 109 375 inhabitants, 53 596 [49%] men, average age 44.1 years), two were from the northern part of Croatia (Varaždin County, with 175 951 inhabitants, 85 645 [48.7%] men, average age 41.2 years, and Koprivnica-Križevci County, with 115 584 inhabitants, 55 964 [48.4%] men, average age 41.6 years), and one from the eastern (Slavonian) part of Croatia (Brod-Posavina County, with 158 574 inhabitants, 77 115 [48.6%] men, average age 40.6 years). These 4 counties have 559 485 inhabitants (272 320 (48.7%) men), which represents 13.1% of Croatian population (22).

Statistical analysis

Categorical data are presented with absolute (N) and relative (%) frequencies and compared using χ^2 -test, and numerical data are presented with median and range values and compared using *t* test. Only $P < 0.050$ was considered significant. Annual incidence per 1000 inhabitants was calculated using the number of recorded new VTE cases as the numerator and the population at risk in the 4 counties as the denominator, according to the 2011 Census (22). Data were analyzed using MedCalc program (MedCalc Statistical Software version 15.6.1, MedCalc Software bvba, Ostend, Belgium).

RESULTS

In 2011, there were 663 new cases of VTE in 4 Croatian counties: 408 (61.54%) of them had DVT, 219 (33.03%) had PE,

and 36 (5.43%) had both conditions (Table 1). The majority of patients were elderly people (median age 71 years, range 13-97 years), and there were more women (N=373, 56.3%) (Table 1). Female patients were significantly older (median 74, range 21-97 years) than male (median 65, range 13-90 years; $P < 0.001$). Patients with PE or with both PE and DVT were significantly older than patients with DVT alone (median and range; 74 [21-97] vs 69 [13-92] years, $P = 0.006$).

Recurrent VTE was diagnosed in 79 (11.9%) patients (Table 2). There were more patients with secondary (N=380, 57.3%) than with idiopathic VTE (N=283, 42.7%) (Tables 2-3). Patients with idiopathic VTE were significantly older than those with secondary VTE (median and range; 74 [26-92] vs 69 [13-97] years, $P = 0.001$). There were no differences between patients with secondary and patients with idiopathic VTE in disease recurrence and sex.

The most frequent causes of secondary VTE were cancer (40.8% of secondary VTE) and trauma, surgery, and immobilization (38.2% of secondary VTE). Other causes included sepsis, other diseases (eg, autoimmune diseases, inflammatory bowel diseases, neurological diseases with paralysis), central venous catheter, drugs, pregnancy and puerperium, and >6 h airplane flight. Patients with trauma, surgery, or immobilization were significantly younger than other patients with VTE (median and range; 67 [13-90] vs 72 [21-97] years, $P = 0.003$). Many of the 380 cases of secondary VTE (N=163, 42.9% of secondary VTE) had more than one underlying condition (Table 3).

TABLE 1. Diagnosis, sex, and age of patients with newly diagnosed venous thromboembolism (VTE) in 4 Croatian counties in 2011*

County	Total VTE (N)	Diagnosis VTE N (% VTE)			Sex		Age in years (median, range)
		DVT	PE	DVT+PE	male N (%)	female N (%)	
Šibenik-Knin	151	99 (65.56)	43 (28.46)	9 (5.96)	64 (42.4)	87 (57.62)	72 (13-92)
Koprivnica-Križevci	87	54 (62.07)	26 (29.89)	7 (8.05)	39 (44.8)	48 (55.2)	69 (26-90)
Varaždin	298	159 (53.36)	128 (42.95)	11 (3.69)	125 (41.9)	173 (58.1)	73 (21-97)
Brod-Posavina	127	96 (75.59)	22 (17.32)	9 (7.09)	62 (48.82)	65 (51.2)	67 (33-88)
Total	663	408 (61.54)	219 (33.03)	36 (5.43)	290 (43.7)	373 (56.3)	71 (13-97)

*PE – pulmonary embolism; DVT – deep venous thrombosis.

TABLE 2. Idiopathic and recurrent venous thromboembolism (VTE), death, and cause of death of patients with newly diagnosed VTE in 4 Croatian counties in 2011

County	Idiopathic N (%)	Recurrent N (%)	Death N (% VTE)	Reason of death N (% of death)	
				VTE	other diseases
Šibenik-Knin	74 (49.0)	18 (11.9)	11 (7.29)	4 (36.4)	7 (63.6)
Koprivnica-Križevci	40 (46.0)	10 (11.5)	6 (6.9)	4 (66.7)	2 (33.3)
Varaždin	122 (40.9)	39 (13.1)	38 (12.8)	16 (42.1)	22 (57.9)
Brod-Posavina	47 (37.0)	12 (9.5)	15 (11.8)	11 (73.3)	4 (26.7)
Total	283 (42.7)	79 (11.9)	70 (10.6)	35 (50.0)	35 (50.0)

There were 59 (8.9%) VTE patients aged 45 or younger (Table 4). Among them there were significantly more men than women (59.3% vs 40.7%, $P=0.011$), significantly more cases of secondary than of idiopathic VTE (11.6% vs 5.3%, $P=0.005$), and 22 cases (3.3% of all VTE patients) of idiopathic or recurrent VTE. There was no significant difference in VTE incidence between seasons or months of the year (data not shown).

Among all 663 VTE patients, 70 (10.6%) died, 50.0% of them with VTE as the main cause of death (Table 2). There was no difference in sex and no difference in VTE recurrence between patients who died and those who survived. Patients who died were significantly older than those who survived (median and range; 77 [26-90] vs 70 [13-97] years, $P=0.001$). Among patients who died, significantly more patients had secondary (81.4%) than idiopathic (18.6%) VTE ($P<0.001$).

When we compared the counties according to the type of VTE (DVT, PE, or both), Varaždin County had the highest percentage and Brod-Posavina County the lowest percentage of PE ($P<0.001$). Furthermore, Brod-Posavina County had the highest percentage and Varaždin County the lowest percentage of DVT ($P<0.001$, Table 1). There were no significant differences between counties in age and sex of the patients, recurrence of VTE, percentage of idiopathic or secondary VTE, mortality, and cause of death. When we compared the counties according to the cause of secondary VTE, Brod-Posavina County had the highest percentage of trauma, surgery and immobilization (29.1%, $P=0.020$), and Varaždin County the highest percentage of sepsis (9.4%, $P<0.001$).

Varaždin County (1.694 per 1000 persons) and Šibenik-Knin County (1.381 per 1000 persons) had significantly higher incidence of VTE than Brod-Posavina County (0.8 per 1000

TABLE 3. Causes of secondary newly diagnosed venous thromboembolism (VTE) in 4 Croatian counties in 2011

County	Secondary (N, % all VTE in county and total)	Causes of secondary VTE* (N, % of secondary VTE in county and total)		
		cancer	trauma, surgery, immobilization	other†
Šibenik-Knin	77 (51.0)	39 (50.7)	23 (29.9)	28 (36.4)
Koprivnica-Križevci	47 (54.0)	20 (42.6)	24 (51.1)	19 (40.4)
Varaždin	176 (59.06)	68 (38.6)	61 (34.7)	87 (49.4)
Brod-Posavina	80 (63.0)	28 (22.1)	37 (46.3)	45 (56.3)
Total	380 (57.3)	155 (40.8)	145 (38.2)	179 (47.1)

*163 (42.9%) secondary VTE had >1 cause of secondary VTE.

†Other causes of VTE included other diseases (eg, autoimmune diseases, inflammatory bowel diseases, neurological diseases with paralysis), sepsis, central venous catheter, drugs, pregnancy and puerperium, >6 h airplane flight.

TABLE 4. Patients aged 45 years or younger with newly diagnosed venous thromboembolism (VTE) in 4 Croatian counties in 2011

County	Younger ≤45 years (N, % of all VTE in county and total)			
	total	idiopathic	recurrent	idiopathic or recurrent
Šibenik-Knin	16 (10.6)	3 (1.9)	3 (1.9)	6 (3.9)
Koprivnica-Križevci	9 (10.3)	1 (1.15)	2 (2.3)	3 (3.45)
Varaždin	22 (7.4)	8 (2.7)	2 (0.7)	10 (3.4)
Brod-Posavina	12 (9.5)	3 (2.4)	1 (0.8)	3 (2.4)
Total	59 (8.9)	15 (2.3)	8 (1.2)	22 (3.3)

TABLE 5. Incidence of venous thromboembolism (VTE) in 4 Croatian counties and estimated VTE incidence for the entire Croatia in 2011

County	Number of inhabitants in 2011		VTE (N)	Incidence of VTE per 1000 inhabitants
	(N, % of whole Croatia)			
Šibenik-Knin*	109 375 (2.6)		151	1.381
Koprivnica-Križevci	115 584 (2.7)		87	0.753
Varaždin*	175 951 (4.1)		298	1.694
Brod-Posavina	158 575 (3.7)		127	0.800
Total	559 485 (13.1)		663	1.185

*Counties with higher incidence of VTE than other counties, $P<0.001$.

persons) and Koprivnica-Križevci County (0.753 per 1000 persons; $P < 0.001$, Table 5). Estimated annual incidence of VTE in Croatia according to these 4 counties was 1.185 per 1000 people.

DISCUSSION

Our study showed that the estimated incidence of VTE in Croatia in 2011 was 1.185 per 1000 people. This estimate, as well as the characteristics of VTE observed in this study, are similar to the findings of large international studies, although there are also some differences. Age is an important risk factor for VTE. Several studies showed that the incidence of VTE markedly increased with increasing age for both sexes, from fewer than 5 first-time VTE cases per 100 000 per year among children younger than 15 years to 450-600 cases per 100 000 per year among people older than 80 years (1,3,5). Indeed, in our study the majority of patients were older people (median age 71 years). In the past 50 years, the average age in Croatia increased by almost 10 years (from 32.5 in 1961 to 41.7 years in 2011), mostly due to a continuous decrease in fertility and increase in life expectancy (22).

In our study, patients with PE or with both PE and DVT were older than patients with DVT alone. Other studies showed that the elderly more often had PE and had more serious VTE than younger patients, and that in their case massive PE was particularly life-threatening (23).

Although other studies mostly found no consistent differences in the incidence of VTE between men and women (5), our study found more female (56.3%) than male (43.7%) patients with newly diagnosed VTE. In Croatia there are more women (51.8%) than men (48.2%), but there are also different proportions of men and women in different age groups. For example, there are more men in younger age groups and more women in older age groups, starting with the age group 45-49 years (22). In our study, which mostly included elderly patients, female patients were significantly older than male, but in the subgroup of patients aged 45 years or younger there were more male patients, reflecting the age-sex structure in Croatia.

In our study, recurrent VTE was diagnosed in 11.9% of patients. Recurrent VTE has been already recognized as a serious problem. For example, in a prospective cohort study of 355 patients with DVT, Prandoni et al found recurrent VTE in 8.6% of cases after 6 months and in 30.3% of cases after 8 years (12), while Heit et al reported recur-

rent VTE in 10.1% of cases after 6 months, in 12.9% after 1 year, and in 30.4% after 10 years (18).

VTE is a multifactorial disease resulting from a complex interaction between different inherited (genetic) and acquired prothrombotic factors. Inherited thrombophilia is a predisposition to thrombosis and exposes carriers to increased risks for VTE compared with non-carriers. Coen et al showed that the prevalence of FV Leiden and prothrombin G20210A mutations were higher in Croatian patients with VTE than in healthy subjects (24), and Jukic et al confirmed the association of non-O0 blood group genotypes with increased risk of VTE in Croatia (25). However, it is still a matter of intense debate under which circumstances comprehensive and expensive laboratory genetic testing for inherited thrombophilia is useful in clinical practice. Some authors suggest that thrombophilia screening should be considered in patients with VTE aged 45 years or younger, especially in those with idiopathic or recurrent VTE (26). Therefore, we analyzed the patients aged 45 years or younger and found 59 (8.9%) patients in that subgroup, with more cases of secondary than idiopathic VTE, and 22 of them (just 3.3% of all VTE patients) had idiopathic or recurrent VTE, representing possible candidates for genetic thrombophilia testing.

In our study, there were more patients with secondary than with idiopathic VTE. In other studies, the proportion of patients with idiopathic VTE was between 26% and 47% of first-time cases, partly depending on the definition of idiopathic and secondary VTE used in the studies (5). In our study patients with idiopathic VTE were significantly older than those with secondary VTE, although both groups included mostly elderly patients. There was no difference in disease recurrence or sex between patients with secondary and idiopathic VTE. The most frequent cause of secondary VTE was cancer (40.8% of secondary VTE), but 42.9% of cases with secondary VTE had more than one underlying condition associated with VTE.

Cancer is one of the strongest risk factors for VTE and, vice versa, VTE is a frequent complication of malignancy and may be the earliest manifestation of an occult cancer (27). The association of cancer and VTE is due to multiple factors: malignant cells can produce hypercoagulable state through multiple mechanisms, and the risk for VTE is further increased by chemotherapy and other drugs for cancer treatment, the use of central venous catheters, surgery, and prolonged immobility (28,29). These patients require appropriate thromboprophylaxis and adequate long term anticoagulant treatment if VTE occurs (30).

The second most frequent cause of secondary VTE in our study was trauma, surgery, and immobilization (38.2% of secondary VTE). It is known that surgery is related to a 6-fold increased risk of VTE (7,9). The 9th edition of Antithrombotic Therapy and Prevention of Thrombosis Guidelines of the American College of Chest Physicians state that the high incidence of postoperative VTE mandates that thromboprophylaxis should be considered in every surgical patient (31).

Although some studies showed a higher incidence of fatal PE during winter, others did not confirm this finding (5). Similarly, we did not find a significant difference in VTE incidence either between seasons or between the months of the year.

In our study, 70 (10.6%) VTE patients died and in 50% of them VTE was the main cause of death. Patients who died and those who survived did not differ in sex and recurrence of VTE. The literature reports that the risk of dying is highest shortly after the VTE event, and that during the first year after the VTE it gradually approaches that in the general population (7). In our study, patients who died were significantly older than those who survived. Among patients who died, there were more patients with secondary than with idiopathic VTE. In patients with cancer, VTE is the second leading cause of death (32), and cancer was the most frequent underlying condition for secondary VTE in our study.

Diagnosis and treatment of VTE was performed in each hospital according to the standard clinical practice and local policy. Therefore, the decision to perform Color Doppler ultrasound analysis of peripheral veins to exclude DVT in patients with confirmed PE was made by the local physician depending of the clinical presentation and local policy, and was not part of this study. This might explain the finding that 219 (33.03%) patients had PE alone and just 36 (5.43%) patients had confirmed both PE and DVT. However, some studies also reported a small percentage (6%) of patients with both PE and DVT (7) and most clinical studies that did not include autopsy data described the incidence of clinically diagnosed DVT to be approximately twice that of PE (5), which is similar to our finding.

Varaždin County and Šibenik-Knin County had higher incidence of VTE than Brod-Posavina County and Koprivnica-Križevci County. These differences might be explained by the fact that Šibenik-Knin County has the oldest population among the studied counties (average age 44.1 years

and is the second oldest county in Croatia after Lika-Senj County (22). On the other hand, General Hospital Varaždin, the only general hospital in Varaždin County, admits many chronically ill patients with immobility transferred from other regional special hospitals for chronic diseases or from rehabilitation centers, who are at a greater risk for VTE event (personal communication).

The estimated annual incidence of VTE in our study was 1.185 per 1000 people (or 118.5 per 100000 people), which is comparable to other literature data, where VTE incidence rates varied from 1 to 2 per 1000 individuals per year (1-5,7,9). The epidemiology of VTE is always challenging: beside real differences, differences between studies may also arise from differences in study design, patients cohorts, and case definition (7).

This study has some limitations. First, the true incidence of VTE may have been underestimated for several reasons: we reviewed inpatient medical records and did not include VTE cases treated outside of the county general hospitals (for example cases from nursing homes or sudden deaths from community). However, outpatient treatment of newly DVT and especially PE is extremely rare in Croatia. It is also likely that there were fatal and/or non diagnosed VTE events that were not included in this analysis. Another potential limitation is that assessment of associated conditions and classification as idiopathic or secondary VTE were limited by contents of the medical record, and some data were missing for many patients such as smoking, obesity, diet, and family history for VTE.

Strengths of this study include involvement of 4 different counties from different parts of Croatia, representing 13.1% of Croatian population according to the national population Census that was done in the same year as this study, thus representing the most accurate data. We believe that our study made a reliable estimate of the burden of VTE in Croatia, describing its characteristics and associated substantial mortality, which might help policy-makers to develop strategies dealing with this major health problem.

However, further research is needed to estimate VTE incidence in Croatia more accurately, to describe trends in incidence over time, and to implement optimal VTE prevention and management. The literature data showed that the type, duration, and intensity of prophylaxis were frequently inappropriate and suboptimal (33-35). In our study, the most common risk factors for secondary VTE were cancer, trauma, surgery, and immobilization,

and many cases of secondary VTE had more than one underlying condition. Since VTE is a preventable disease and effective prophylaxis is widely available – in addition to the development of new anticoagulants (36) – improved thromboprophylaxis in these settings might substantially lower the incidence and mortality of VTE.

Funding None.

Ethical approval received from the medical ethics committee of each hospital where this study was conducted.

Declaration of authorship The study was initiated by DP and VGK. VGK, ZNL, HH, BC, RBF, MK, and DK contributed to acquisition, analysis, and interpretation of data. DP, MP, SZS, BL, and DN contributed to conception and design of the study with analysis and interpretation of data. MP performed statistical analysis. The manuscript was drafted by DP. All authors revised the work critically for important intellectual content, approved the final version of the manuscript, and are accountable for all aspects of the work.

Competing interests MP is a statistical editor in the *Croatian Medical Journal*. To ensure that any possible conflict of interest relevant to the journal has been addressed, this article was reviewed according to best practice guidelines of international editorial organizations. All authors have completed the Unified Competing Interest form at www.icmje.org/doi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- Anderson FA Jr, Wheeler HB, Goldberg RJ, Hosmer DW, Patwardhan NA, Jovanovic B, et al. A population-based perspective of the hospital incidence and case-fatality rates of deep vein thrombosis and pulmonary embolism. The Worcester DVT Study. *Arch Intern Med*. 1991;151:933-8. [Medline:2025141](#) [doi:10.1001/archinte.1991.00400050081016](#)
- Kniffin WD Jr, Baron JA, Barrett J, Birkmeyer JD, Anderson FA Jr. The epidemiology of diagnosed pulmonary embolism and deep venous thrombosis in the elderly. *Arch Intern Med*. 1994;154:861-6. [Medline:8154949](#) [doi:10.1001/archinte.1994.00420080053005](#)
- Silverstein MD, Heit JA, Mohr DN, Petterson TM, O'Fallon WM, Melton LJ III. Trends in the incidence of deep vein thrombosis and pulmonary embolism: a 25-year population-based cohort study. *Arch Intern Med*. 1998;158:585-93. [Medline:9521222](#) [doi:10.1001/archinte.158.6.585](#)
- Heit JA, Silverstein MD, Mohr DN, Petterson TM, Lohse CM, O'Fallon WM, et al. The epidemiology of venous thromboembolism in the community. *Thromb Haemost*. 2001;86:452-63. [Medline:11487036](#)
- White RH. The epidemiology of venous thromboembolism. *Circulation*. 2003;107 Suppl 1:14-8. [Medline:12814979](#) [doi:10.1161/01.CIR.0000078468.11849.66](#)
- Lijfering WM, Rosendaal FR, Cannegieter SC. Risk factors for venous thrombosis – current understanding from an epidemiological point of view. *Br J Haematol*. 2010;149:824-33. [Medline:20456358](#) [doi:10.1111/j.1365-2141.2010.08206.x](#)
- Naess IA, Christiansen SC, Romundstad P, Cannegieter SC, Rosendaal FR, Hammerström J. Incidence and mortality of venous thrombosis: a population-based study. *J Thromb Haemost*. 2007;5:692-9. [Medline:17367492](#) [doi:10.1111/j.1538-7836.2007.02450.x](#)
- Cohen AT, Agnelli G, Anderson FA, Arcelus JL, Bergqvist D, Brecht JG, et al. Venous thromboembolism (VTE) in Europe. The number of VTE events and associated morbidity and mortality. *Thromb Haemost*. 2007;98:756-64. [Medline:17938798](#)
- Cushman M, Tsai AW, White RH, Heckbert SR, Rosamond WD, Enright P, et al. Deep vein thrombosis and pulmonary embolism in two cohorts: the longitudinal investigation of thromboembolism etiology. *Am J Med*. 2004;117:19-25. [Medline:15210384](#) [doi:10.1016/j.amjmed.2004.01.018](#)
- Heit JA, Silverstein MD, Mohr DN, Petterson TM, O'Fallon WM, Melton LJ III. Predictors of survival after deep vein thrombosis and pulmonary embolism: a population-based, cohort study. *Arch Intern Med*. 1999;159:445-53. [Medline:10074952](#) [doi:10.1001/archinte.159.5.445](#)
- Goldhaber SZ, Visani L, De Rosa M. Acute pulmonary embolism: clinical outcomes in the International Cooperative Pulmonary Embolism Registry (ICOPER). *Lancet*. 1999;353:1386-9. [Medline:10227218](#) [doi:10.1016/S0140-6736\(98\)07534-5](#)
- Prandoni P, Lensing AW, Cogo A, Cuppini S, Villalta S, Carta M, et al. The long term clinical course of acute deep venous thrombosis. *Ann Intern Med*. 1996;125:1-7. [Medline:8644983](#) [doi:10.7326/0003-4819-125-1-199607010-00001](#)
- Asbeutah AM, Riha AZ, Cameron JD, McGrath BP. Five year outcome study of deep vein thrombosis in the lower limbs. *J Vasc Surg*. 2004;40:1184-9. [Medline:15622373](#) [doi:10.1016/j.jvs.2004.10.025](#)
- Roumen-Klappe EM, den Heijer M, Janssen MC, van der Vleuten C, Thien T, Wollersheim H. The post-thrombotic syndrome: incidence and prognostic value of non-invasive venous examinations in a six-year follow-up study. *Thromb Haemost*. 2005;94:825-30. [Medline:16270638](#)
- Wille-Jørgensen P, Jørgensen LN, Crawford M. Asymptomatic postoperative deep vein thrombosis and the development of postthrombotic syndrome. A systematic review and meta-analysis. *Thromb Haemost*. 2005;93:236-41. [Medline:15711738](#)
- Pengo V, Lensing AW, Prins MH, Marchiori A, Davidson BL, Tiozzo F, et al. Incidence of chronic thromboembolic pulmonary hypertension after pulmonary embolism. *N Engl J Med*. 2004;350:2257-64. [Medline:15163775](#) [doi:10.1056/NEJMoa032274](#)
- Ribeiro A, Lindmarker P, Johnsson H, Juhlin-Dannfelt A, Jorfeldt L. Pulmonary embolism: one-year follow-up with echocardiography doppler and five-year survival analysis. *Circulation*. 1999;99:1325-30. [Medline:10077516](#) [doi:10.1161/01.CIR.99.10.1325](#)
- Heit JA, Mohr DN, Silverstein MD, Petterson TM, O'Fallon WM, Melton LJ III. Predictors of recurrence after deep vein thrombosis and pulmonary embolism: a population-based cohort study.

- Arch Intern Med. 2000;160:761-8. [Medline:10737275](#) [doi:10.1001/archinte.160.6.761](#)
- 19 Griffin J. Deep vein thrombosis and pulmonary embolism. London: Office of Health Economics. 1996;48–59.
 - 20 Bergqvist D, Jendteg S, Johansen L, Persson U, Odegaard K. Cost of long-term complications of deep venous thrombosis of the lower extremities: an analysis of a defined patient population in Sweden. *Ann Intern Med.* 1997;126:454-7. [Medline:9072931](#) [doi:10.7326/0003-4819-126-6-199703150-00006](#)
 - 21 Zakai NA, McClure LA. Racial differences in venous thromboembolism. *J Thromb Haemost.* 2011;9:1877-82. [Medline:21797965](#) [doi:10.1111/j.1538-7836.2011.04443.x](#)
 - 22 Census of Population. Household and Dwellings 2011. Statistical Report. Croatian Bureau of Statistics. Zagreb, 2013.
 - 23 Spirk D, Husmann M, Hayoz D, Baldi T, Frauchiger B, Engelberger R, et al. Predictors of in-hospital mortality in elderly patients with acute venous thrombo-embolism: the SWISS Venous ThromboEmbolic Registry (SWIVTER). *Eur Heart J.* 2012;33:921-6. [Medline:22036872](#) [doi:10.1093/eurheartj/ehr392](#)
 - 24 Coen D, Zadro R, Honovic L, Banfic L, Stavljenic Rukavina A. Prevalence and association of the factor V Leiden and prothrombin G20210A in healthy subjects and patients with venous thromboembolism. *Croat Med J.* 2001;42:488-92. [Medline:11471205](#)
 - 25 Jukic I, Bingulac-Popovic J, Dogic V, Babic I, Culej J, Tomicic M, et al. ABO blood groups and genetic risk factors for thrombosis in Croatian population. *Croat Med J.* 2009;50:550-8. [Medline:20017223](#) [doi:10.3325/cmj.2009.50.550](#)
 - 26 Pulanic D, Zupancic Salek S, Labar B. Thrombophilia-inherited and acquired susceptibility to thrombosis. *Lijec Vjesn.* 2009;131:236-7. in Croatian. [Medline:19769289](#)
 - 27 Rodger RL, Bick MD. Cancer-associated thrombosis. *N Engl J Med.* 2003;349:109-11. [Medline:12853582](#) [doi:10.1056/NEJMp030086](#)
 - 28 Prandoni P, Piccioli A, Girolami A. Cancer and venous thromboembolism: an overview. *Haematologica.* 1999;84:437-45. [Medline:10329923](#)
 - 29 Fimognari FL, Repetto L, Moro L, Gianni W, Incalzi RA. Age, cancer and the risk of venous thromboembolism. *Crit Rev Oncol Hematol.* 2005;55:207-12. [Medline:15979886](#) [doi:10.1016/j.critrevonc.2005.04.011](#)
 - 30 Vucic N, Ostojic R, Sviric T. Treatment of deep vein thrombosis with oral anticoagulants in patients with malignancy: prospective cohort study. *Croat Med J.* 2002;43:296-300. [Medline:12035135](#)
 - 31 Gould MK, Garcia DA, Wren SM, Karanicolas PJ, Arcelus JL, Heit JA, et al. Prevention of VTE in nonorthopedic surgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141 (2 Suppl):e227S-77S.
 - 32 Noble S, Pasi J. Epidemiology and pathophysiology of cancer-associated thrombosis. *Br J Cancer.* 2010;102 Suppl 1:S2-9. [Medline:20386546](#) [doi:10.1038/sj.bjc.6605599](#)
 - 33 Arnold DM, Kahn SR, Shrier I. Missed opportunities for prevention of venous thromboembolism: an evaluation of the use of thromboprophylaxis guidelines. *Chest.* 2001;120:1964-71. [Medline:11742929](#) [doi:10.1378/chest.120.6.1964](#)
 - 34 Ageno W, Squizzato A, Ambrosini F, Dentali F, Marchesi C, Mera V, et al. Thrombosis prophylaxis in medical patients: a retrospective review of clinical practice patterns. *Haematologica.* 2002;87:746-50. [Medline:12091126](#)
 - 35 Rashid ST, Thursz MR, Razvi NA, Voller R, Orchard T, Rashid ST, et al. Venous thromboprophylaxis in UK medical inpatients. *J R Soc Med.* 2005;98:507-12. [Medline:16260800](#) [doi:10.1258/jrsm.98.11.507](#)
 - 36 Trkulja V, Kolundzic R. Rivaroxaban vs dabigatran for thromboprophylaxis after joint-replacement surgery: exploratory indirect comparison based on meta-analysis of pivotal clinical trials. *Croat Med J.* 2010;51:113-23. [Medline:20401953](#) [doi:10.3325/cmj.2010.51.113](#)