

Combined Cataract and Corneal Transplantation Surgery Without Viscosurgical Devices

Kuzman, Tomislav; Bakrac, Ana; Meter, Ana; Gabrić, Ivana; Gaćina, Dina; Pauk, Sanja; Jukić, Anđela; Škegro, Ivan; Masnec, Sanja; Kalauz, Miro

Source / Izvornik: **Acta Informatica Medica, 2023, 31, 186 - 190**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.5455/aim.2023.31.186-190>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:105:016528>

Rights / Prava: [Attribution-NonCommercial 4.0 International/Imenovanje-Nekomercijalno 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-11-27**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



Combined Cataract and Corneal Transplantation Surgery Without Viscoelastic Devices

Tomislav Kuzman¹, Ana Pupic-Bakrac², Ana Meter², Ivana Gabric¹, Dina Lesin Gacina¹, Sanja Vidas Pauk¹, Andjela Jukic², Ivan Skegro¹, Sanja Masnec¹, Miro Kalauz¹

¹Department of Ophthalmology, University Hospital Center Zagreb, School of Medicine, University of Zagreb, Zagreb, Croatia

²Department of Ophthalmology, University Hospital Dubrava, Zagreb, Croatia

Corresponding author: Ana Pupic-Bakrac, Department of Ophthalmology, University Hospital Dubrava, Avenija Gojka Šuška 6, 10000, Zagreb, Croatia. anapupicbakrac@gmail.com, ORCID ID: <https://orcid.org/0000-0002-8493-7305>

doi: 10.5455/aim.2023.31.186-190

ACTA INFORM MED. 2023, 31(3): 186-190

Received: JUN 15, 2023

Accepted: SEP 05, 2023

© 2023 Tomislav Kuzman, Ana Pupic-Bakrac, Ana Meter, Ivana Gabric, Dina Lesin Gacina, Sanja Vidas Pauk, Andjela Jukic, Ivan Skegro, Sanja Masnec, Miro Kalauz

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: The most common complications after performing the triple Descemet's stripping automated endothelial keratoplasty (DSAEK), which combines the cataract phacoemulsification, intraocular lens implantation and DSAEK procedure, are detachment or decentration of the donor lamella and postoperative interface haze. One reason for this is the retained viscoelastic used during surgery. **Objective:** This study aimed to describe triple DSAEK procedure without the usage of viscoelastic and to discuss its potential benefits on surgical outcomes. **Methods:** The surgical procedures and outcomes of patients with Fuchs' dystrophy and lens opacification who underwent the triple DSAEK were retrospectively reviewed. The surgical procedure was described, and postoperative complications were studied. **Results:** The study included 10 eyes of 10 patients. Capsulorhexis and IOL implantation performed in locally potentiated anesthesia compared to general anesthesia did not significantly differ ($P > 0,05$). The mean preoperative best-corrected visual acuity was 0.75 LogMar. The mean postoperative best-corrected visual acuity was 0.2 LogMar. The central graft thickness before surgery was 129.6 μm and 6 months after surgery was 114.2 μm . Successful attachment of the donor lamellae was observed in all 10 patients. None of the patients had postoperative interface haze or any other possible viscoelastic caused complication. **Conclusion:** Although viscoelastic can facilitate certain aspects of the triple DSAEK procedure, we conclude that this procedure can be performed completely without its use. If performed by a trained surgeon, the procedure can be feasible without the complications of donor lamella detachment, decentration, or interface haze.

Keywords: Triple procedure, Viscoelastic, DSAEK, Phacoemulsification.

1. BACKGROUND

The triple procedure combining phacoemulsification, intraocular lens (IOL) implantation and Descemet's stripping automated endothelial keratoplasty (DSAEK) in patients with Fuchs' dystrophy (FECD) and cataract provides more rapid visual improvement. The additional advantage is the avoidance of second surgical procedure. In patients with FECD and cataracts requiring both DSAEK and cataract surgery, operation can be performed concurrently or sequentially with DSAEK shortly after phacoemulsification and IOL implantation (1).

Many ophthalmic viscosurgical devices (OVDs) are used worldwide to facilitate cataract surgery. The advantages of viscoelastic during cataract surgery

are well known, and phacoemulsification and IOL implantation without the use of viscoelastic are not performed in everyday surgical practice. The main viscoelastic function is to maintain anterior chamber depth, facilitate capsulorhexis, protect and stabilize intraocular tissues (corneal endothelium and the posterior capsule) and facilitate IOL implantation. A deep anterior chamber reduces the chance of donor endothelial contact with the iris and IOL (2).

The use of viscoelastic during DSAEK can simplify descementorhexis in certain cases (3). However, some potential side effects of residual OVDs have been observed postoperatively after cataract surgery as well as DSAEK. Residual viscoelastic can cause increased intraocular pressure owing to the blockage of

the trabecular meshwork and postoperative elevated intraocular eye pressure, toxic anterior segment syndrome (TASS), and posterior capsule opacification (4-7). OVD needs to be completely removed at the end of the surgery, which increases the duration of surgery. Moreover, viscoelastic substances increase the final cost of surgery. In cases of the DSAEK procedure, regardless combined with phacoemulsification or not residual viscoelastic may remain at the graft-host interface.

One of the most common complication of DSAEK is graft detachment, with average incidence rate of 15% (8, 9). The causes of graft dislocation can be multifactorial and include tissue storage conditions, surgical techniques, pre-existing conditions and lack of patient cooperation. It is suggested that surgeons should minimize the amount of viscoelastic used during procedure (10). Retained viscoelastic may impact graft adherence and lead to posterior graft dislocation and detachment (11). Poor graft adherence can also cause graft failure and rejection (12, 13). Further, retained viscoelastic can affect visual outcomes by producing haze at the interface. Several authors reported cases of residual viscoelastic as a direct cause of interface haze (12, 14), however there is a lack of studies which show exact rate of viscoelastic caused graft dislocation and detachment. To the best of our knowledge, there is insufficient information about performing the triple DSAEK procedure without any viscoelastic.

2. OBJECTIVE

Our study aimed to describe the triple DSAEK procedure without viscoelastic agents and to discuss its potential benefits on surgical outcomes. Further, we aimed to investigate whether we would have above mentioned complications if we avoided the use of viscoelastic.

3. MATERIAL AND METHODS

A retrospective analysis (retrospective chart review) of the medical records of patients who underwent a triple procedure, including phacoemulsification, IOL implantation, and DSAEK without ophthalmic viscosurgical devices, was conducted. Ethical committee approval was obtained from the institution before performing the research.

The operations were performed by a single surgeon at a single tertiary referral center. All participants were diagnosed with Fuchs' endothelial dystrophy grade 4–6 and nuclear opacification (NO3, NO4, or NO5), and cortical opacification (C2 or C3), according to the Lens Opacities Classification System III scale. Ten eyes of 10 patients who underwent the triple procedure without viscoelastic treatment were identified. The patients did not have concomitant ophthalmic diseases, such as glaucoma or retinal diseases. Patients with a postoperative follow-up period of < 6 months were excluded.

Demographic data such as sex and age at the time of diagnosis were recorded.

The patients underwent complete ophthalmological examination preoperatively and postoperatively (6 months), which included best-corrected visual acuity measured (Snellen table, ETDRS – logMAR table), slit-lamp examination, Goldmann tonometry, and fundus examination. The thickness of the lamella and cornea were measured with anterior segment optical coherence tomography (OCT) Visante (Carl Zeiss Meditec, Jena, Germany).

Patients underwent phacoemulsification with IOL implantation and DSAEK (triple procedure), IOL power calculation using IOL Master 700 (Carl Zeiss, Jena, Germany), and SRK/T formula. After the triple DSAEK occurs hyperopic shift, which we calculated during IOL measurement. Implanted IOLs were +1.25 diopters higher, which compensated for the reduction in corneal power by DSAEK.

Donor corneal lamellas were pre-cut and preserved using conventional eye bank techniques. The corneas were stored in the eye bank in either hypothermic storage or tissue culture media. Corneas in a hypothermic storage medium were stored at a temperature of 4°C for a maximum of seven days, and were then prepared for lamellar keratoplasty immediately before planned operation. The other corneas were kept in a tissue culture media at a temperature of 31°C for a maximum of 28 days. Corneas in the storage become thicker and in order to return to the physiological thickness, it must be stored in a transmedia containing dextran, for at least for 24 hr before the preparation for lamellar keratoplasty. The corneas were microbiologically tested during storage. The procedure of preparing corneas for lamellar keratoplasty was performed by specially trained employees using the automatic microkeratome (Gebauer Slc Original, Neuhausen, Germany). Each cornea was placed on the artificial eye chamber filled with corneal storage media to preserve the endothelial cell viability. The epithelium of each cornea was removed to make the cut as precise as possible. The permissible variation of the knife, which was 30 µm, should be considered. The cornea was carefully removed from the artificial eye chamber and stored in a transport media for delivery to transplantation center.

The surgeries were performed under general anesthesia or locally potentiated anesthesia.

One side port entrance made at the beginning of the cataract surgery at the 6 o'clock position was used for continuous irrigation with the balanced salt solution (BSS). The bottle was raised to a height of 60–80 cm of water and adjusted according to the anterior chamber depth. The other side port was made at the 2 o'clock position for the chopper. Capsulorhexis was started using a 26-gauge cystitome needle and continued with forceps under BSS irrigation through the main 2.75 mm port at the 12 o'clock position. Stop and chop phacoemulsification were performed using the Alcon Centurion device (Alcon, Fort Worth, TX, USA). IOL implantation (Johnson & Johnson Sensar One-piece AAB00) into the capsular bag was performed under continuous BSS irrigation. The cartridge was also filled with BSS.

Based on the surgeons' personal experience, we graded the complexity of performing capsulorhexis without viscoelastic during the procedure from 1 to 3, where 1 = easy, 2 = moderate, and 3 = difficult to perform. The same gradation was used with IOL implantation (1 = easy; 2 = moderate; and 3 = difficult to implant) (Table 1).

After phacoemulsification and IOL implantation, Mio-stat (Carbachol 0.01%, Alcon) was injected into the anterior chamber to perform miosis. Subsequently, under continuous irrigation, peripheral iridotomy at the 6 o'clock position was performed with vertical vitreoretinal scissors to avoid the potential postoperative pupillary block.

An 8.0 mm diameter descemetorhexis under an air bubble

Complexity grade	Capsulorexis	IOL implantation
Easy (grade 1)	Continuous capsulorexis in one act	IOL implantation in one act
Moderate (grade 2)	Discontinuation of capsulorexis 1-2 times and waiting for formation of the anterior chamber	Short-term collapse of the anterior chamber
Difficult (grade 3)	Multiple discontinuation of capsulorexis >2 times and waiting for formation of the anterior chamber	Short-term collapse of the anterior chamber with difficulty positioning the IOL

Table 1. Criteria used to evaluate complexity grade of performing capsulorhexis and IOL implantation without viscoelastic.

Complexity grade	Easy	Moderate	Difficult	Fisher's exact test
Capsulorexis:				
GA	3	2	0	
LPA	2	2	1	
Total	5	4	1	P > 0,05
IOL implantation:				
GA	3	2	0	
LPA	2	2	1	
Total	5	4	1	P > 0,05

Table 2. Gradation of the complexity of performing capsulorhexis and IOL implantation based on surgeons' personal experience. GA – general anesthesia; LPA – locally potentiated anesthesia; IOL – intraocular lens.

was made by the reverse Sinsky hook through the side port at the 2 o'clock position. The central part of the Descemet membrane was removed from the eye using endothelial 23-gauge grasping forceps.

An 8.0 mm trephine was used to separate the donor graft lamella. A Busin glider was used to insert the donor graft into the anterior chamber through the main incision. Air bubble was then injected into the anterior chamber to attach the graft to the stroma. Finally, reconstituted cefuroxime (1 mg/0.1 mL) was injected into the anterior chamber. We performed control examinations of the patients after 8 hours, the next day, in 7 days, in 2 weeks, and then every month after the procedure. Postoperative follow up was without complications.

After obtaining the necessary data, statistical analysis was conducted. SPSS for Windows (version 13.0, SPSS Inc. Chicago, IL, USA) and Microsoft Excel (version of Office 2007; Microsoft Corporation, Redmond, WA, USA) were used. Data are expressed as counts and percentages for categorical variables and as means with standard deviations or medians with ranges for continuous variables. Fisher's exact test was used to analyze the differences between categorical variables. Statistical significance was set at P<0.05.

4. RESULTS

The present study included 10 eyes of 10 patients. Within the study population, there were nine women and one man. The mean age of the patients who underwent the procedure was 68 years (range 54–81 years). Four patients had Fuchs' dystrophy grade 4, four had Fuchs' dystrophy grade 5, and two presented with grade 6. The mean preoperative best-corrected visual acuity was 0.75 LogMar, ranging from 1.3 to 0.5 LogMar. The central graft thickness before surgery, measured by ultrasound pachymetry, ranged from 101 to 165 μm, with a mean of 129.6 μm. The central graft thickness 6 months after

surgery, measured by anterior OCT, ranged from 89 to 135 μm, mean of 114.2 μm.

Gradation of the complexity of performing capsulorhexis and IOL implantation in general and locally potentiated anesthesia is shown in Table 1. As seen from the Table 2, both capsulorhexis and IOL implantation performed in locally potentiated anesthesia compared to general anesthesia did not significantly differ (P > 0,05).

The mean postoperative best-corrected visual acuity was 0.2 LogMar, ranging from 0.22 to 0.1 LogMar. The mean intraocular pressure (IOP) on the first postoperative day measured with Goldman applanation tonometry was 14.6 mmHg, ranging from 12 to 18 mmHg. 6None of the patients had postoperative complications, including graft dislocation and interface fluid; therefore, none of the patients underwent postoperative rebubbling. In addition, none of the patients experienced interfacial haze.

5. DISCUSSION

The most common complications of DSAEK surgery are graft dislocation and detachment (8, 12, 13). These complications usually require rebubbling or recentering the graft postoperatively. The American Academy of Ophthalmology reports a wide range frequency of detachment after DSAEK, from 0% to 82%; however, the average rates are approximately 15% (8). The rebubbling rate differs among studies from 5% to 50% (15).

Various potential causes of graft dislocation or detachment are mentioned in the literature, such as improper tissue storage conditions, surgical techniques and patients preexisting conditions. Postoperative patient behavior is also important as some patients can squeeze and rub their eyes which leads to potential graft dislocation (9, 16-18). Among the many causes of graft dislocation and detachment, one of them could be viscoelastic, which is used during surgery. OVDs used during the triple procedure should be completely evacuated because incomplete removal and retained viscoelastic disables good adherence of donor cornea to the recipient stroma and may lead to graft dislocation or detachment. It is also suggested that surgeons should minimize the amount of viscoelastic used during procedure (10). By analyzing literature, we found case reports of residual viscoelastic as a direct cause of interface haze (12, 14). However, we did not find empirical data on exact incidence of residual viscoelastic causing graft dislocation and detachment. Further, in cases of retained viscoelastic after DSAEK, as shown by certain cases in partial detachment, surgeons recommend removal in second surgical procedure to minimize interface haze (12). Our idea was to remove the viscoelastic as one of the potential causes of above mentioned complications during the triple DSAEK procedure, since other causes are difficult to influence (such as postoperative patient behavior, donor tissue quality). We wondered what percentage of graft dislocation and detachment would be if we completely excluded the viscoelastic from the triple DSAEK procedure.

Owing to possible complications, many surgeons avoid use of OVDs during DSAEK; however, the triple procedure includes phacoemulsification in which viscoelastic is used almost inevitably for safe capsulorhexis and IOL implantation (19).

To the best of our knowledge, this is the first report on a triple procedure without viscoelastic in the literature. Several studies have been conducted on phacoemulsification without the use of viscoelastic (20-25). Goles et al. indicated that phacoemulsification can be performed by trained surgeons without viscoelastic with the same level of safety regarding endothelial cell damage (23). These studies claimed that phacoemulsification without viscoelastic is safe, with very low postoperative complications. Moreover, the reduced time of the surgery and the reduced costs of procedures stand out as advantages.

Prompted by possible complications and encouraged by phacoemulsification studies without viscoelastic, we started to perform a triple procedure without viscoelastic agents.

In our study, all procedures were successfully performed without postoperative complications (graft dislocation or detachment, increased intraocular pressure, TASS, and interface haze). In experienced surgeon hands, three basic steps in which viscoelastic is normally used can be successfully performed without, by using a careful surgical technique. Capsulorhexis and IOL implantation in this study were performed under continuous infusion, and Descemetorexis was performed under an air bubble.

According to Table 2, it seems that capsulorhexis and IOL implantation are equal to perform under general anesthesia or locally potentiated anesthesia. In general anesthesia, the impact of patient cooperation during the procedure is minimized so it is expected that the procedure would be easier. However, if the anterior chamber is too unstable intraoperatively, we can always add viscoelastic and finish the triple DSAEK procedure with viscoelastic.

Limitations of the study

This study has several limitations. First limitation is small sample size, however it showed feasibility to conduct research with larger sample size. Basing on this study in larger sample size we should generate more accurate results. Considering that we do not have many patients whose condition requires a triple procedure, it is difficult to collect a larger number for research. Therefore, a multicenter study should be conducted to have a larger sample. Further, we are lacking previous studies in the this topic, and we can not compare our results with others. There are studies in the literature describing phacoemulsification surgery and DSAEK without viscoelastic, however we did not find any study about triple DSAEK procedure without viscoelastic.

6. CONCLUSION

To the best of our knowledge, this is the first study on a triple procedure without viscoelastic in the literature. Although viscoelastic can facilitate certain parts of the operation, we can conclude that this procedure can be performed completely without its use. If performed by a trained surgeon, the procedure can be feasible without complications of donor lamella detachment, decentration, or interface haze

The future controlled trials are recommended to see if the triple DSAEK procedure without viscoelastic has fewer complications.

• **Acknowledgements:** The authors thank Dr. Jure Pupić-Bakrač for careful reading of the manuscript, insightful suggestions and help with

table design.

- **Patient Consent Form:** Case reports are retrospective. Only archived patients' data and samples processed for diagnostic purposes were used. Prior the processing, written consent with examination and diagnostic tests, as well as with potential further use of the data and samples for scientific or educational purposes including publication of anonymized data was provided by each particular patient.
- **Athors contribution:** TK: Conceptualization, Investigation, Writing–Original Draft, Writing – Review and editing. APB: Investigation, Writing–Original Draft, Visualization. AM, IG, DLG, SVP, AJ, IŠ, SM, MK: Resources, Review and editing.
- **Conflict of interest statement:** The authors have no conflicts of interest to disclose.
- **Financial support and sponsorship:** The authors received no specific funding for this work.

REFERENCES

1. Covert DJ, Koenig SB. New triple procedure: Descemet's stripping and automated endothelial keratoplasty combined with phacoemulsification and intraocular lens implantation. *Ophthalmology*. 2007; 114(7): 1272-1277.
2. Bissen-Miyajima H. Ophthalmic viscosurgical devices. *Curr Opin Ophthalmol*- 2008; 19:50.
3. Melki SA, Harissi-Dagher M, Wu S, Fava MA: Viscoelastic-assisted DSAEK. *Ophthalmology*. 2011; 118: 1003.
4. Van den Bruel A, Gailly J, Devriese S, Welton NJ, Shortt AJ, Vrijens F: The protective effect of ophthalmic viscoelastic devices on endothelial cell loss during cataract surgery: a meta-analysis using mixed treatment comparisons. *Br J Ophthalmol* 95:5, 2011
5. Dick HB, Augustin AJ, Pakula T, Pfeiffer N: Endotoxins in ophthalmic viscosurgical devices. *Eur J Ophthalmol* 13:176, 2003
6. Bissen-Miyajima H: In vitro behavior of ophthalmic viscosurgical devices during phacoemulsification. *J Cataract Refract Surg* 32:1026, 2006
7. Altintas AK, Ciritoglu MY, Beyazyildi ZO, Can CU, Polat S: Toxic Anterior Segment Syndrome Outbreak after Cataract Surgery Triggered by Viscoelastic Substance. *Middle East Afr J Ophthalmol* 24:43, 2017
8. Lee WB, Jacobs DS, Musch DC, Kaufman SC, Reinhart WJ, Shtein RM: Descemet's stripping endothelial keratoplasty: safety and outcomes: a report by the American Academy of Ophthalmology. *Ophthalmology* 116:1818, 2009
9. Price MO, Price FW, Jr.: Endothelial keratoplasty—a review. *Clin Exp Ophthalmol* 38:128, 2010
10. Yoo SH, Chang V: Prevention and management of Descemet's stripping automated endothelial keratoplasty complications. In Golnik KC (ed) *The Ophthalmic News and Education Network, Clinical Education / News / Current Insight, Cornea/ External Disease*, (ed. San Francisco, CA, US, American Academy of Ophthalmology, 2009,
11. Terry MA, Ousley PJ: Deep lamellar endothelial keratoplasty: early complications and their management. *Cornea* 25:37, 2006
12. Anshu A, Planchard B, Price MO, da RPC, Price FW, Jr.: A cause of reticular interface haze and its management after descemet stripping endothelial keratoplasty. *Cornea* 31:1365, 2012
13. Kymionis GD, Voulgari N, Kontadakis GA, Mikropoulos D, Petrovic A, Droutsas K: Surgical management of post-Descemet stripping automated endothelial keratoplasty interface

- haze associated with interface deposits. *Indian J Ophthalmol* 68:174, 2020
14. Vira S, Shih CY, Ragusa N, Sheyman A, Feder R, Weisenthal RW, Rosenwasser GO, Hannush SB, Udell IJ, Bouchard CS: Textural interface opacity after descemet stripping automated endothelial keratoplasty: a report of 30 cases and possible etiology. *Cornea* 32:e54, 2013
 15. Parekh M, Leon P, Ruzza A, Borroni D, Ferrari S, Ponzin D, Romano V: Graft detachment and rebubbling rate in Descemet membrane endothelial keratoplasty. *Surv Ophthalmol* 63:245, 2018
 16. Suh LH, Yoo SH, Deobhakta A, Donaldson KE, Alfonso EC, Culbertson WW, O'Brien TP: Complications of Descemet's stripping with automated endothelial keratoplasty: survey of 118 eyes at One Institute. *Ophthalmology* 115:1517, 2008
 17. Price FW, Jr., Price MO: Descemet's stripping with endothelial keratoplasty in 200 eyes: Early challenges and techniques to enhance donor adherence. *J Cataract Refract Surg* 32:411, 2006
 18. Dirisamer M, van Dijk K, Dapena I, Ham L, Oganés O, Frank LE, Melles GR: Prevention and management of graft detachment in descemet membrane endothelial keratoplasty. *Arch Ophthalmol* 130:280, 2012
 19. Higashide T, Sugiyama K: Use of viscoelastic substance in ophthalmic surgery—focus on sodium hyaluronate. *Clin Ophthalmol* 2:21, 2008
 20. Ogurel T, Ogurel R, Onaran Z, Ornek K: Safety of hydroimplantation in cataract surgery in patients with pseudoexfoliation syndrome. *Int J Ophthalmol* 10:723, 2017
 21. Taskin I, Aslan L: Effects of phacoemulsification with versus without viscoelastic devices on surgical outcomes. *Int Ophthalmol* 38:5, 2018
 22. Rao A, Padhy D, Das G, Sarangi S: Viscoless Manual Small Incision Cataract Surgery with Trabeculectomy. *Semin Ophthalmol* 33:552, 2018
 23. Goles N, Nerancic M, Konjik S, Pajic-Eggspuehler B, Pajic B, Cvejic Z: Phacoemulsification and IOL-Implantation without Using Viscoelastics: Combined Modeling of Thermo Fluid Dynamics, Clinical Outcomes, and Endothelial Cell Density. *Sensors (Basel)* 21, 2021
 24. Bardoloi N, Sarkar S, Paliana A, Das H: Pure phaco: phacoemulsification without ophthalmic viscosurgical devices. *J Cataract Refract Surg* 46:174, 2020
 25. Ozcura F, Cevik S: Hydroimplantation versus viscoimplantation: comparison of intraocular lens implantation with and without ophthalmic viscoelastic device in phacoemulsification. *Rom J Ophthalmol* 62:282, 2018