

Polymorphisms in Survivin (BIRC5 Gene) Are Associated with Age of Onset in Breast Cancer Patients

Sušac, Ilona; Ozretić, Petar; Gregorić, Maja; Levačić Cvok, Mirela; Sabol, Maja; Levanat, Sonja; Trnski, Diana; Eljuga, Domagoj; Seiwerth, Sven; Aralica, Gorana; ...

Source / Izvornik: **Journal of Oncology, 2019, 2019**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1155/2019/3483192>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:955145>

Rights / Prava: [Attribution 4.0 International](#) / [Imenovanje 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-07-23**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



Research Article

Polymorphisms in Survivin (*BIRC5* Gene) Are Associated with Age of Onset in Breast Cancer Patients

Ilona Sušac,¹ Petar Ozretić ,² Maja Gregorić,³ Mirela Levačić Cvok,^{2,4} Maja Sabol,² Sonja Levanat,² Diana Trnski,² Domagoj Eljuga,^{1,5} Sven Seiwerth,^{6,7} Gorana Aralica,^{6,8} Mladen Stanec,⁵ and Vesna Musani ²

¹Eljuga Polyclinic, 10000 Zagreb, Croatia

²Division of Molecular Medicine, Ruđer Bošković Institute, 10000 Zagreb, Croatia

³Zagreb Health School, 10000 Zagreb, Croatia

⁴Kardinal Alojzije Stepinac Krašić Primary School, 10454 Krašić, Croatia

⁵Department for Oncoplastic and Reconstructive Surgery, University Hospital for Tumors, 10000 Zagreb, Croatia

⁶Institute of Pathology, University of Zagreb School of Medicine, 10000 Zagreb, Croatia

⁷Clinical Department of Pathology and Cytology, University Hospital Centre Zagreb, 10000 Zagreb, Croatia

⁸Department of Pathology, Clinical Hospital Dubrava, 10000 Zagreb, Croatia

Correspondence should be addressed to Vesna Musani; vmusani@irb.hr

Received 9 October 2018; Revised 7 June 2019; Accepted 25 June 2019; Published 28 July 2019

Academic Editor: Giandomenico Roviello

Copyright © 2019 Ilona Sušac et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Survivin, encoded by *BIRC5* gene (baculoviral IAP repeat containing 5), belongs to the family of inhibitors of apoptosis proteins (IAPs). In mammalian cells it participates in the control of mitosis, apoptosis regulation, and cellular stress response. Its expression is increased in almost all types of cancers. The aim of this study was to investigate the role of *BIRC5* polymorphisms in breast cancer (BC) and to connect survivin expression with various clinicopathological characteristics of BC patients. Blood and archival tumour tissue samples were collected from 26 BC patients from Croatia. Survivin expression was determined immunohistochemically. *BIRC5* promoter, coding region, and 3'UTR were genotyped. DNA from 74 healthy women was used as control. *BIRC5* polymorphisms and survivin expression were tested against age of onset, histological grade, tumour type and size, lymph node status, oestrogen, progesterone, Her2, and Ki67 status. Numbers of samples with weak, moderate, and strong survivin expression were 9 (33.3%), 11 (40.7%), and 7 (25.9%), respectively. Most patients had nuclear survivin staining (92.6%). High survivin expression was significantly associated with negative oestrogen receptor status ($p=0.007$) and positive Ki67 expression ($p=0.032$). Ki67 expression was also positively correlated with histological grade ($p=0.0009$). Fourteen polymorphisms were found in BC samples, located mostly in promoter and 3'UTR of *BIRC5*. There was no significant difference in the distribution of polymorphisms between BC and control samples. Among clinicopathological characteristics of BC patients, alleles of five *BIRC5* polymorphisms were associated with younger age of onset: c.-644T>C (55.8 years [y] vs. 48.1 y; $p=0.006$), c.-241C>T (54.2 y vs. 45.0; $p=0.029$), c.9809T>C (55.8 y vs. 48.1 y; $p=0.006$), c.-1547C>T (58.3 y vs. 50.9 y; $p=0.011$), and c.9386T>C (50.8 y vs. 59.5 y; $p=0.004$). To assess the significance of *BIRC5* polymorphisms and survivin expression as predictive and prognostic biomarkers for BC further research with a larger sample size is needed.

1. Introduction

Breast cancer (BC) is one of the most common malignancies among women worldwide. Despite generally good prognosis for BC patients, there is a wide variation in survival [1]. Some of the common risk factors for BC include age, genetic

background, hormonal factors, reproductive and menstrual history, excessive alcohol consumption, radiation, benign breast disease, and obesity [2]. Genetic variation has been shown to affect both susceptibility and prognosis of BC [1]. Genome wide association studies (GWAS) have recently been used to identify new loci for BC, and so far almost

100 loci have been identified [2]. In addition to providing new prognostic markers, genetic determinants of prognosis can also give new biological insight for progression of BC [3]. Survivin, encoded by *BIRC5* (baculoviral IAP repeat containing 5) gene, belongs to the family of inhibitors of apoptosis (IAP) proteins [4]. It is the smallest of eight IAP proteins and contains a single Baculovirus IAP Repeat (BIR), which is the hallmark of these molecules [5, 6]. Survivin is a multifunctional protein and in mammalian cells it participates in at least three cellular processes: the control of mitosis, the regulation of apoptosis, and the cellular stress response [5]. Survivin is strongly expressed during development, while in healthy organisms it is not expressed in differentiated tissues, and its expression is markedly increased in almost all types of cancers (including bladder cancer, lung cancer, breast cancer, stomach, oesophagus, pancreas, liver, uterine, ovarian, and haematological cancers) [7]. Elevated expression of survivin in BC is associated with increased resistance to chemo- and radiotherapy, increased grade, lymph node invasion, and tumour size, as well as decreased survival rate [8]. According to Marsicano et al., breast cancer patients with higher expression of *BIRC5* have significantly greater chance of developing metastasis than patients with lower expression [9]. Survivin is expressed in a cell cycle-regulated manner, being maximally expressed during the G2/M phase, followed by rapid decline of both the mRNA and protein levels at the G1 phase [10]. Several *BIRC5* polymorphisms were shown to be associated with susceptibility (gastric [11], bladder [12], and hepatocellular [13]), survival (colorectal [14] and breast [15]), or age of onset (ovarian cancer [4]). Some of these polymorphisms create binding sites for either transcription factors or regulatory miRNA and potentially can influence survivin levels. Survivin has been proposed as a potential target for anticancer intervention and several survivin inhibitors and survivin-related molecular therapies are in development [16]. The purpose of this preliminary study was to investigate the role of *BIRC5* polymorphisms in BC. The connection between polymorphisms found and protein expression in tumour tissue and their association was also assessed.

2. Materials and Methods

2.1. Study Subjects. This study included BC patients who had their regular check-up at the Eljuga Polyclinic and agreed to participate. Upon signing the consent forms their blood was drawn. 26 patients, diagnosed between 2000 and 2012 and whose archival paraffin embedded tumour tissue was available from archive at the University Hospital for Tumours, University Hospital Centre “Sestre Milosrdnice,” were included in the study. One patient had a bilateral BC and both tumours were available for testing. Demographic and clinicopathological data on patients was collected from medical records. Demographic characteristics included age of onset and time since operation, and clinicopathological characteristics included histological grade, tumour size, oestrogen and progesterone receptor status, Her2 status, lymph node status, type, and Ki67 status (determined routinely as part of the diagnostic procedure in the clinic). Two

5- μ m sections of tumour tissue were cut from paraffin block for immunohistochemical analysis. DNA samples from 74 healthy women with no history of BC (median age 80, range 65-101) were collected in our previous study [17]. Although usual case-control studies are age matched, in this case it was more appropriate to use elderly healthy population as control. This population is less likely to have cancer predisposing polymorphisms. The study was approved by the Institutional Ethics Committee (EP-6820/13-11).

2.2. Immunohistochemical Staining. Survivin protein was analysed as follows: slides were deparaffinized, prewarmed in Epitope Retrieval Solution (Dako, Glostrup, Denmark) and kept at 95-99°C for 20 min, left to cool for additional 20 min, and then washed in PBS 3x5 min. Endogenous peroxidase was blocked with 3 % H₂O₂/methanol for 10 min and then washed 3x5 min in PBS. The sample was circled by PAP-PEN (Kiyota, Baltimore, MD). Protein block serum-free (DAKO) was added for 10 min.

Immunohistochemical staining was performed using monoclonal rabbit anti-survivin antibody (1:200, 71G4B7, Cell Signalling Technology, Danvers, MA). Following an overnight incubation, the immunodetection was completed using the LSAB Visualization System (DAKO) utilizing 3, 3'-diaminobenzidine (DAB) chromogen as substrate, according to the manufacturer's instructions. All sections were counterstained with haematoxylin (DAKO). Negative controls were obtained by omitting the primary antibody.

Stained slides were examined by two experienced pathologists. Samples were scored between 1 and 3 based on survivin expression, taking into account the percentage of positive cells and the staining intensity, as previously described by Xu et al. [18]. Staining intensity was expressed as + (weak), ++ (moderate), and +++ (intense). Percentage of tumor cells positive for survivin staining was categorized as 0 (no positive cells), 1 (<10 %), 2 (10-50 %), and 3 (>50 %). Final score was then generated by multiplying the staining intensity with the percentage of positive cells and ranged from 0 (no stained tumor cells) to 9 (>50 % of intensively stained tumor cells). The final score of 0 was considered as negative (0), 1-3 as weak (1), 4-6 as moderate (2), and 7-9 points as strong (3).

2.3. SNP Selection and Genotyping. Whole coding region of the *BIRC5* gene (including alternative exons 2alpha, 2B, and 3B) was genotyped, including the most common SNPs in *BIRC5* promoter and the 3' untranslated region (3'UTR). Polymorphisms that were selected from the National Center for Biotechnology Information SNP database (<http://www.ncbi.nlm.nih.gov/snp>) are those usually used in association studies (see [4, 10, 13-15, 19-21]). The majority of these studies have found that polymorphisms in 5' and 3' regions have the greatest impact on survivin expression [22] and might contribute to risk of various cancers [21]. Genomic DNA from blood was extracted by the salting out method. Thirteen PCR fragments were analysed using high resolution melting analysis on High-Resolution Melter (HR-1, Idaho Technology, USA) as described by Cvok et al. [17] followed by Sanger sequencing (ABI PRISM 310 Genetic Analyzer, Applied Biosystems, USA). Due to presence of 5 different

TABLE 1: Patient clinicopathological characteristics.

Characteristic	No. of patients (%)
Age at diagnosis (years)	
< 40	1 (3.8)
40-50	9 (34.6)
50-60	8 (30.8)
> 60	8 (30.8)
Years since operation	
5	8 (30.8)
10	15 (57.7)
15	3 (11.5)
Histological grade	
1	4 (14.8)
2	12 (44.4)
3	10 (37.0)
N.A. *	1 (3.7)
Tumour size	
1 (≤ 2 cm)	20 (74.1)
2 ($> 2 \leq 5$ cm)	6 (22.2)
3 (> 5 cm)	1 (3.7)
Oestrogen receptor status	
Positive	21 (77.8)
Negative	6 (22.2)
Progesterone receptor status	
Positive	16 (59.3)
Negative	11 (40.7)
Her2 status	
Positive	2 (7.4)
Negative	22 (81.5)
N.A.	3 (11.1)
Lymph node status	
Positive	11 (40.7)
Negative	16 (59.3)
Pathologic Characteristics	
Invasive ductal carcinoma	22 (81.5)
Medullary carcinoma	2 (7.4)
Invasive lobular carcinoma	1 (3.7)
Mixed invasive ductal/lobular carcinoma	1 (3.7)
Papillary carcinoma	1 (3.7)
Ki67 status	
Positive	12 (44.4)
Negative	12 (44.4)
N.A.	3 (11.1)

* Patient data not available.

polymorphisms in the PCR product of exon 4, it was directly sequenced. PCR primer sequences and cycling conditions are listed in Supplementary Table 1.

2.4. Statistical Analysis. Genotype and allele frequencies of patients and controls were compared using χ^2 test. Deviation from the Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium were calculated using the online tool SHEsis

TABLE 2: Levels and localization of survivin expression.

Characteristic	No. of samples (%)
Survivin immunoreactivity score	
weak	9 (33.3)
moderate	11 (40.7)
strong	7 (25.9)
Survivin cellular localization	
cytoplasmic	1 (3.7)
nuclear	25 (92.6)
cytoplasmic+nuclear	1 (3.7)

[<http://analysis.bio-x.cn/myAnalysis.php>] [23]. The χ^2 test was used for testing the association between genetic data and categorical clinicopathological variables. Independent samples t-test and one-way analysis of variance with Tukey-Kramer post hoc test were used for testing the difference between continuous (age of onset and logarithmic transformed tumour size) and categorical variables. Two-tailed p-values less than 0.05 were considered statistically significant. Statistical analysis was performed using MedCalc v18.2.1 (MedCalc Software bvba).

3. Results and Discussion

3.1. Patient Characteristics. A total of 26 patients with BC were recruited in this study. The main clinical and histopathological features are summarized in Table 1. The median age of patients at diagnosis was 53 years (range 37-69): only one patient (3.8%) was younger than 40 years, 34.6% were between 40 and 50 years, 30.8% were between 50 and 60, and 30.8% were older than 60. The majority of patients (96.2%) were still alive at the end of the study so survival analysis was not performed.

Tumour size was associated with positive lymph nodes ($p=0.004$). Histological grade was associated with Ki67 expression ($p=0.0009$). All samples with histological grade 1 were negative for Ki67, while only one sample with histological grade 3 was negative for Ki67. Progesterone receptor (PR) and oestrogen receptor (ER) status were weakly positively associated (0.053). These results are in agreement with previously published international data [24-27].

3.2. Survivin Expression. Survivin expression was analysed in 27 BC samples from 26 patients using immunohistochemical staining. Samples were scored between 1 and 3 based on survivin expression, taking into account the percentage of positive cells and the staining intensity (Figure 1). The numbers of samples exhibiting survivin expression with a score of 1 (weak), 2 (moderate), and 3 (strong) were 9 (33.34%), 11 (40.72%), and 7 (25.94%), respectively (Table 2). The majority of patients had nuclear survivin staining (92.6%), so we combined all the samples together for the statistical evaluation.

The high nuclear staining of survivin is different than most published data, where the nuclear staining is mostly lower than cytoplasmic [18, 28-31] and only rarely has been described as equal or higher than cytoplasmic [32, 33].




FIGURE 1: Examples of scored staining intensities: (a) negative control, (b) score 1, (c) score 2, and (d) score 3.

3.3. Association of Survivin Expression with Clinicopathological Parameters. Clinicopathological findings (age at diagnosis, time since operation, histological grade, tumour size, oestrogen and progesterone receptors, Her2 status, lymph node status, and Ki67 status) were compared with the immunohistochemical characterization of survivin expression. Since the majority of samples (22 of 27 samples, 84.6%) were ductal BC all samples were grouped together. The statistical analyses revealed a significant association between survivin expression and ER status. High survivin expression was significantly associated with negative ER status ($p=0.007$). No samples with low survivin expression (score 1), 18.2% samples with medium expression (score 2), and 57.1% of samples with high survivin expression (score 3) had negative ER status (Figure 2). Survivin expression was weakly associated with negative PR status ($p=0.051$). Furthermore, another statistically significant association was confirmed between survivin expression and Ki67 expression. High survivin expression was associated with positive Ki67 expression ($p=0.032$).

There is controversial data reported about association between high survivin expression and negative ER and/or PR status in BC. While some studies [34, 35] found the same association as here, other studies report these parameters as not significant [32, 36, 37]. The similar situation is with the association between high survivin expression and Ki67 expression. While some studies [38] showed the same results as here, others showed there was no significant association between survivin and Ki67 [31, 39].




FIGURE 2: Association between survivin expression and ER status. Survivin expression was ranked weak, moderate, and strong, and percentage of ER negative and ER positive samples per group is shown.

3.4. BIRC5 Polymorphisms in Breast Cancer Patients. Fifteen different polymorphisms were found in the constitutional DNA of 26 BC patients and 74 healthy controls.

The whole *BIRC5* gene coding region and selected promoter and 3'UTR regions were investigated for selected ten polymorphisms. One of the ten polymorphisms (c.-235G>A) was not found in BC patients. Polymorphisms and their frequencies are listed in Table 3.

According to the TRANSFAC database (<http://genexplan.com/transfac>), promoter and 5'UTR region of *BIRC5* gene

TABLE 3: Comparison of minor allele frequencies of *BIRC5* polymorphisms between breast cancer patients and healthy control samples. Nucleotide changes are named according to the literature.

gene region	SNP ID number	nucleotide change	minor allele frequency controls (n/N, %)*	minor allele frequency BC samples (n/N, %)*	p (for allele frequencies)
promoter	rs3764383	c.-1547C>T**	37/148 (25.0)	16/52 (30.8)	0.466
promoter	rs8073903	c.-644T>C	49/148 (33.1)	18/52 (34.6)	0.865
promoter	rs8073069	c.-625G>C	33/148 (22.3)	13/52 (25.0)	0.704
promoter	rs17878731	c.-267G>A	1/148 (0.7)	1/52 (1.9)	0.453
promoter	rs17878467	c.-241C>T	16/148 (10.8)	6/52 (11.5)	1.000
promoter	rs17887126	c.-235G>A	2/148 (1.3)	0/52 (0)	
5'UTR	rs9904341	c.-31G>C	55/148 (37.2)	18/52 (34.6)	0.867
intron 2	rs4789551	c.221+209T>C	7/148 (4.7)	1/52 (1.9)	0.683
exon 4	rs2071214	c.9194G>A***	5/148 (3.4)	1/52 (1.9)	1.000
3'UTR	rs17885521	c.9288G>C	3/148 (2.0)	1/52 (1.9)	1.000
3'UTR	rs17882627	c.9342G>A	2/148 (1.3)	2/52 (3.8)	0.278
3'UTR	rs2239680	c.9386T>C	34/148 (23.0)	14/52 (26.9)	0.575
3'UTR	rs17882139	c.9387_9388insAA	3/148 (2.0)	2/52 (3.8)	0.606
3'UTR	rs1042489	c.9809T>C	53/148 (35.8)	18/52 (34.6)	1.000
3'UTR	rs2661694	c.10611C>A	38/148 (25.7)	11/52 (21.1)	0.578

* n – number of minor alleles, N – number of analyzed alleles
 ** C is the minor allele
 *** G is the minor allele.




FIGURE 3: Structure of *BIRC5* gene with positions of analyzed polymorphisms. Positions of known binding sites for transcription factors and miRNA are shown according to TRANSFAC database [40]. Chromosome 17 positions are numbered according to human genome build GRCh38/hg38.

contain 29 binding sites for 14 different transcription factors. Chromosome positions of the detected polymorphisms revealed that polymorphisms rs17878467 (c.-241C>T) and rs17887126 (c.-235G>A) are located in the shared binding site of transcription factors EGR1 and SP1 (Figure 3). 3'UTR

region also contains 16 binding sites for 11 different miRNAs, and only polymorphism rs1042489 (c.9809T>C) is located in the binding site of hsa-miR-708-5p (Figure 3) [40].

Fourteen different polymorphisms were found in BC samples while in controls one more polymorphism was




FIGURE 4: Pairwise linkage disequilibrium (LD) of eight *BIRC5* sequence variants with highest minor allele frequency. The location of each sequence variant along the *BIRC5* gene is relative to the real nucleotide position. The number in each diamond indicates the intensity of LD (r^2) between respective pairs of sequence variants. The LD strength is also represented by shades of grey (0 [white] < r^2 < 1 [black]). (a) Controls. (b) Cases.

present (c.-235G>A). The majority of detected polymorphisms were located either upstream or downstream of the coding region of the gene, and only one in the coding region. In addition to ten polymorphisms selected for this study, five more rare polymorphisms were found: one in the promoter (c.-267G>A), one in intron 2 (c.221+209T>C), and 3 in the 3'UTR (c.9288G>C, c.9342G>A, and c.9387_9388insAA). There was no significant difference in genotype or allele frequencies between BC and control samples.

The observed allele frequencies for most of the polymorphisms are in accordance with previously observed frequencies for either global or European populations [15, 19, 20, 22, 41–43], or in NCBI database (<http://www.ncbi.nlm.nih.gov>), and only for c.221+209T>C the frequency was lower in this study than previously observed [41, 44]. c.-235G>A polymorphism found only in healthy controls is known to create a new “alternative” binding site for the transcription factor GATA-1 and is suggested to increase the BC aggressiveness [45]. Since most of the patients in this study are still alive, it is possible that this is a reason this polymorphism was not found in the patient group.

Analysis showed there is a difference in number of polymorphisms in linkage disequilibrium (LD) between BC samples and controls. BC cases showed increased nonrandom association of alleles at multiple loci. In case of c.-644C>T and c.9809T>C, these two polymorphisms always appear in the same combination in all tested BC samples showing complete LD ($r^2=0.99$) (Figure 4). Only one pair of polymorphisms, c.9386T>C and c.10611C>A, shows decreased association in BC samples compared to control samples.

LD between various *BIRC5* polymorphisms have often been reported in different populations: some observed here,

but also others. LD between c.-1547C>T and c.9386T>C found here has been reported by Shi et al. [15], as were c.-1547C>T with c.10611C>A and c.9386T>C with c.10611C>A. The latter one was also reported by Pu et al. [46]. LD between c.-644T>C and c.-625G>C was reported twice, by Jang et al. [19] and Yang et al. [42], and between c.-625G>C and c.9809T>C once [47]. Interestingly, the polymorphisms in total LD, c.-644C>T and c.9809T>C, found in BC patients have not been described before.

3.5. Association of *BIRC5* Polymorphisms with Survivin Expression and Clinicopathological Parameters. *BIRC5* polymorphisms were not associated with survivin expression, but several polymorphisms were found to be associated with age of onset. Genotype frequencies of polymorphisms c.-644T>C, c.9386T>C, and c.9809T>C were significantly associated with age of onset, while c.-1547C>T was weakly associated. Allele frequencies of these four, as well as SNP c.-241C>T, were also associated with age of onset (Table 4). When genotype frequencies were plotted against age of onset, a clear trend could be seen for the five polymorphisms (Figure 5) (the results are presented as in Han et al.'s study [4]).

For the c.-1547C>T polymorphism, the minor allele T was associated with earlier age of onset. The same trend is visible for the c.-644C>T allele and the c.9809T>C allele, where in both cases the minor C allele is associated with earlier age of onset. For the c.9386T>C allele, the major allele T is associated with earlier age of onset. For the final, c.-241C>T polymorphism, where the difference was significant only on allele but not on genotype frequencies, the minor allele T also seems to be associated with earlier age of onset but the




FIGURE 5: Association of age of onset with the significantly associated polymorphisms. (a) $c.-1547C>T$. (b) $c.-644T>C$. (c) $c.-241C>T$. (d) $c.9386T>C$. (e) $c.9809T>C$.

number of samples in the TT group is too low to make this comparison.

Until now, only two studies showed association between *BIRC5* polymorphisms and age of onset. Han et al. [4] found significant connection of age of onset for the $c.-1547A>G$ and less significantly $c.-31G>C$ polymorphisms in ovarian cancer, while Hmeljak et al. [20] found that $c.-241C>T$ was associated with the age of diagnosis of malignant pleural mesothelioma patients, but not with total survival. These

results are comparable with our results in BC, where we show that the minor allele of the $c.1547A>G$ polymorphism is also associated with age of onset, and in our case the results for the $c.-241C>T$ are significant when comparing allele, but not for genotype frequencies. This discrepancy may be due to a small number of samples with the TT genotype (only one in our dataset). In BC, amplifications in *BIRC5* gene were also associated with the younger age of onset [48].

TABLE 4: Association of genotype and allele frequencies of polymorphisms with age of onset.

polymorphism			mean±SD (years)	p-value	
c.-1547C>T*	genotype	CC	62.3±6.5	0.054	
		CT	55.8±8.1		
		TT	49.0±10.1		
	allele	C	58.3±7.9		0.011
		T	50.9±9.8		
c.-644T>C	genotype	TT	59.4±8.2	0.006	
		TC	50.7±8.6		
		CC	39.0±2.8		
	allele	T	55.8±9.2		0.006
		C	48.1±9.1		
c.-241C>T	genotype	CC	54.7±10.0	0.143	
		CT	49.0±4.5		
		TT	37.0±0.0		
	allele	C	54.2±9.6		0.029
		T	45.0±7.1		
c.9386T>C	genotype	TT	49.1±9.3	0.029	
		TC	57.4±8.4		
		CC	62.3±6.5		
	allele	T	50.8±9.6		0.004
		C	59.5±7.6		
c.9809T>C	genotype	TT	59.4±8.2	0.006	
		TC	50.7±8.6		
		CC	39.0±2.8		
	allele	T	55.8±9.2		0.006
		C	48.1±9.1		

* C is the minor allele.

4. Conclusions

This was the first study investigating the possible role of *BIRC5* polymorphisms in breast cancer etiology conducted in Croatia. In our study no association of any of *BIRC5* polymorphism with BC or level of survivin expression was observed. High survivin expression was significantly associated with negative ER status. However, as many as five *BIRC5* polymorphisms were associated with the age of onset, a phenomenon which was previously observed only in ovarian cancer patients. Two of those five polymorphisms (c.-644T>C and c.9809T>C) were also shown to be in complete LD in breast cancer patients. We are aware that the sample size of this study was limited, but it should be noted that the allele frequencies for polymorphisms found were in accordance with previous reports, except for c.-235G>A, which was not present in BC samples. To assess the significance of *BIRC5* polymorphisms and survivin expression as predictive and prognostic biomarkers for BC further research with a larger sample size is needed.

Data Availability

This is not applicable, since there are no big datasets connected with this article.

Disclosure

Results published in this paper were presented as poster at HDIR-5: The Fifth Meeting of the Croatian Association for Cancer Research with International Participation, Zagreb in November, 2018.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Authors' Contributions

Ilona Sušac and Petar Ozretić are equally contributing authors.

Acknowledgments

This study was funded and supported by Terry Fox Foundation (Terry Fox Run Croatia 2012 and 2016) and Foundation of the Croatian Academy of Sciences and Arts (10-102/168-3-2015). The authors are grateful to Croatian League Against Cancer and the Terry Fox Foundation for their administrative and financial support. They are also extremely grateful to all participating patients, without whose help and willingness to participate this study would not be possible.

Supplementary Materials

Supplementary Table 1: PCR primer sequences and cycling conditions. (*Supplementary Materials*)

References

- [1] X. O. Shu, J. Long, W. Lu et al., "Novel genetic markers of breast cancer survival identified by a genome-wide association study," *Cancer Research*, vol. 72, no. 5, pp. 1182–1189, 2012.
- [2] T. A. Masoodi, B. Banaganapalli, V. Vaidyanathan, V. R. Talluri, and N. A. Shaik, "Computational analysis of breast cancer GWAS loci identifies the putative deleterious effect of STXBP4 and ZNF404 gene variants," *Journal of Cellular Biochemistry*, vol. 118, no. 12, pp. 4296–4307, 2017.
- [3] M. Ghousaini, P. D. Pharoah, and D. F. Easton, "Inherited genetic susceptibility to breast cancer: the beginning of the end or the end of the beginning?" *The American Journal of Pathology*, vol. 183, no. 4, pp. 1038–1051, 2013.
- [4] C. H. Han, Q. Wei, K. K. Lu, Z. Liu, G. B. Mills, and L.-E. Wang, "Polymorphisms in the survivin promoter are associated with age of onset of ovarian cancer," *International Journal of Clinical and Experimental Medicine*, vol. 2, no. 4, pp. 289–299, 2009.
- [5] D. C. Altieri, "Targeting survivin in cancer," *Cancer Letters*, vol. 332, no. 2, pp. 225–228, 2013.
- [6] P. K. Jaiswal, A. Goel, and R. D. Mittal, "Survivin: A molecular biomarker in cancer," *Indian Journal of Medical Research*, vol. 142, no. April, pp. 389–397, 2015.
- [7] D. C. Altieri, "Validating survivin as a cancer therapeutic target," *Nature Reviews Cancer*, vol. 3, no. 1, pp. 46–54, 2003.
- [8] K. Tanaka, S. Iwamoto, G. Gon, T. Nohara, M. Iwamoto, and N. Tanigawa, "Expression of survivin and its relationship to loss of

- apoptosis in breast carcinomas," *Clinical Cancer Research*, vol. 6, no. 1, pp. 127–134, 2000.
- [9] S. R. Marsicano, R. K. Kuniyoshi, F. S. Gehrke, B. C. Alves, L. A. Azzalis, and F. L. Fonseca, "Survivin expression in patients with breast cancer during chemotherapy," *Tumor Biology*, vol. 36, no. 5, pp. 3441–3445, 2015.
- [10] Y. Xu, F. Fang, G. Ludewig, G. Jones, and D. Jones, "A mutation found in the promoter region of the human survivin gene is correlated to overexpression of survivin in cancer cells," *DNA and Cell Biology*, vol. 23, no. 9, pp. 527–537, 2004.
- [11] Z.-J. Cheng, L.-H. Hu, and S.-J. Huang, "Correlation of -31G/C polymorphisms of survivin promoter to tumorigenesis of gastric carcinoma," *Chinese Journal of Cancer*, vol. 27, no. 3, pp. 258–263, 2008.
- [12] N. Kawata, N. Tsuchiya, Y. Horikawa et al., "Two survivin polymorphisms are cooperatively associated with bladder cancer susceptibility," *International Journal of Cancer*, vol. 129, no. 8, pp. 1872–1880, 2011.
- [13] Y. Hsieh, C. Tsai, C. Yeh, S. Yang, Y. Hsieh, and C. Weng, "Survivin T9809C, an SNP located in 3'-UTR, displays a correlation with the risk and clinicopathological development of hepatocellular carcinoma," *Annals of Surgical Oncology*, vol. 19, no. S3, pp. 625–633, 2012.
- [14] A. G. Antonacopoulou, K. Floratou, V. Bravou et al., "The survivin -31 Snp in human colorectal cancer correlates with survivin splice variant expression and improved overall survival," *Analytical Cellular Pathology*, vol. 33, no. 5, Article ID 673592, pp. 177–189, 2010.
- [15] H. Shi, M. Bevier, R. Johansson et al., "Prognostic impact of polymorphisms in the MYBL2 interacting genes in breast cancer," *Breast Cancer Research and Treatment*, vol. 131, no. 3, pp. 1039–1047, 2012.
- [16] M. S. Coumar, F. Tsai, J. R. Kanwar, S. Sarvagalla, and C. H. Cheung, "Treat cancers by targeting survivin: Just a dream or future reality?" *Cancer Treatment Reviews*, vol. 39, no. 7, pp. 802–811, 2013.
- [17] M. L. Cvok, M. Cretnik, V. Musani, P. Ozretic, and S. Levanat, "New sequence variants in BRCA1 and BRCA2 genes detected by high-resolution melting analysis in an elderly healthy female population in Croatia," *Clinical Chemistry and Laboratory Medicine*, vol. 46, no. 10, 2008.
- [18] C. Xu, M. Yamamoto-Ibusuki, Y. Yamamoto et al., "High survivin mRNA expression is a predictor of poor prognosis in breast cancer: a comparative study at the mRNA and protein level," *Breast Cancer*, vol. 21, no. 4, pp. 482–490, 2014.
- [19] J. S. Jang, K. M. Kim, K. H. Kang et al., "Polymorphisms in the survivin gene and the risk of lung cancer," *Lung Cancer*, vol. 60, no. 1, pp. 31–39, 2008.
- [20] J. Hmeljak, N. Erčulj, V. Dolžan, I. Kern, and A. Cör, "BIRC5 promoter SNPs do not affect nuclear survivin expression and survival of malignant pleural mesothelioma patients," *Journal of Cancer Research and Clinical Oncology*, vol. 137, no. 11, pp. 1641–1651, 2011.
- [21] X. Wang, L. Huang, Y. Xu et al., "Association between survivin -31G > C promoter polymorphism and cancer risk: a meta-analysis," *European Journal of Human Genetics*, vol. 20, no. 7, pp. 790–795, 2012.
- [22] E. Aynaci, E. Coskunpinar, A. Eren et al., "Association between survivin gene promoter -31G/C and -644C/T polymorphisms and non-small cell lung cancer," *Genetics and Molecular Research*, vol. 12, no. 3, pp. 3975–3982, 2013.
- [23] Y. Y. Shi and L. He, "SHEsis, a powerful software platform for analyses of linkage disequilibrium, haplotype construction, and genetic association at polymorphism loci," *Cell Research*, vol. 15, no. 2, pp. 97–98, 2005.
- [24] W. Foulkes, "Size surprise? Tumour size, nodal status, and outcome after breast cancer," *Current Oncology*, vol. 19, no. 5, pp. 241–243, 2012.
- [25] E. Orang, E. T. Marzony, and A. Afsharfard, "Predictive role of tumor size in breast cancer with axillary lymph node involvement - can size of primary tumor be used to omit an unnecessary axillary lymph node dissection?" *Asian Pacific Journal of Cancer Prevention*, vol. 14, no. 2, pp. 717–722, 2013.
- [26] N. A. Soliman and S. M. Yussif, "Ki-67 as a prognostic marker according to breast cancer molecular subtype," *Cancer Biology & Medicine*, vol. 13, no. 4, pp. 496–504, 2016.
- [27] D. C. Allred, J. M. Harvey, M. Berardo, and G. M. Clark, "Prognostic and predictive factors in breast cancer by immunohistochemical analysis," *Modern Pathology*, vol. 11, no. 2, pp. 155–168, 1998.
- [28] M. Adamkov, E. Halasova, K. Kajo et al., "Survivin: a promising biomarker in breast carcinoma," *Neoplasma*, vol. 57, no. 6, pp. 572–577, 2010.
- [29] M. Adamkov, K. Kajo, D. Vybohova, J. Krajcovic, F. Stuller, and J. Rajcani, "Correlations of survivin expression with clinicomorphological parameters and hormonal receptor status in breast ductal carcinoma," *Neoplasma*, vol. 59, no. 01, pp. 30–37, 2012.
- [30] M. Adamkov, D. Výbohová, J. Horáček, M. Kovalská, and M. Furjelová, "Survivin expression in breast lobular carcinoma: Correlations with normal breast tissue and clinicomorphological parameters," *Acta Histochemica*, vol. 115, no. 5, pp. 412–417, 2013.
- [31] M. Sarti, S. Pinton, C. Limoni et al., "Differential expression of testin and survivin in breast cancer subtypes," *Oncology Reports*, vol. 30, no. 2, pp. 824–832, 2013.
- [32] S. M. Kennedy, L. O'Driscoll, R. Purcell et al., "Prognostic importance of survivin in breast cancer," *British Journal of Cancer*, vol. 88, no. 7, pp. 1077–1083, 2003.
- [33] L. Kleinberg, V. A. Flørenes, J. M. Nesland, and B. Davidson, "Survivin, a member of the inhibitors of apoptosis family, is down-regulated in breast carcinoma effusions," *American Journal of Clinical Pathology*, vol. 128, no. 3, pp. 389–397, 2007.
- [34] N. S. Youssef, I. H. Hewedi, and M. N. Abd Raboh, "Immunohistochemical expression of survivin in breast carcinoma: relationship with clinicopathological parameters, proliferation and molecular classification," *Journal of the Egyptian National Cancer Institute*, vol. 20, no. 4, pp. 348–357, 2008.
- [35] P. N. Span, F. C. Sweep, E. T. Wiegerinck et al., "Survivin is an independent prognostic marker for risk stratification of breast cancer patients," *Clinical Chemistry*, vol. 50, no. 11, pp. 1986–1993, 2004.
- [36] S. Yamashita, Y. Masuda, T. Kurizaki et al., "Survivin expression predicts early recurrence in early-stage breast cancer," *Anti-cancer Research*, vol. 27, no. 4C, pp. 2803–2808, 2007.
- [37] Y. Zhao, Y. Wang, X. Ni et al., "Clinical significance of Smac and survivin expression in breast cancer patients treated with anthracycline-based neoadjuvant chemotherapy," *Molecular Medicine Reports*, vol. 9, no. 2, pp. 614–620, 2014.
- [38] Q. Lin, Y. Liu, H. Chen et al., "Survivin, Ki-67 and tumor grade as predictors of response to docetaxel-based neoadjuvant chemotherapy in locally advanced breast cancer," *Molecular and Clinical Oncology*, vol. 1, no. 5, pp. 839–844, 2013.

- [39] N. Barnes, P. Haywood, P. Flint, W. F. Knox, and N. J. Bundred, "Survivin expression in in situ and invasive breast cancer relates to COX-2 expression and DCIS recurrence," *British Journal of Cancer*, vol. 94, no. 2, pp. 253–258, 2006.
- [40] E. Wingender, "The TRANSFAC project as an example of framework technology that supports the analysis of genomic regulation," *Briefings in Bioinformatics*, vol. 9, no. 4, pp. 326–332, 2008.
- [41] J. Dai, G. Jin, J. Dong et al., "Prognostic significance of survivin polymorphisms on non-small cell lung cancer survival," *Journal of Thoracic Oncology*, vol. 5, no. 11, pp. 1748–1754, 2010.
- [42] X. Yang, G. Xiong, X. Chen et al., "Polymorphisms of survivin promoter are associated with risk of esophageal squamous cell carcinoma," *Journal of Cancer Research and Clinical Oncology*, vol. 135, no. 10, pp. 1341–1349, 2009.
- [43] R. Boidot, F. Végran, D. Jacob et al., "The expression of BIRC5 is correlated with loss of specific chromosomal regions in breast carcinomas," *Genes, Chromosomes and Cancer*, vol. 47, no. 4, pp. 299–308, 2008.
- [44] D. Casabonne, O. Reina, Y. Benavente et al., "Single nucleotide polymorphisms of matrix metalloproteinase 9 (MMP9) and tumor protein 73 (TP73) interact with Epstein-Barr virus in chronic lymphocytic leukemia: results from the European case-control study EpiLymph," *Haematologica*, vol. 96, no. 2, pp. 323–327, 2011.
- [45] R. Boidot, F. Végran, D. Jacob et al., "The transcription factor GATA-1 is overexpressed in breast carcinomas and contributes to survivin upregulation via a promoter polymorphism," *Oncogene*, vol. 29, no. 17, pp. 2577–2584, 2010.
- [46] F. Pu, Z. Shao, S. Yang et al., "Association between functional variants in BIRC5/survivin gene 3' untranslated region and mRNA expression in lymphoblastoid cell lines," *Oncology Letters*, vol. 10, no. 4, pp. 2319–2322, 2015.
- [47] C. Li, Z. Li, M. Zhu et al., "Clinicopathological and prognostic significance of survivin over-expression in patients with esophageal squamous cell carcinoma: a meta-analysis," *PLoS ONE*, vol. 7, no. 9, Article ID e44764, 2012.
- [48] K. Ghaffari, M. Hashemi, E. Ebrahimi, and R. Shirkoohi, "BIRC5 genomic copy number variation in early-onset breast cancer," *Iranian Biomedical Journal*, vol. 20, no. 4, pp. 241–245, 2016.



Hindawi

Submit your manuscripts at
www.hindawi.com

