

# 2022: The Year in Cardiovascular Disease - The Year of Upfront Lipid Lowering Combination Therapy.

---

Banach, Maciej; Reiner, Željko; Cicero, Arrigo; Sabouret, Pierre; Viigimaa, Margus; Sahebkar, Amirhossein; Postadzhiyan, Arman; Gaita, Dan; Pella, Daniel; Penson, Peter

Source / Izvornik: **Archives of Medical Science, 2022, 18, 1429 - 1434**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.5114/aoms/156147>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:105:073614>

Rights / Prava: [Attribution-NonCommercial-ShareAlike 4.0 International/Imenovanje-Nekomercijalno-Dijeli pod istim uvjetima 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-09-01**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



## 2022: the year in cardiovascular disease – the year of upfront lipid lowering combination therapy

Maciej Banach<sup>1,2,3</sup>, Zeljko Reiner<sup>4</sup>, Arrigo F.G. Cicero<sup>5,6</sup>, Pierre Sabouret<sup>7</sup>, Margus Viigimaa<sup>8,9</sup>, Amirhossein Sahebkar<sup>10, 11,12,13</sup>, Arman Postadzhiyan<sup>14</sup>, Dan Gaita<sup>15,16</sup>, Daniel Pella<sup>17</sup>, Peter E. Penson<sup>18,19</sup> on behalf of *the International Lipid Expert Panel (ILEP)*

<sup>1</sup>Department of Preventive Cardiology and Lipidology, Medical University of Lodz (MUL), Lodz, Poland

<sup>2</sup>Cardiovascular Research Centre, University of Zielona Gora, Zielona Gora, Poland

<sup>3</sup>Department of Cardiology and Congenital Diseases of Adults, Polish Mother's Memorial Hospital Research Institute (PMMHRI), Lodz, Poland

<sup>4</sup>Department of Internal Medicine, University Hospital Centre Zagreb, Zagreb, Croatia

<sup>5</sup>IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

<sup>6</sup>Hypertension and Cardiovascular Risk Factors Research Center, Medical and Surgical Sciences Department, Sant'Orsola-Malpighi University Hospital, Bologna, Italy

<sup>7</sup>Heart Institute and Action Group, Pitié-Salpêtrière, Sorbonne University Paris, Paris, France

<sup>8</sup>Department of Health Technologies, Tallinn University of Technology, Tallinn, Estonia

<sup>9</sup>The North Estonia Medical Centre, Tallinn, Estonia

<sup>10</sup>Applied Biomedical Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>11</sup>Biotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>12</sup>School of Medicine, The University of Western Australia, Perth, Australia

<sup>13</sup>Department of Biotechnology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>14</sup>Department of General Medicine, Emergency University Hospital "St. Anna", Medical University of Sofia, Sofia, Bulgaria

<sup>15</sup>Universitatea de Medicina si Farmacie Victor Babes din Timisoara, Romania

<sup>16</sup>Clinica de Cardiologie, Institutul de Boli Cardiovasculare Timisoara, Romania

<sup>17</sup>2<sup>nd</sup> Department of Cardiology of the East Slovak Institute of Cardiovascular Disease and Faculty of Medicine PJ Safarik University, Kosice, Slovak Republic

<sup>18</sup>Clinical Pharmacy and Therapeutics Research Group, School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Liverpool, UK

<sup>19</sup>Liverpool Centre for Cardiovascular Science, Liverpool, UK

### Corresponding author:

Prof. Maciej Banach  
MD, PhD, FNLA, FAHA  
FESC, FASA  
Department of  
Preventive Cardiology  
and Lipidology  
Medical University  
of Lodz (MUL)  
Rzgowska 281/289  
93-338 Lodz, Poland  
Phone: +48 42 639 37 71  
Fax: +48 42 639 37 71  
E-mail: maciej.banach@  
umed.lodz.pl

**Submitted:** 25 October 2022; **Accepted:** 28 October 2022

**Online publication:** 7 November 2022

Arch Med Sci 2022; 18 (6): 1429–1434

DOI: <https://doi.org/10.5114/aoms/156147>

Copyright © 2022 Termedia & Banach

Recent data clearly show that we are extremely ineffective in the use of lipid-lowering therapy (LLT). Furthermore, the problem seems to be the most serious in the group of patients at very high and extremely high-risk – the patients who stand to benefit most from effective therapy [1]. The results of the DaVinci study showed that the proportion of very high-risk patients reaching the low-density lipoprotein cholesterol (LDL-C) goal (< 55 mg/dl; 1.4 mmol/l) is only 18% in Europe [2], and only 13% in Central and Eastern European (CEE) countries [3]. Only a very small percentage of patients at extremely high-risk, reach the goal of < 40 mg/dl (1 mmol/l) [2–5]. The SANTORINI study, which was performed in 2020–2021, showed a small improvement of 2.7% over the DaVinci study which was completed in 2019 [6]. However, still only 20.7% of atherosclerotic cardiovascular disease (ASCVD) patients achieved their LDL-C goal. This study also clearly showed that these disappointing findings are associated with a very poor therapeutical approach in


patients with cerebral ASCVD and peripheral artery disease (PAD) – with only 15 and 18.6% reaching LDL-C goals, respectively [6]. The latter group present a real unmet need in most of the departments of cardiology in Europe, as PAD is very rarely diagnosed – despite the recognition of a very strong link between PAD and ASCVD and cardiovascular events. Unfortunately, the majority of these patients are now diagnosed and treated by the interventional surgeons [7, 8]. The results of the REALITY cohort study, performed in 2019 based on the nationwide database with real-life Spanish patients, confirmed that only about 3% of individuals achieve the LDL-C target of less than 55 mg/dl (1.4 mmol/l) and less than 15% of ASCVD patients reach the goal of < 70 mg/dl (1.8 mmol/l) [9]. This study again confirmed that the largest unmet needs refer to patients with stroke/transient ischemic attack (TIA) and PAD, for whom the 24-month mortality rate was between 10.2–11.9% [9–11].

The main reason for such a large extent of underperformance is the lack of use of high intensity statin (HIS) therapy. Less than 25% patients received HIS therapy in the SANTORINI study [6]. The corresponding figure was 32% in CEE countries and 38% in the Europe based on the DaVinci study [2, 3]. Comparing this data with the results of the Hyperlipidaemia Therapy in tERTiary Cardiological cEnTer (TERCET) Registry for ACS patients, which includes data up to 2016, it appears that there has been no improvement, as 6 years ago we observed the HIS ratio of 37.4% and 39.1% in patients with STEMI and NSTEMI, respectively [12]. Even more dramatic examples of underuse of effective therapies are seen for patients treated with the combination therapy of statin and ezetimibe, and double/triple therapy with PCSK9 inhibitors [2–6, 13, 14]. In the TERCET registry the proportion of patients on the combination therapy with ezetimibe was only 0.3% of patients with myocardial infarction (MI), and 3.5% in those with stable coronary artery disease [12]. In the DaVinci study, 9% of ASCVD patients were treated with combination therapy, and in the SANTORINI study it was 17.5% (and only 8.5% and 12% in those with cerebral ASCVD and PAD, respectively) [2, 3, 6]. There are still very few patients (only 4.7%) receiving combination therapy with PCSK9 inhibitors [6] (in the DaVinci it was only 1% [1]). This represents an enormous under-use of therapy considering the needs of patients. Based on modelling studies, a PCSK9 targeted approach therapy is expected to be administered in 20–22% of patients to achieve LDL-C goals. This means that presently, only one fifth of patients benefits from these innovative therapies [15, 16].

This relatively small improvement in the number of patients being treated with LLT combination ther-

apy has not been accompanied by an improvement in the proportion of patients achieving LDL-C goals [17]. There are several reasons for this. The first is obviously associated with too few patients being treated with combination therapy – and in particular, insufficient use of upfront combination LLT. This is despite our International Lipid Expert Panel (ILEP) recommendations on upfront combination therapy, which were published in April 2021 [14], and the reiteration of the advice in the European Atherosclerosis Society (EAS) Task Force paper [18] as well as an expert opinion paper published in EHJ [19], and at least several national guidelines in which this approach was approved (e.g., in Poland, Spain, France) [7, 20–22]. Another reason for limited goal achievement is related to the fact that when non-statin drug(s) are added to statin therapy, the statin dose is reduced at the same time in many patients, and/or ezetimibe is discontinued in case of triple therapy with a PCSK9 targeted therapy approach. This was clearly demonstrated in the GOULD Registry, where after 2 years only 17.1% patients had LLT intensification, and in the PCSK9 inhibitors group there was LLT de-escalation in 18.6% of patients, and statin downtitration/discontinuation and ezetimibe discontinuation in 10.2% [23]. The same trend was observed in an analysis of 1499 patients included to the Managed Care for Acute Myocardial Infarction Survivors (MACAMIS; “KOS-Zawat”) program in Poland, where after 12 months it was observed that LLT was deescalated in 23.5% and intensified in only 9.2% of patients after ACS [24]. That is why all forthcoming recommendations should focus not only on when and how to intensify LLT, but to focus on intensification of each step (preferably with the use of FDC with the highest tolerated statin doses), and without discontinuation of the ezetimibe, which has become an unfortunate trend for many physicians treating patients with a PCSK9 targeted therapy approach. The results of the ODYSSEY APPRISE (and many other available reports) clearly showed that in patients treated with alirocumab, keeping patients on high intensity statin therapy and ezetimibe was associated with significantly larger number of patients reaching LDL-C goals (by as much as 3.5% and 13.4%, respectively) [5]. Therefore, to treat our patients effectively, we should always retain high-intensity statin therapy while adding ezetimibe, and HIS and ezetimibe while adding PCSK9 inhibitors. This approach would lead to an expected 80–85% reduction of LDL-C in comparison to the 65–70% seen in most current real-life observations [25, 26] (Figure 1).

Obviously, downtitration is possible in the case of statin-associated muscle symptoms (SAMS)/statin intolerance (SI). But in most cohort- and observational studies SI is overdiagnosed – which report prevalence to be as high as 30–40%. In the



**Figure 1.** How to be effective with lipid lowering therapy in ASCVD patients (the size of the LDL-C reduction for some recommended combinations is an assumption and still needs to be confirmed)

\*FDC of high intensity statin therapy and ezetimibe is a preferable option for all ASCVD patients. ASCVD – atherosclerotic cardiovascular disease, LDL-C – low-density lipoprotein cholesterol, ACS – acute coronary syndrome, PAD – peripheral artery disease, HIS – high intensity statin therapy, FDC – fixed dose combination, EZE – ezetimibe, BA – bempedoic acid, PCSK9i – proprotein convertase subtilisin/kexin 9 targeted approach therapy (PCSK9 inhibitors + inclisiran).

TERCET registry, worryingly, 9.3% of ACS patients received no statin therapy (and a much higher proportion received only low-to moderate intensity statin therapy) [12]. In the DaVinci study 6% received no statin, and additionally 46% only received low to moderate doses of statins in monotherapy) [2]. In the SANTORINI study, as many as 21.4% of ASCVD patients received no statin [6]. It is worth emphasizing that the true prevalence of SI is only < 7% when it is diagnosed using approved definitions [27]. After the exclusion of the placebo effect [28, 29] (using an approach which considers patient education [28], by exclusion of conditions and risk factors that might increase of the SI risk and the use of the SAMS-Clinical Index Score), complete statin intolerance is observed in only about 2% of treated patients [30, 31]. Therefore, to provide effective treatment, it is not enough to use effective drugs at high doses, we must also carefully and effectively manage SI [32].

Finally, we also need to have clear and unanimous guidelines. Unfortunately, the European Society of Cardiology (ESC) prevention guidelines published in September 2021 recommended a stepwise approach to achieve LDL-C goal for ASCVD patients [33]. Meanwhile, evidence-based medicine (EBM) supports the paradigm of “the lower the better for longer” and “the earlier on LDL-C goal, the better” [34], which therefore requires intensive treatment as early as possible. The stepwise guidelines promote an approach, which was applied in most departments of cardiology about 15–20 years ago – and which is a main reason of very small ratio of ASCVD patients reaching LDL-C goals (irrespective what goal was based on the applicable guidelines). The

consequence of missed goals is the high number of recurrent CVD events and mortality (which may be as high as 20% in the first 12 months after ACS) [35, 36]. The reevaluation of the risk, looking again at comorbidities and considering patients’ preferences in those at the very high and extremely high CVD risk (e.g. in those after stroke or ACS), were, based on the authors’ guidelines, intended to improve the effectiveness of therapy [33]. However, the guidelines introduced considerable confusion in practice because they promoted the opposite approach to the ESC/EAS lipid guidelines published in 2019 [37], and contrasted with the upfront combination therapy approach, which has been widely used in last 2 years [28]. This may be the reason that fewer and fewer patients will achieve LDL-C goals (e.g., those who, based on these guidelines express a preference against further LLT intensification). Alternatively, the goal be met, but achievement of the target will be delayed (against the rule – ‘the earlier the better’).

Taking this opportunity, it is also worth at least briefly discussing at least 3 issues, that are raised by supporters of the stepwise approach:

1. *Cost-effectiveness.* It is said that the stepwise approach is more cost-effective. However, this claim cannot be substantiated because the approach essentially prolongs the time to achievement of the LDL-C target, as well as reduces the number of patients achieving the goal at all (see below). Abundant data indicates the direct link between unachieved therapeutic goals, and increased risk of CVD events (and therefore long-term costs). This does not even require a health technology assessment (HTA) analysis. An approach allowing earlier achievement and better maintenance

- of the LDL-C goals (assuming better adherence with the fixed doses combination (FDC) therapy, and earlier introduction of PCSK9 targeted approach) is the best way to reduce the risk of recurrent CVD events (and their consequences – costs of the interventional procedures, costs of heart failure therapy, sick leave/disability, and rehabilitation) and mortality [38, 39].
2. *Potential risk in the elderly and in frailty patients (over 75 years of age).* We now have convincing data suggesting that intensive LLT therapy in elderly patients with ASCVD (especially after an event) is as beneficial in this group as it is for younger groups [7, 40]. Additionally, an approach involving the administration of the upfront LLT with intensive statin therapy (but not necessarily at the highest-possible doses – atorvastatin 40 mg and rosuvastatin 20 mg) with ezetimibe as FDC may not only decrease the risk of possible SAMS (in comparison to HIS with the highest doses) [27] but may also increase adherence and the number of patients reaching LDL-C targets. Clinical practice clearly shows us that in most of these patients, statin therapy and combination therapy with statin (even in moderate intensity doses) and ezetimibe is sufficient for most of these patients to reach LDL-C goals. This is probably due to physiologically lower baseline LDL-C levels in comparison to younger groups [7, 37]. Obviously individualization of the therapy for this group of patients is highly recommended.
  3. *Lack of data confirming that the upfront lipid lowering therapy is effective, especially in the context of the CVD events reduction.* Since the introduction of ezetimibe in 2002 (and in Poland since 2006) we have had numerous data suggesting that combination therapy (also as FDC) is significantly more effective in comparison to maximally tolerated statin therapy and/or to double the dose of statin [7, 12]. Additionally, high intensity statin therapy is associated with approximately 50% reduction of LDL-C, whereas for the combination therapy the expected reduction is as high as 65% [7, 37]. In a systematic review and meta-analysis of 13 studies with 5080 patients, the authors showed a significantly greater percentage reduction in LDL-C levels in patients treated with ezetimibe-statin vs. statin monotherapy (by -14.1%,  $p < 0.001$ ) [41]. Reduction of LDL-C levels attributed to add-on ezetimibe was significantly greater than that for statin dose doubling (by -15.3%,  $p < 0.001$ ), and LDL-C goal achievement favoured add-on ezetimibe over statin titration by as much as 2.5 times ( $p = 0.007$ ) [41]. In a pooled analysis of 27 clinical trials with 21,000 subjects therapy with ezetimibe plus statin produced significantly greater reductions in LDL-C, total-cholesterol, non-high density lipoprotein cholesterol (HDL-C), apolipoprotein (Apo)B, triglycerides, and significantly higher achievement of LDL-C  $< 70$  mg/dl by 18.2%, and  $< 100$  mg/dl by 23.4% ( $p < 0.0001$  for all) [42]. The same results were observed for both coronary heart disease patients (LDL-C  $< 70$  mg/dl was achieved more frequently in statin/ezetimibe group by 21.8%), and in diabetic patients ( $< 100$  mg/dl by 20.2%) [42]. Katzmann *et al.* studied prescription trends in oral non-statin LLT and their effects on LDL-C in Germany [43]. Data from 311,242 patients were analysed. They confirmed that addition of ezetimibe in patients already prescribed a statin reduced LDL-C by an additional 23.8% ( $32.3 \pm 38.4$  mg/dl), with a greater reduction with FDC (reduction 28.4% ( $40.0 \pm 39.1$  mg/dl)) as compared to separate pills (19.4% ( $27.5 \pm 33.8$  mg/dl));  $p < 0.0001$ . Despite a small proportion of patients reaching the recommended LDL-C level of  $< 70$  mg/dl, a greater proportion of patients treated with FDC reached the goal (31.5%) compared to those with separate pills (21.0%) [43]. In a randomised, open-label, non-inferiority trial, 3780 ASCVD patients were randomly assigned to receive either moderate-intensity statin with ezetimibe combination therapy (rosuvastatin 10 mg with ezetimibe 10 mg;  $n = 1894$ ) or high-intensity statin monotherapy (rosuvastatin 20 mg;  $n = 1886$ ) [44]. The primary endpoint was the 3-year composite of cardiovascular death, major cardiovascular events, or non-fatal stroke. The primary endpoint occurred in 172 (9.1%) patients in the combination therapy group and 186 (9.9%) patients in the high-intensity statin monotherapy group (absolute difference -0.78%; 95% CI: -2.39–0.83). LDL-C concentrations of  $< 70$  mg/dl at 1, 2, and 3 years were observed in 73%, 75%, and 72% of patients in the combination therapy group, and 55%, 60%, and 58% of patients in the high-intensity statin monotherapy group (all  $p < 0.0001$ ). The combination therapy was also significantly better tolerated – discontinuation or dose reduction of the study drug by intolerance was observed in 88 (4.8%) patients and 150 (8.2%) patients, respectively ( $p < 0.0001$ ) [44]. Finally, based on data from 38,023 consecutive patients with ACS from the Polish Registry of Acute Coronary Syndromes (PL-ACS) we showed that there was a significant difference between statin monotherapy (atorvastatin or rosuvastatin) and upfront combination therapy of statin and ezetimibe after 1 year (5.9 vs. 3.6%,  $p = 0.049$ ), 2 (7.8 vs. 4.4%,  $p = 0.023$ ) and 3 years (10.2 vs. 5.6%,  $p = 0.03$ ) of follow-up (as well as for the overall period) – in favor for the upfront combination therapy [45]. Upfront combination therapy was associated with a significant reduction



of all-cause mortality in comparison to statin monotherapy (OR = 0.490, 95% CI: 0.351–0.683) with absolute risk reduction (ARR) of 4.6% after 3 years (number needed to treat – NNT = 22) [45].

In conclusion, in recent decades we have made many mistakes in the management of lipid disorders, from completely ignoring this very common risk factor, to incorrect risk stratification, late diagnosis, and ineffective treatment with low doses and without the use of combination therapy [46–49]. In 2022, we have so many effective drugs (including new agents close on the horizon), and effective and scientifically confirmed approaches to treatment. Therefore, we simply cannot afford to lose our patients due to our own ineffectiveness and therapeutic inertia [50]. So, starting from now #ThinkUpfront!

### Conflict of interest

Maciej Banach: speakers bureau: Amgen, Daiichi Sankyo, KRKA, Polpharma, Mylan/Viatrix, Novartis, Novo-Nordisk, Sanofi-Aventis, Teva, Zentiva; consultant to Amgen, Daiichi Sankyo, Esperion, NewAmsterdam, Novartis, Novo-Nordisk, Polfarmex, Sanofi-Aventis; Grants from Amgen, Daiichi Sankyo, Mylan/Viatrix, Sanofi and Valeant; Zeljko Reiner: speakers bureau: Sanofi, Novartis; Margus Viigimaa: speakers bureau: Amgen, Novartis, Novo-Nordisk, Sanofi-Aventis, Menarini; Arman Postadzhyan: speakers bureau: Amgen, Novartis, Novo-Nordisk, Teva, Zentiva; Daniel Pella: received honoraria for events sponsored by Amgen, Jamieson, Novartis, MSD, Pfizer, Servier; Peter E. Penson owns four shares in AstraZeneca PLC and has received honoraria and/or travel reimbursement for events sponsored by AKCEA, Amgen, AMRYT, Link Medical, Mylan, Napp, Sanofi; all other authors have no conflict of interest.

### References

- Sabouret P, Angoulvant D, Cannon CP, Banach M. Low levels of low-density lipoprotein cholesterol, intracerebral haemorrhage, and other safety issues: is there still a matter of debate? *Eur Heart J Open* 2022; 2: oeac038.
- Ray KK, Molemans B, Schoonen WM, et al.; DA VINCI study. EU-Wide cross-sectional observational study of lipid-modifying therapy use in secondary and primary care: the DA VINCI study. *Eur J Prev Cardiol* 2021; 28: 1279-89.
- Vrablik M, Seifert B, Parkhomenko A, et al. Lipid-lowering therapy use in primary and secondary care in Central and Eastern Europe: DA VINCI observational study. *Atherosclerosis* 2021; 334: 66-75.
- Lyall DM, Ward J, Banach M, et al. PCSK9 genetic variants and cognitive abilities: a large-scale Mendelian randomization study. *Arch Med Sci* 2021; 17: 241-4.
- Banach M, López-Sendon JL, Averna M, et al. Treatment adherence and effect of concurrent statin intensity on the efficacy and safety of alirocumab in a real-life setting: results from ODYSSEY APPRISE. *Arch Med Sci* 2022; 18: 285-92.
- Catapano AL, Manu MC, Burden A, Ray KK; the SANTORINI Investigators. LDL-C goal achievement and lipid-lowering therapy in patients by atherosclerotic cardiovascular disease subtype: the SANTORINI study. *Eur Heart J* 2022; 43 Suppl\_2: ehac544.2373.
- Banach M, Burchardt P, Chlebun K, et al. PoLA/CFPiP/PCS/PSLD/PSD/PSH guidelines on diagnosis and therapy of lipid disorders in Poland 2021. *Arch Med Sci* 2021; 17: 1447-547.
- Sahebkar A, Mikhailidis DP, Banach M. CETP inhibition in the REVEAL trial: could we aim higher? *Arch Med Sci* 2020; 16: 1229-30.
- Campuzano R, Barrios V, Mostaza JM, et al. Preliminary results from REALITY: a nation-wide study of a database with 1.8 million real-life patients to study atherosclerotic cardiovascular disease and familial hypercholesterolemia in Spain. *Eur Heart J* 2022; 43 Suppl\_2: ehac544.2672.
- Banach M, Shekoohi N, Mikhailidis DP, Lip GYH, Hernandez AV, Mazidi M. Relationship between low-density lipoprotein cholesterol, lipid-lowering agents and risk of stroke: a meta-analysis of observational studies (n = 355,591) and randomized controlled trials (n = 165,988). *Arch Med Sci* 2022; 18: 912-29.
- Patsourakos NG, Kouvari M, Kotidis A, et al. The incidence of recurrent cardiovascular events among acute coronary syndrome patients treated with generic or original clopidogrel in relation to their sociodemographic and clinical characteristics. The Aegean study. *Arch Med Sci* 2020; 16: 1013-21.
- Dyrbus K, Gasior M, Desperak P, Nowak J, Osadnik T, Banach M. Characteristics of lipid profile and effectiveness of management of dyslipidaemia in patients with acute coronary syndromes – data from the TERCET registry with 19,287 patients. *Pharmacol Res* 2019; 139: 460-6.
- Ugovšek S, Zupan J, Rehberger Likozar A, Šebeštjen M. Influence of lipid-lowering drugs on inflammation: what is yet to be done? *Arch Med Sci* 2022; 18: 855-69.
- Banach M, Penson PE, Vrablik M, et al.; ACS EuroPath Central & South European Countries Project. Optimal use of lipid-lowering therapy after acute coronary syndromes: A Position Paper endorsed by the International Lipid Expert Panel (ILEP). *Pharmacol Res* 2021; 166: 105499.
- Cannon CP, Khan I, Klimchak AC, Reynolds MR, Sanchez RJ, Sasiela WJ. Simulation of lipid-lowering therapy intensification in a population with atherosclerotic cardiovascular disease. *JAMA Cardiol* 2017; 2: 959-66.
- Momtazi-Borojeni AA, Jaafari MR, Afshar M, Banach M, Sahebkar A. PCSK9 immunization using nanoliposomes: preventive efficacy against hypercholesterolemia and atherosclerosis. *Arch Med Sci* 2021; 17: 1365-77.
- Banach M, Penson PE. Lipid-lowering therapies: better together. *Atherosclerosis* 2021; 320: 86-8.
- Averna M, Banach M, Bruckert E, et al. Practical guidance for combination lipid-modifying therapy in high- and very-high-risk patients: a statement from a European Atherosclerosis Society Task Force. *Atherosclerosis* 2021; 325: 99-109.
- Ray KK, Reeskamp LF, Laufs U, et al. Combination lipid-lowering therapy as first-line strategy in very high-risk patients. *Eur Heart J* 2022; 43: 830-3.
- Sabouret P, Lemesle G, Bellemain-Appaix A, et al. Post-discharge and long-term follow-up after an acute coronary syndrome: International Collaborative Group of CNCF position paper. *Arch Med Sci* 2022; 18: 839-54.

21. Solnica B, Sygitowicz G, Sitkiewicz D, et al. 2020 Guidelines of the Polish Society of Laboratory Diagnostics (PSLD) and the Polish Lipid Association (PoLA) on laboratory diagnostics of lipid metabolism disorders. *Arch Med Sci* 2020; 16: 237-52.
22. Barrios V, Escobar C, Banach M. Primary prevention. The cornerstone to reduce the burden of cardiovascular disease. *Rev Esp Cardiol* 2021; 74: 827-8.
23. Cannon CP, de Lemos JA, Rosenson RS, et al.; GOULD Investigators. Use of lipid-lowering therapies over 2 years in GOULD, a registry of patients with atherosclerotic cardiovascular disease in the US. *JAMA Cardiol* 2021; 6: 1-9.
24. Gasior M, et al. Dyslipidemia prevalence, treatment, and attainment of LDL-cholesterol treatment goals in patients participating in the Managed Care for Acute Myocardial Infarction Survivors. *Kardiol Pol* 2022; (in press).
25. Lewek J, Konopka A, Starostecka E, Penson PE, Maciejewski M, Banach M. Clinical features of familial hypercholesterolemia in children and adults in EAS-FHSC Regional Center for rare diseases in Poland. *J Clin Med* 2021; 10: 4302.
26. Vicente-Valor J, García-González X, Ibáñez-García S, et al. PCSK9 inhibitors revisited: effectiveness and safety of PCSK9 inhibitors in a real-life Spanish cohort. *Biomed Pharmacother* 2022; 146: 112519.
27. Bytyçi I, Penson PE, Mikhailidis DP, et al. Prevalence of statin intolerance: a meta-analysis. *Eur Heart J* 2022; 43: 3213-23.
28. Penson PE, Bruckert E, Marais D, et al.; International Lipid Expert Panel (ILEP). Step-by-step diagnosis and management of the nocebo/drucebo effect in statin-associated muscle symptoms patients: a position paper from the International Lipid Expert Panel (ILEP). *J Cachexia Sarcopenia Muscle* 2022; 13: 1596-622.
29. Banach M, Penson PE. Drucebo effect – the challenge we should all definitely face! *Arch Med Sci* 2021; 17: 542-43.
30. Banach M. Statin intolerance: time to stop letting it get in the way of treating patients. *Lancet* 2022; 400: 791-3.
31. Serban MC, Colantonio LD, Manthripragada AD, et al. Statin intolerance and risk of coronary heart events and all-cause mortality following myocardial infarction. *J Am Coll Cardiol* 2017; 69: 1386-95.
32. Banach M, Cannon CP, Paneni F, Penson PE; endorsed by the International Lipid Expert Panel (ILEP). Individualized therapy in statin intolerance: the key to success. *Eur Heart J* 2022; doi: 10.1093/eurheartj/ehac556.
33. Visseren FLJ, Mach F, Smulders YM, et al.; ESC National Cardiac Societies; ESC Scientific Document Group. 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* 2021; 42: 3227-337.
34. Banach M, Penson PE. Statins and LDL-C in secondary prevention-so much progress, so far to go. *JAMA Netw Open* 2020; 3: e2025675.
35. Piątek Ł, Wilczek K, Kurzawski J, et al. Gender-related disparities in the treatment and outcomes in patients with non-ST-segment elevation myocardial infarction: results from the Polish Registry of Acute Coronary Syndromes (PL-ACS) in the years 2012-2014. *Arch Med Sci* 2020; 16: 781-8.
36. Zandecki L, Janion M, Sadowski M, et al. Associations of changes in patient characteristics and management with decrease in mortality rates of men and women with ST-elevation myocardial infarction – a propensity score-matched analysis. *Arch Med Sci* 2020; 16: 772-80.
37. Mach F, Baigent C, Catapano AL, et al.; ESC Scientific Document Group. 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J* 2020; 41: 111-88.
38. Sasidharan A, Bagepally BS, Kumar SS, Jagadeesh KV, Natarajan M. Cost-effectiveness of ezetimibe plus statin lipid-lowering therapy: a systematic review and meta-analysis of cost-utility studies. *PLoS One* 2022; 17: e0264563.
39. Sharrock T. The cost-effectiveness of fixed-dose combinations for preventive cardiovascular pharmacotherapy. A thesis for the Degree of Masters of Public Health; University of Otago, New Zealand, 2018, 1-104.
40. Awad K, Mohammed M, Zaki MM, et al.; Lipid and Blood Pressure Meta-analysis Collaboration (LBPMC) Group and the International Lipid Expert Panel (ILEP). Association of statin use in older people primary prevention group with risk of cardiovascular events and mortality: a systematic review and meta-analysis of observational studies. *BMC Med* 2021; 19: 139.
41. Mikhailidis DP, Lawson RW, McCormick AL, et al. Comparative efficacy of the addition of ezetimibe to statin vs statin titration in patients with hypercholesterolaemia: systematic review and meta-analysis. *Curr Med Res Opin* 2011; 27: 1191-210.
42. Morrone D, Weintraub WS, Toth PP, et al. Lipid-altering efficacy of ezetimibe plus statin and statin monotherapy and identification of factors associated with treatment response: a pooled analysis of over 21,000 subjects from 27 clinical trials. *Atherosclerosis* 2012; 223: 251-61.
43. Katzmann JL, Sorio-Vilela F, Dornstauder E, et al. Non-statin lipid-lowering therapy over time in very-high-risk patients: effectiveness of fixed-dose statin/ezetimibe compared to separate pill combination on LDL-C. *Clin Res Cardiol* 2022; 111: 243-52.
44. Kim BK, Hong SJ, Lee YJ, et al.; RACING investigators. Long-term efficacy and safety of moderate-intensity statin with ezetimibe combination therapy versus high-intensity statin monotherapy in patients with atherosclerotic cardiovascular disease (RACING): a randomised, open-label, non-inferiority trial. *Lancet* 2022; 400: 380-90.
45. Lewek J, Niedziela J, Desperak D, et al. Intensive statin therapy versus upfront combination therapy of statin and ezetimibe in acute coronary syndrome patients. A propensity score matching analysis based on the PL-ACS Registry data. *Eur Heart J* (data unpublished, in review).
46. Fras Z, Tršan J, Banach M. On the present and future role of Lp-PLA<sub>2</sub> in atherosclerosis-related cardiovascular risk prediction and management. *Arch Med Sci* 2021; 17: 954-64.
47. Shakour N, Ruscica M, Hadzadeh F, et al. Statins and C-reactive protein: in silico evidence on direct interaction. *Arch Med Sci* 2020; 16: 1432-9.
48. Penson PE, Pirro M, Banach M. LDL-C: lower is better for longer-even at low risk. *BMC Med* 2020; 18: 320.
49. Banach M, Stulc T, Dent R, Toth PP. Statin non-adherence and residual cardiovascular risk: there is need for substantial improvement. *Int J Cardiol* 2016; 225: 184-96.
50. Banach M, Kaźmierczak J, Mitkowski P, et al. Which patients at risk of cardiovascular disease might benefit the most from inclisiran? Polish experts' opinion. The compromise between EBM and possibilities in health-care. *Arch Med Sci* 2022; 18: 569-76.