

Immune effector cell-associated hematotoxicity: EHA/EBMT consensus grading and best practice recommendations

Rejeski, Kai; Subklewe, Marion; Aljurf, Mahmoud; Bachy, Emmanuel; Balduzzi, Adriana; Barba, Pere; Bruno, Benedetto; Benjamin, Reuben; Carrabba, Matteo G.; Chabannon, Christian; ...

Source / Izvornik: **Blood**, 2023, 142, 865 - 877

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1182/blood.2023020578>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:656065>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International](#)/[Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)


Download date / Datum preuzimanja: **2024-05-26**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine](#)
[Digital Repository](#)





Immune effector cell–associated hematotoxicity: EHA/EBMT consensus grading and best practice recommendations

Kai Rejeski,^{1,*} Marion Subklewe,^{1,*} Mahmoud Aljurf,² Emmanuel Bachy,³ Adriana Balduzzi,⁴ Pere Barba,^{5,6} Benedetto Bruno,⁷ Reuben Benjamin,⁸ Matteo G. Carrabba,⁹ Christian Chabannon,¹⁰ Fabio Ciceri,⁹ Paolo Corradini,¹¹ Julio Delgado,¹² Roberta Di Blasi,¹³ Raffaella Greco,⁹ Roch Houot,¹⁴ Gloria Iacoboni,^{5,6} Ulrich Jäger,¹⁵ Marie José Kersten,¹⁶ Stephan Mielke,¹⁷ Arnon Nagler,¹⁸ Francesco Onida,¹⁹ Zinaida Peric,²⁰ Claire Roddie,²¹ Annalisa Ruggeri,⁹ Fermín Sánchez-Guijo,²² Isabel Sánchez-Ortega,²³ Dominik Schneidawind,²⁴ Maria-Luisa Schubert,²⁵ John A. Snowden,²⁶ Catherine Thieblemont,¹³ Max Topp,²⁷ Pier Luigi Zinzani,²⁸ John G. Gribben,²⁹ Chiara Bonini,³⁰ Anna Sureda,³¹ and Ibrahim Yakoub-Agha³²

¹Department of Medicine III, LMU University Hospital, LMU Munich, Munich, Germany; ²Oncology Center, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia; ³Department of Hematology, Hospices Civils de Lyon and Université Claude Bernard Lyon 1, Lyon, France; ⁴Pediatric Transplantation Unit, Department of Medicine and Surgery, University of Milan-Bicocca-Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; ⁵Department of Hematology, Vall d'Hebron University Hospital, Experimental Hematology, Vall d'Hebron Institute of Oncology, Barcelona, Spain; ⁶Department of Medicine, Universitat Autònoma de Barcelona, Bellaterra, Spain; ⁷Division of Hematology and Cell Therapy Unit, Department of Molecular Biotechnology and Health Sciences, University of Turin, Turin, Italy; ⁸School of Cancer & Pharmaceutical Sciences, King's College London, London, United Kingdom; ⁹Unit of Hematology and Bone Marrow Transplantation, IRCCS San Raffaele Hospital, Milan, Italy; ¹⁰Institut Paoli-Calmettes Comprehensive Cancer Centre and Module Biothérapies du Centre d'Investigations Cliniques de Marseille, INSERM-Aix-Marseille Université-AP-HM-IPC, CBT-1409, Marseille, France; ¹¹Division of Hematology and Stem Cell Transplantation, Fondazione IRCCS Istituto Nazionale dei Tumori, University of Milan, Milan, Italy; ¹²Oncoimmunotherapy Unit, Department of Hematology, Hospital Clinic de Barcelona, Barcelona, Spain; ¹³Université de Paris, Assistance Publique-Hôpitaux de Paris, Service d'Hématologie, Paris, France; ¹⁴Department of Hematology, CHU Rennes, University of Rennes, INSERM U1236, Rennes, France; ¹⁵Department of Medicine I, Medical University of Vienna, Vienna, Austria; ¹⁶Department of Hematology, Amsterdam University Medical Centers, Amsterdam, The Netherlands; ¹⁷Department of Cellular Therapy and Allogeneic Stem Cell Transplantation, Department of Laboratory Medicine and Medicine Huddinge, Karolinska University Hospital and Institute, Stockholm, Sweden; ¹⁸Division of Hematology, Chaim Sheba Medical Center, Tel Aviv University, Tel-Hashomer, Israel; ¹⁹Hematology and Bone Marrow Transplantation Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, University of Milan, Milan, Italy; ²⁰Department of Hematology, University Hospital Centre Zagreb, Zagreb, Croatia; ²¹Department of Hematology, University College London Hospital, London, United Kingdom; ²²University of Salamanca, IBSAL-University Hospital of Salamanca, Salamanca, Spain; ²³Executive Office, European Society for Blood and Marrow Transplantation, Barcelona, Spain; ²⁴Department of Medical Oncology and Hematology, University Hospital Zurich, Zurich, Switzerland; ²⁵Department of Medicine V, University Hospital Heidelberg, Heidelberg, Germany; ²⁶Department of Haematology, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, United Kingdom; ²⁷Medizinische Klinik und Poliklinik II, Universitätsklinikum Würzburg, Würzburg, Germany; ²⁸IRCCS Azienda Ospedaliero-Universitaria di Bologna, Istituto di Ematologia "Seràgnoli," Dipartimento di Medicina Specialistica, Diagnostica e Sperimentale, Università di Bologna, Bologna, Italy; ²⁹Barts Cancer Institute, Queen Mary, University of London, London, United Kingdom; ³⁰Division of Immunology, Transplantation and Infectious Disease, Experimental Hematology Unit, IRCCS San Raffaele Hospital, Vita-Salute San Raffaele University, Milan, Italy; ³¹Clinical Hematology Department, Institut Català d'Oncologia-L'Hospitalet, Barcelona, Spain; and ³²CHU de Lille, University of Lille, INSERM U1286, Lille, France

Hematological toxicity is the most common adverse event after chimeric antigen receptor (CAR) T-cell therapy. Cytopenias can be profound and long-lasting and can predispose for severe infectious complications. In a recent worldwide survey, we demonstrated that there remains considerable heterogeneity in regard to current practice patterns. Here, we sought to build consensus on the grading and management of immune effector cell–associated hematotoxicity (ICAHT) after CAR T-cell therapy. For this purpose, a joint effort between the European Society for Blood and Marrow Transplantation (EBMT) and the European Hematology Association (EHA) involved an international panel of 36 CAR T-cell experts who met in a series of virtual conferences, culminating in a 2-day meeting in Lille, France. On the basis of these deliberations, best practice recommendations were developed. For the grading of ICAHT, a classification

system based on depth and duration of neutropenia was developed for early (day 0-30) and late (after day +30) cytopenia. Detailed recommendations on risk factors, available preinfusion scoring systems (eg, CAR-HEMA-TOTOX score), and diagnostic workup are provided. A further section focuses on identifying hemophagocytosis in the context of severe hematotoxicity. Finally, we review current evidence and provide consensus recommendations for the management of ICAHT, including growth factor support, anti-infectious prophylaxis, transfusions, autologous hematopoietic stem cell boost, and allogeneic hematopoietic cell transplantation. In conclusion, we propose ICAHT as a novel toxicity category after immune effector cell therapy, provide a framework for its grading, review literature on risk factors, and outline expert recommendations for the diagnostic workup and short- and long-term management.

Introduction and state of the art

The last decade has firmly established chimeric antigen receptor (CAR) T-cell therapy as a practice-changing immunotherapy platform for an increasing number of refractory B-cell malignancies.¹⁻⁷ Although durable remissions can be achieved, this comes with the caveat of a unique spectrum of side effects ranging from cytokine release syndrome (CRS), to immune effector cell-associated neurotoxicity syndrome (ICANS), and immune effector cell-associated hemophagocytic lymphohistiocytosis-like syndrome (IEC-HS).⁸⁻¹¹ Real-world evidence has underlined the growing importance of hematological toxicity as the most frequent Common Terminology Criteria for Adverse Events (CTCAE) grade ≥ 3 adverse event after CAR T-cell therapy.¹²⁻¹⁴ Similarly high rates of cytopenias have been reported for other T-cell-based immunotherapies such as bispecific antibodies.¹⁵⁻¹⁹ Notably, profound and often long-lasting cytopenias can add to the immunosuppression conferred by B-cell aplasia and consecutive hypogammaglobulinemia.²⁰ Importantly, severe infections are a major driver of both morbidity and nonrelapse mortality following CAR T-cell therapies.²¹⁻²³

Hematological side effects have been described after CAR T-cell therapy regardless of the target antigen (eg, CD19 vs CD22 vs B-cell maturation antigen [BCMA]) and across various disease entities (eg, large B-cell lymphoma [LBCL], B-cell precursor acute lymphoblastic leukemia [BCP-ALL], mantle cell lymphoma [MCL], multiple myeloma [MM], and follicular lymphoma [FL]).^{3-5,24-29} Several features underline the unique nature of CAR T-cell-related hematotoxicity. First, cytopenias can persist long after the resolution of clinical CRS, and have been reported as long as months to years after CAR T-cell infusion.³⁰ Hematopoietic count recovery often follows a biphasic trajectory, with intermittent recovery followed by second, or multiple, dips.^{12,13} Second, patients can develop very severe bone marrow (BM) aplasia that is often refractory to therapeutic measures such as growth factor support.^{13,31,32} Finally, the underlying pathophysiology remains to be elucidated, although recent evidence points toward the importance of both baseline hematopoietic reserve and the systemic inflammatory state of the host.¹³ Moreover, the inflammatory stress conferred by severe CRS and the associated alterations in cytokine patterns can exert myelosuppressive effects.³³⁻³⁵

In a recent international survey led by the European Hematology Association (EHA) and European Society for Blood and Marrow Transplantation (EBMT), we identified a high degree of heterogeneity in regard to both the grading and management of cytopenias.³⁶ Current grading systems, such as the CTCAE, describe cytopenias predominantly in quantitative terms by assigning severity grades based on the depth of cytopenia. However, they are not clinically actionable and fail to capture the distinct nature of post-CAR T-cell hematopoietic reconstitution, such as the biphasic and/or delayed course. Furthermore, the cumulative risk of secondary complications (eg, infections or bleeding) primarily increases with the respective duration of observed cytopenia.^{22,37} Classification systems that were developed for cytopenia following classic cytotoxic chemotherapies may not apply to patients receiving novel T-cell-based immunotherapies. To accommodate these unique features of hematological side effects in adult patients receiving such therapies, we herein introduce the concept of immune

effector cell-associated hematotoxicity (ICAHT). Based on a novel framework for grading, we outline expert recommendations for its diagnostic workup and management.

Methodology

This workshop is based on the EBMT Practice Harmonization and Guidelines committee method.³⁸ In September 2022, K.R. and M.S. proposed to set up a workshop to issue European recommendations regarding the grading and management of ICAHT, particularly after autologous CAR T-cell therapy. As a first step, an international survey on current practices at >50 global CAR T-cell therapy centers was sent out and results were analyzed.³⁶ Experts from different countries and belonging to EBMT and EHA were subsequently invited to join the workshop. As a second step, several teleconferences took place to discuss and advance the first draft. Along with the results of the international survey, a comprehensive literature review was carried out by the workshop participants within each subgroup, which served as the basis for the discussions. The third step consisted of a 2-day face-to-face meeting, which took place in Lille, France in March 2023.

These recommendations are intended to be general in scope and applicable to all diseases and types of CAR T-cell therapies or other T-cell-based immunotherapies (eg, bispecific antibody constructs) adopted as standard clinical practice. They are intended to reflect current best practices in this new and rapidly evolving field and aim to help clinicians and other health care professionals in providing consistent, high-quality patient care. These recommendations were created because of the growing number of autologous CAR T-cell therapies currently available outside of clinical trials for the treatment of hematological malignancies. Given the lack of high-quality evidence from randomized trials in this area (expected evidence levels 3-5, Oxford Centre for Evidence Based Medicine), the decision was made not to grade these recommendations. They therefore represent the consensus point of view of the authors. When administering CAR T-cell therapies within clinical trials, physicians are advised to follow respective trial protocols.

Consensus recommendations

ICAHT grading

On the basis of the results of the international survey on behalf of EHA and EBMT, the expert panel defined early ICAHT as cytopenia occurring during the first 30 days after CAR T-cell infusion. Conversely, late ICAHT was classified as cytopenia observed beyond day +30. The expert panel resolved that the main clinical action points of post-CAR T-cell cytopenias concerned profound and/or prolonged neutropenia, and that isolated thrombocytopenia or anemia represent rare occurrences. Concomitantly, a grading system based on neutropenia was pursued. For early ICAHT (day 0-30), a grading system based on both depth and duration of neutropenia was defined due to the associated clinical sequelae (Table 1, top). Late ICAHT was graded based on the elapsed time from CAR T-cell infusion (eg, occurring after day +30) with the severity (grade 1-4) defined by the depth of neutropenia (Table 1, bottom). For anemia and thrombocytopenia, the expert panel refers to existing grading systems and recommends that institutional guidelines should

Table 1. ICAHT grading

Grading	1	2	3	4
Early ICAHT (day 0-30)				
ANC $\leq 500/\mu\text{L}$	<7 d	7-13 d	≥ 14 d	Never above $500/\mu\text{L}$
ANC $\leq 100/\mu\text{L}$	—	—	≥ 7 d	≥ 14 d
Late ICAHT (after day +30)*				
ANC	$\leq 1500/\mu\text{L}$	$\leq 1000/\mu\text{L}$	$\leq 500/\mu\text{L}$	$\leq 100/\mu\text{L}$

*Measured ≥ 2 time points, or nontransient neutropenia.

be followed, as further outlined in “Management of cytopenias” and Table 3 (refer to “Transfusions”).

Risk factors for developing post-CAR T-cell cytopenias

The overall incidence of hematological toxicity in the key registrational trials for CAR T-cell products endorsed by the European Medicines Agency are outlined in the supplemental Table 1, available on the *Blood* website. Furthermore, we performed an extensive literature review of prominent real-world studies with a specific focus on correlative studies and potential risk factors (supplemental Table 2). Overall, a plethora of factors contribute to the development of cytopenias after CAR T-cell therapy, some of

which remain incompletely understood. Broadly, they relate to the underlying disease and its previous treatments, baseline risk factors (eg, hematopoietic reserve, BM infiltration, and systemic inflammation), as well as CAR T-cell product features and CRS-related inflammatory patterns (summarized in Table 2 and the supplemental Material).^{12,13,23,30,33,34,39-56}

What scoring systems to use

Based on several of the risk factors delineated earlier, the CAR-HEMATOTOX score was developed to identify patients at high risk for prolonged neutropenia, and especially the development of the aplastic phenotype of neutrophil recovery.¹³ An online calculator can be found on the website of the German

Table 2. Risk factors associated with an increased risk of post-CAR T-cell cytopenias

	Risk factors	Comments	References
Disease-related features	Underlying disease (BCP-ALL > B-NHL) Disease burden prior to CAR T-cell infusion (progressive disease, high LDH)	Evidence concerning the rate of cytopenias in patients with MM still emerging Especially BM disease burden	39 14,102
Prior therapies	Number of prior therapy lines Prior hematopoietic stem cell transplantation Bridging therapy	Associated with baseline hematopoietic function	39 104 41
Baseline marrow status	BM infiltration Preexisting cytopenias Clonal hematopoiesis of indeterminate potential (ChIP)?	Particularly preexisting thrombocytopenia Has been linked to increased inflammation, potential emerging risk factor	42,40 13,33 46,47,85
Baseline inflammatory status	Increased serum CRP Increased serum ferritin		13 13
CAR T-cell product and postinfusion risk factors	Costimulatory molecule (CD28 > 41BB) Type of construct (tandem > single target) Severe CRS Sustained increased inflammatory markers Oligoclonal T-cell expansion Active infection CRS/MAS or IEC-HS	May also reflect differences in lymphodepletion dosing (cyclophosphamide dosing) In select patients; the success of autologous stem cell boost argues against this as a general mechanism Mainly viral or in case of concomitant sepsis Cytopenia as overlapping symptomology	39 39 33,34 33 35 105 10,11,106

BCP-ALL, B-cell precursor acute lymphoblastic leukemia; B-NHL, B-cell non-Hodgkin lymphoma; CRP, C-reactive protein; LDH, lactate dehydrogenase; MAS, macrophage activation syndrome.

Lymphoma Alliance (<https://www.german-lymphoma-alliance.de/Scores.html>). The score incorporates factors related to hematopoietic reserve (absolute neutrophil count [ANC], hemoglobin, and platelet count) and baseline inflammatory state (C-reactive protein and ferritin) and was validated for a primary end point of severe neutropenia (ANC < 500/ μ L) lasting ≥ 14 days during the first 60 days after CAR T-cell infusion. Importantly, the CAR-HEMATOTOX score is determined before lymphodepleting chemotherapy and thus enables early risk-stratification into a high vs low risk of developing severe hematotoxicity after CAR T-cell treatment (Figure 1). In subsequent studies, the score also identified patients at risk for severe infections and poor treatment outcomes across multiple disease entities (eg, LBCL, MCL, and MM).^{22,42-44,57} However, it is important to note that the score remains to be validated prospectively and for adult and pediatric patients with B-cell precursor acute lymphoblastic leukemia. Furthermore, the test characteristics (high sensitivity, but lower specificity) indicate a lower positive predictive value, meaning that not all patients deemed high risk will develop severe hematotoxicity. Conversely, the high negative predictive value suggests that the score is particularly helpful in ruling out patients at risk for severe hematotoxicity.

Assessment and diagnostic workup of ICAHT

In patients with a high-risk profile for developing ICAHT (Table 2; Figure 1), baseline BM studies (prior to apheresis or lymphodepletion) should be considered to risk-stratify patients for hematological toxicity and to identify underlying marrow infiltration as a pertinent risk factor. Cryopreservation of the BM aspirate and/or peripheral blood mononuclear cells is optional but may provide useful information in case the patient develops secondary BM failure (eg, presence of CHiP clone).

In case of cytopenia that persists beyond the expected reconstitution of lymphodepleting chemotherapy (typically after week 2-3 after CAR T-cell infusion), the first step in the workup lies in defining the differential diagnosis, which can include

drug-induced cytopenia, vitamin deficiencies, infectious causes, sustained inflammatory stressors, relapse, and/or active BM disease. The expert panel recommends performing an incremental diagnostic workup, with an initial tier 1 assessment comprising standard diagnostic tests that should be performed in all cases of severe, or grade ≥ 3 , ICAHT (Figure 2). In case the tier 1 results are inconclusive and cytopenias persist and/or are granulocyte colony-stimulating factor (G-CSF) refractory (absence of count recovery despite ≥ 5 days of G-CSF support), a subsequent tier 2 diagnostic workup can be pursued. Importantly, this includes extended viral studies, as well as BM aspiration and biopsy. The expert panel would reserve cytogenetics and next-generation sequencing to rule out an underlying myeloid malignancy to either cases of profound, long-lasting marrow aplasia (eg, no count recovery above an ANC of ≥ 500 / μ L by day +30, pancytopenia), or new-onset pancytopenia that is refractory to therapeutic measures late after CAR T-cell infusion.

Hemophagocytosis associated with severe hematotoxicity after CAR T-cell therapy

Hemophagocytic lymphohistiocytosis (HLH) is a hyper-inflammatory condition resulting from abnormal immune activation, which is associated with high fever, hyperferritinemia, prolonged cytopenia, and eventually multiorgan failure. HLH remains a diagnostic quandary because unique biomarkers are still lacking and/or not readily available. In the context of CAR T-cell therapy, the incidence of HLH-like symptoms ranges from 1% to 3.4%.^{10,58} Two entities, CRS/MAS and IEC-HS, can be distinguished based on time of onset and presence of concomitant CRS/ICANS symptoms.^{11,29,59,60} In patients with severe ICAHT that present with aplastic neutrophil recovery and rising serum ferritin, the diagnosis of HLH should be considered as both can present with profound immune dysregulation and increased interferon signaling.^{42,54} A comprehensive workup is recommended in order to identify additional abnormalities such as new-onset hepatosplenomegaly, hypertriglyceridemia, coagulopathy, and hypofibrinogenemia, as well as hemophagocytosis features on BM biopsy or in other tissues (Figure 2). Existing




Figure 1. The CAR-HEMATOTOX score as a risk stratification tool. Data presented in the tables are based on multicenter retrospective analyses examining hematotoxicity and infectious complications in patients receiving CAR-T for relapsed/refractory LBCL,^{13,22} MCL,⁴³ or MM.⁴⁴ CI, confidence interval; HT, CAR-HEMATOTOX score; MCL, mantle cell lymphoma; MM, multiple myeloma.

	Categories	Putative causes	Test	Time points	Comments
TIER 1	Lower threshold to perform – minimal workup				
	Poor bone marrow reserve	Prior treatments including allo-HCT, fludarabine, marrow infiltration	Complete blood count (CBC), reticulocyte production index (RPI), peripheral blood smear	Routinely	Recommended
	Medication – drug side effects	Check for concomitant myelosuppressive medications		Routinely	
	Vitamin deficiencies	Vitamin B12, folic acid	Serum levels	Routinely	Recommended
	Rule out infections	Bacterial/viral/fungal infections	Blood cultures, CMV PCR, procalcitonin, CD4 ⁺ T-cell, IgG, B-cell levels	Routinely	Recommended
	Rule out macrophage-activation syndrome*	CRS/MAS or IEC-HS	Serum ferritin, triglycerides	Routinely	Recommended
TIER 2	Subsequent work-up – In case of G-CSF refractory state, if tier 1 results are negative and/or risk factors are present				
	Viral PCR considering the clinical presentation	Parvovirus	Parvovirus B19 PCR	In case of prolonged anemia	Recommended
		HHV6, JCV	HHV6, JCV PCR blood/CSF	In case of neurologic symptoms	Recommended
		EBV, adenovirus, HSV	PCR	In case of HLH	Recommended
	Bone marrow disease	(MDS/AML/myelofibrosis) or relapse	BM aspirate, biopsy, flow cytometry, immunohistochemistry, cytogenetics, NGS	In case of prolonged cytopenia	Recommended
		Relapse of leukemia/lymphoma	Flow cytometry peripheral blood / bone marrow, including B-cell panel	Routinely	Recommended
	Other causes	Other rare hematologic diseases, myeloid diseases, PNH, autoimmune processes	Myeloid panel, GPI-linked structures, direct antiglobulin test (DAT)	In case of suspected MPN/ PNH/autoimmune processes	Recommended

Figure 2. Step-by-step diagnostic workup depending on ICAHT severity. *In case of elevated ferritin and clinical suspicion of MAS, refer to supplemental Table 3 and supplemental Figure 1. AML, acute myeloid leukemia; CMV, cytomegalovirus; CRS/MAS, cytokine release syndrome with macrophage activation syndrome; EBV, Epstein-Barr virus; GPI, glycosylphosphatidylinositol; HHV6, human herpesvirus 6; HSV, herpes simplex virus; IgG, immunoglobulin G; JCV, JC virus; MDS, myelodysplastic syndrome; MPN, myeloproliferative neoplasm; NGS, next-generation sequencing; PCR, polymerase chain reaction; PNH, paroxysmal nocturnal hemoglobinuria.

scoring systems that can guide the diagnosis of HLH in the context of severe ICAHT include HLH-2004 criteria, the H-score, and the optimized HLH inflammatory index.^{61–63} Furthermore, supplemental Table 3 outlines the MD Anderson criteria,⁵⁸ EBMT/EHA recommendations,⁵⁹ and IEC-HS criteria,¹¹ which were deemed more specific to CAR T-cell therapy by the expert panel. In patients in whom ICAHT manifests in the form of HLH, anti-inflammatory measures should be promptly initiated to mitigate cytokine storm and its clinical sequelae. Patients should be treated with anakinra, a recombinant humanized interleukin-1 receptor antagonist, in combination with high-dose corticosteroids (supplemental Figure 1). In refractory cases, ruxolitinib, cytokine adsorption, and emapalumab (an interferon- γ inhibitor) can be considered, albeit data remain scarce.^{64–66}

Management of cytopenias

The management of ICAHT can broadly be separated into an initial phase, which addresses the (expected) early cytopenias and aims to mitigate the risk of infections and/or other complications, as well as a later phase that is initiated in case of persistent and/or therapy-refractory cytopenias. An overview of the expert recommendations for early ICAHT management is provided in Table 3.

Transfusions

Because of the frequent nature of severe anemia and thrombocytopenia after CAR T-cell therapy, transfusions are an essential part of supportive care and include either packed red blood cell concentrates or platelet concentrates.

Transfusion-associated graft-versus-host disease (ta-GVHD) is a rare complication of transfusion wherein viable donor T lymphocytes in cellular blood products mount an immune response against the recipient.⁶⁷ Considering the high mortality rate (>90%), prevention of ta-GVHD is recommended, although there is no internationally agreed upon consensus on the duration of the use of irradiated blood products across cellular therapies. In the setting of hematopoietic cell transplantation (HCT), standard practice is to use irradiated blood for (1) at least 2 weeks before stem cell collection until at least 3 months after auto-HCT, and (2) at the start of conditioning at the latest until at least 6 months after allogeneic HCT (allo-HCT), or until immune reconstitution.⁶⁸ In the context of CAR T-cell therapy, the expert panel recommended the irradiation of blood products from 7 days before leukapheresis until at least 90 days after CAR T-cell infusion unless conditioning, disease, or previous treatment determine indefinite duration (Table 3). Of note, the use of the purine analog fludarabine as a component of lymphodepletion before CAR T-cell infusion may affect local guidance for irradiated blood products.⁶⁸ Given its relative rarity, we recommend reporting cases of ta-GVHD after CAR T-cell therapy to regulatory authorities.

Growth factor support

GM-CSF Granulocyte-macrophage colony-stimulating factor (GM-CSF) is typically elevated in patients with CRS and ICANS receiving CAR T-cell therapy. The use of GM-CSF as a growth factor for patients with low blood counts should be avoided as it may promote inflammatory toxicity and induce neuroinflammation after CAR T-cell therapy.^{69,70}

Table 3. Short-term management of cytopenias

	When	How	Precautions	Comments
pRBC/platelet transfusions	As per institutional standards, based on patient risk profile	As per institutional standards For pRBC: consider using 1 product per time to reduce iron overload ⁶⁸	Irradiation of blood products; start 7 d before leukapheresis until at least 90 d after CAR T-cell infusion	Because of the use of fludarabine
G-CSF	Prophylactic G-CSF: on day +2 in patients with a high-risk profile for ICAHT (eg, high CAR-HEMATOTOX score and risk profile per Table 2) Therapeutic G-CSF: severe neutropenia (ANC < 500/ μ L) neutropenia with or without infectious complications	Based on individual risk profile: consider early G-CSF administration (from day +2) as prophylaxis in those at high risk for ICAHT Dosing: 5 μ g/kg once daily In case of prolonged neutropenia with/without infectious complications Dosing: 5 μ g/kg once daily, consider increasing dose in case of nonresponse	In patients at low risk for ICAHT, G-CSF probably not necessary	Reduced risk of febrile neutropenia (without increasing the risk of severe, or grade ≥ 3 , CRS nor ICANS) No detrimental effect on CAR T-cell expansion kinetics or treatment outcomes ^{73,74} Patients with intermittent neutrophil recovery often rapidly respond to G-CSF stimulation, whereas those who are aplastic are often G-CSF unresponsive
Antibacterial prophylaxis	In patients with a low risk for ICAHT, not recommended In patients with a high-risk profile for ICAHT, prophylaxis may be considered once ANC is <500/ μ L	As per institutional standards (eg, levofloxacin or ciprofloxacin)	Warning in case of colonization by MDR pathogens	Look at local bacterial epidemiology. High local prevalence of MDR GNB might prevent the use of antibacterial prophylaxis.
Antiviral	All patients	Start from LD conditioning until 1 y after CAR T-cell infusion AND/OR until CD4 ⁺ count is >0.2 $\times 10^9$ /L Valaciclovir 500 mg twice a day or acyclovir 800 mg twice a day		
Antipneumocystis	All patients	Start from LD conditioning until 1 y after CAR T-cell infusion AND/OR until CD4 ⁺ count is >0.2 $\times 10^9$ /L Co-trimoxazole 480 mg once daily or 960 mg 3 times each week	In case of co-trimoxazole allergy, pentamidine inhalation (300 mg once every month), dapsone 100 mg daily or atovaquone 1500 mg once daily can be considered	Can be started later depending on center guidelines
Systemic primary antifungal prophylaxis	Prophylaxis may be considered in case of severe neutropenia (ANC < 500/ μ L) and a high-risk profile for ICAHT (eg, CAR-HEMATOTOX score and risk profile per Table 2) and/or prolonged neutropenia	Mold-active prophylaxis for 1-3 mo (depending on the duration of neutropenia and use of steroids): posaconazole (300 mg/d) or micafungin (50 mg per day, IV)		In patients with prior allo-HCT, prior invasive aspergillosis, and those receiving corticosteroids (long-term >72 h, or high-dose), prophylaxis is recommended

LD, lymphodepletion; MDR GNB, multidrug-resistant gram-negative bacteria; pRBC, packed red blood cell.

G-CSF Because of the concerns for the use of GM-CSF and the hypothesized, but largely unknown, risks of exacerbating toxicities, early guidance suggested generally deferring G-CSF until resolution of acute CAR T-cell-related immunotoxicity (typically week 3). However, several recent reports question this as a general rule and point toward an acceptable safety profile for the early use of G-CSF, with no increase of high-grade (grade ≥ 3) CRS/ICANS.⁷¹⁻⁷⁵ In the largest retrospective analysis by Miller et al (n = 197), prophylactic G-CSF before CAR T-cell therapy (mostly pegylated G-CSF) was associated with faster neutrophil recovery, comparable treatment outcomes, and similar rates of severe ICANS.⁷⁴ Although prophylactic G-CSF was associated with a higher rate of grade ≥ 2 CRS, this observation did not

extend to the clinically relevant grade ≥ 3 CRS. In a subgroup analysis, the authors found that G-CSF did not worsen severity of CRS in patients who already present with low-grade (grade 1) toxicity. In a further study by Lievin et al, early G-CSF administration (from day +2) in patients with neutropenia was associated with a reduced risk of febrile neutropenia without increasing the risk of severe CRS or ICANS.⁷³ Notably, G-CSF was also safe in maintaining CAR T-cell expansion kinetics and antilymphoma activity, without any deleterious impact on the quality of response and outcomes.^{72,73} Appraising the aforementioned evidence and weighing the benefits and risks, early G-CSF administration on day +2 can be considered in high-risk patients to shorten the length of expected severe neutropenia (see

Table 2; Figure 1). Therapeutic G-CSF in case of prolonged severe neutropenia ($\text{ANC} < 500/\mu\text{L}$) can also be considered and can be of diagnostic benefit for identifying the aplastic neutrophil recovery phenotype,^{13,32} which is often G-CSF unresponsive. The large majority (>80%) of patients receiving CAR T-cell therapy ultimately respond to growth factor support with count recovery.^{32,34} However, recurrent neutrophil dips (biphasic course) can necessitate intermittent application of therapeutic G-CSF (Figure 3). Finally, a uniform consensus was reached on the necessity of prospective, and ideally multicenter, clinical trials that evaluate the safety and optimal treatment protocol for G-CSF (prophylactic vs early, and pegylated vs nonpegylated) in the context of CAR T-cell therapy and across disease entities (BCP-ALL vs B-NHL vs MM).

TPO agonists

Thrombopoietin (TPO) agonists (eg, eltrombopag and romiplostim) are considered primarily in patients with prolonged and late thrombocytopenia, with the thrombocytopenic nadir typically occurring in the second month after CAR T-cell therapy.^{12,13} Data supporting the use of TPO agonists in the CAR T-cell setting are extremely limited and are restricted to a few case series from single centers with limited patient numbers.⁷⁶⁻⁷⁸ In these limited reports, improvement in platelets and also

hemoglobin and ANC was noted, with some patients becoming transfusion independent both for platelets and packed red blood cell concentrates, similar to improvement in hematopoiesis observed with TPO agonist use in cases of acquired BM failure.^{79,80} Because of the limited available data, the expert panel advises that the use of TPO agonists should parallel the practice for HCT.⁸¹ They can also be used in G-CSF-refractory cases of ICAHT (Figure 3).

Infection prophylaxis

Regarding the administration of anti-infectious prophylaxis during cytopenia, the expert panel broadly recommends adherence to the general EHA/EBMT guidelines for patients receiving CAR T-cell therapy.⁵⁹ The following specific recommendations were issued (Table 3):

- Adherence to current EHA/EBMT guidelines regarding antiviral and antipneumocystis pneumonia prophylaxis, as well as intravenous immunoglobulin substitution for post-CAR T-cell therapy hypogammaglobulinemia.⁵⁹
- The expert panel does not recommend the use of a neutropenic diet to reduce the risk of infection in patients with neutropenia receiving CAR T-cell therapy.⁸²⁻⁸⁴




Figure 3. Treatment algorithm for immune effector cell-associated hematotoxicity. *High risk defined as previous history of hematopoietic stem cell transplantation, baseline cytopenia, high tumor burden, systemic inflammation, and presence of BM infiltration. **Antifungal prophylaxis particularly recommended in patients with previous invasive fungal disease, previous allo-HCT, and receiving corticosteroids (long-term >72 hour or high dose). Decision for/against antibacterial prophylaxis should incorporate local bacterial epidemiology (eg, prevalence for multidrug-resistant gram-negative bacteria); not recommended for patients with a low-risk profile for ICAHT. †Also extends to late ICAHT if these criteria are met.

- Antibacterial prophylaxis: the panel proposes a risk-adapted strategy based on the patient-individual risk profile for infections including the expected incidence rate of protracted, profound neutropenia (ANC of $<100/\mu\text{L}$ for ≥ 7 days), in line with the consensus American Society of Clinical Oncology/Infectious Diseases Society of America (ASCO/IDSA) recommendations for adult patients with cancer.⁸⁵ Antibacterial prophylaxis with a fluoroquinolone (eg, levofloxacin or ciprofloxacin) is not recommended in patients who are at low risk of severe (grade ≥ 3) ICAHT (Tables 1 and 3) and should be avoided because of fluoroquinolone-specific side effects, the potential emergence of resistant strains, and selection for *Clostridium difficile* and enterococci.^{37,86-89} Furthermore, recent publications have demonstrated that antibiotic exposure before CAR T-cell therapy reduces microbiome diversity and is associated with inferior outcomes, potentially because of the multifunctional and immunomodulatory role of the gut microbiome.⁹⁰⁻⁹³ In contrast, antibacterial prophylaxis can be considered in patients at high risk once the ANC is $<500/\mu\text{L}$ to mitigate the risk of severe infections. The CAR-HEMATOTOX score may be useful for guidance and identification of candidates at high risk.^{13,57} In a large retrospective analysis of patients with LBCL receiving CD19 CAR T cells, a significant reduction of severe bacterial infections with fluoroquinolone prophylaxis was observed in patients who were characterized as CAR-HEMATOTOX^{high} but not in those characterized as CAR-HEMATOTOX^{low}, supporting a risk-adapted approach. Importantly, the panel recommends adherence to institutional guidelines that take into account local epidemiology and resistance patterns. In this context, monitoring for multidrug-resistant gram-negative bacteria colonization (ie, active surveillance through rectal swab culture) may be useful both for baseline risk assessment and during prolonged neutropenia.
- Antifungal prophylaxis: to reduce the risk of invasive fungal disease, antimold prophylaxis (eg, micafungin or posaconazole) can be considered in patients at high risk for severe ICAHT (grade ≥ 3) once the ANC is $<500/\mu\text{L}$ (Table 3). Additional risk factors to consider are prior allo-HCT, prior invasive aspergillosis, and receipt of corticosteroids (either long-term, ≥ 72 hours; or high-dose, eg, >10 mg of dexamethasone or equivalent). The low overall incidence rate for invasive fungal disease in the context of CAR T-cell therapy should be taken into account,⁹⁴ although fungal infections represent a frequent cause of fatal infectious complications.^{22,95} Systemic primary antifungal prophylaxis should be continued until stable count recovery (ANC of $>500/\mu\text{L}$ over 3 days) and discontinuation of steroids for CRS/ICANS management.

Hematopoietic cell boost (HCB)

Patients who are unresponsive and/or refractory to G-CSF beyond day +14 after CAR T-cell infusion represent a clinically challenging subgroup of patients at high risk for severe and even fatal infectious complications. Although the evidence remains limited, TPO agonists can be offered in this setting, especially in cases of associated thrombocytopenia.⁷⁸ In cases of severe ICAHT in which an inflammatory stressor is deemed contributory (severe CRS/ICANS or CRS/MAS),

anti-inflammatory strategies such as pulse-dose corticosteroids and/or anticytokine therapies (eg, tocilizumab or anakinra) should be used. A promising strategy pertains to the use of cryopreserved autologous or allogeneic CD34⁺ hematopoietic cells from prior collection (either prior auto- or allo-HCT).⁹⁶⁻⁹⁸ Three recent case series shed light on both the safety and clinical feasibility of this approach across a broad population of pediatric and adult patients (summarized in supplemental Table 4). High rates of sustained neutrophil and platelet engraftment were noted across studies. Although HCB has been successfully applied during active infection,⁹⁹ clinicians should be aware of the possibility of immune reconstitution inflammatory syndrome in patients with prolonged BM aplasia.³¹ Because the earlier application of an available HCB was associated with superior survival outcomes,⁹⁸ the expert panel recommends considering the application of an HCB without prior conditioning chemotherapy for grade ≥ 3 ICAHT beyond day +14 if (1) a boost is readily available, and (2) G-CSF refractoriness has been established. At the same time, the survey results highlighted that even when HCB were considered a viable treatment option in a patient with prior auto-HCT, the cell products were often not available. Although prophylactic collection in high-risk candidates has been proposed as a potential mitigating strategy, the panel cautioned that the collection process may add to the already high logistic burden of CAR T-cell therapy (eg, coordination of apheresis slots and storage capacity), which could negatively impact vein-to-vein times in a state of high disease burden. Furthermore, the process could incur unnecessary collection- and storage-associated costs.^{100,101} Ultimately, it was concluded that further research is needed to assess the number needed to treat for prophylactic stem cell collection.

Allo-HCT

If the above options remain ineffective or elusive and grade 4 ICAHT persists beyond day +30, the expert panel recommends initiating a donor search for a potential allo-HCT as a last resort (ultima ratio). In such cases of life-threatening ICAHT, the benefit and risks of allo-HCT need to be carefully weighed and aligned with the patient's goals of care. Furthermore, the possibility of spontaneous count recovery needs to be considered seriously.^{34,102,103} Accordingly, the expert panel suggested that the ultimate trigger for allo-HCT needs to be discussed on a case-by-case basis. Month 3 to 6 after CAR T-cell infusion was deemed a reasonable time frame to balance both the risk of infection and possibility of spontaneous count recovery. Once the decision for allo-HCT has been made, details regarding donor selection, conditioning regimens, and immunosuppression have to be discussed. Experience and evidence are very limited and only general considerations can be reviewed here. As for every allo-HCT, the same basic principles should apply, keeping in mind that the primary indication is severe and persistent cytopenia although essentially all patients currently receive commercially available CAR T cells to treat malignant lymphoid disorders. Most importantly, salvage allo-HCT can also provide tumor control through the conditioning regimen and graft-versus-tumor effects and current standard procedures will most likely lead to eradication of CAR T cells at the latest when full donor chimerism has been established. Therefore, remission status must be determined before allo-HCT and may guide the choice of conditioning regimen and the taper of

immunosuppression. As usual, performance status, comorbidities, prior therapies, and expected antitumor activity should be carefully considered when discussing the transplantation modalities, donor choice, and selection.

Conclusions and outlook

Much progress has been made in the last years in defining hematological toxicity as a distinct toxicity entity of CAR T-cell therapy. Although the underlying pathophysiology remains incompletely understood, growing evidence points toward critical interactions between host hematopoiesis and CAR T-cell function and efficacy. By defining ICAHT and delineating a specific grading system, we herein provide a nomenclature that enables cross-trial comparisons and invites severity-based management strategies.

In this international consensus guidelines document, we have proposed a structured approach to diagnosis, grading/staging, and clinical management of ICAHT. This endeavor has also set the stage for areas of future development that will require collaboration between various European and non-European stakeholders involved in CAR T-cell therapy. Structured sample collection across multiple centers represents the basis for translational projects that delineate the underlying mechanisms of ICAHT by leveraging novel technologies such as multiomics and single-cell approaches. One area of particular interest lies in identifying early determinants of ICAHT by studying the peripheral blood immune contexture and/or the local BM microenvironment from pre-CAR T-cell samples. Furthermore, large retrospective real-world analyses may shed light on some of the differences in the clinical management of ICAHT that were identified by the EHA/EBMT survey. Residual questions relate to the optimal timing of G-CSF initiation as well as the optimal protocol to use (eg, prophylactic vs early G-CSF). The question of prophylactic collection of CD34⁺ hematopoietic cells in candidates at high risk and the optimal trigger time point for both HCB and allo-HCT represent unresolved issues that warrant further systematic study. Ultimately, prospective clinical trials will be needed that determine the potential benefits and evidence-base of treatment strategies that mitigate ICAHT.

Acknowledgment

The authors are grateful for the support by European Hematology Association and European Society for Blood and Marrow Transplantation, without which this work would not have been possible.

Authorship

Contribution: K.R. and M.S. conceptualized the study; K.R., M.S., M.A., E.B., A.B., P.B., B.B., R.B., M.G.C., C.C., F.C., P.C., J.D., R.D.B., R.G., R.H., G.I., U.J., M.J.K., S.M., A.N., F.O., Z.P., C.R., A.R., F.S.-G., I.S.-O., D.S., M.-L.S., J.A.S., C.T., M.T., P.L.Z., J.G.G., C.B., A.S., and I.Y.-A. performed the investigations; K.R., M.S., R.G., R.H., M.J.K., S.M., C.R., F.S.-G., D.S., M.-L.S., C.T., A.S., and I.Y.-A. performed formal analysis and visualization; K.R., M.S., I.S.-O., A.S., and I.Y.-A. developed the methodology; K.R. and M.S. wrote the original manuscript draft; K.R., M.S., M.A., E.B., A.B., P.B., B.B., R.B., M.G.C., C.C., F.C., P.C., J.D., R.D.B., R.G., R.H., G.I., U.J., M.J.K., S.M., A.N., F.O., Z.P., C.R., A.R., F.S.-G., I.S.-O., D.S., M.-L.S., J.A.S., C.T., M.T., P.L.Z., J.G.G., C.B., A.S., and I.Y.-A. wrote, reviewed, and edited the manuscript; and all authors read and approved the final manuscript.

Conflict-of-interest disclosure: K.R. reports research funding and travel support from Kite/Gilead; received honoraria from Novartis; consultants for and received honoraria from BMS/Celgene. M.S. receives industry research support from Amgen, BMS/Celgene, Kite/Gilead, Janssen, Miltenyi Biotech, Novartis, Roche, Seattle Genetics, and Takeda; serves as a consultant/adviser to AvenCell, CDR-Life, Ichnos Sciences, Incyte Biosciences, Janssen, Miltenyi Biotech, Molecular Partners, Novartis, Pfizer, and Takeda; and serves on the speakers' bureau at Amgen, AstraZeneca, BMS/Celgene, Gilead, GSK, Janssen, Novartis, Pfizer, Roche, and Takeda. E.B. received honorarium from, and serves on the advisory board of, Kite/Gilead, BMS/Celgene, Novartis, Pfizer, Incyte, ADC Therapeutics, Roche, and Takeda; receives personal fees from Kite/Gilead, BMS/Celgene, Novartis, and Pfizer; and reports research funding from Amgen and BMS. A.B. serves on the speakers' bureau of Novartis, Amgen, and Medac; and received meeting support from Novartis, Amgen, Medac, and Neovii. P.B. received honoraria and travel support from Allogene, Amgen, BMS, Kite/Gilead, Incyte, Jazz Pharmaceuticals, Miltenyi Biomedicine, Nektar, Novartis, and Pierre Fabre. R.B. received research funding from Servier and Allogene; and participated in advisory boards for BMS/Celgene, Janssen, Allogene, Takeda, Enara Bio, Securabio, Menarini Stemline, Arovella, and Kite/Gilead. M.C. reports consultancy for Novartis and Janssen. C.C. is a consultant adviser to Bellicum Pharmaceuticals, BMS, Kite/Gilead, Janssen Pharmaceuticals, and Novartis; serves on the speakers' bureau of BMS, Kite/Gilead, Janssen Pharmaceuticals, Jazz, Miltenyi Biotech, and Novartis; and received travel support from BMS and Kite/Gilead. P.C. received honoraria from AbbVie, Amgen, Celgene, Daiichi Sankyo, Janssen, Kite/Gilead, Novartis, Roche, Sanofi, Servier, and Takeda (consulting, advisory role, or lecturer); and has had travel and accommodations paid by for-profit health care companies during the past 2 years (Novartis, Janssen, BMS/Celgene, Takeda, Kite/Gilead, and Roche). R.D. serves on the scientific advisory board of, and as a conference speaker for, Novartis; serves on the scientific advisory board of, as a conference speaker for, and has received travel accommodation from, Kite/Gilead; serves on the scientific advisory board of Janssen; is a conference speaker for Pfizer; serves on the scientific advisory board of BMS/Celgene; and is a conference speaker for AbbVie and Incyte. R.G. discloses speaking honoraria from Biotech, Pfizer, Medac, and Magenta. R.H. received honoraria from Kite/Gilead, Novartis, Incyte, Janssen, MSD, Takeda, and Roche; and serves in a consultancy role at Kite/Gilead, Novartis, Bristol-Myers Squibb/Celgene, ADC Therapeutics, Incyte, and Miltenyi. G.I. received honoraria from BMS/Celgene, Novartis, Janssen, AstraZeneca, AbbVie, Miltenyi, and Roche. U.J. reports consultancy with Novartis, Janssen, and Roche; and having received honoraria from Gilead, BMS/Celgene, and Miltenyi. M.J.K. received honoraria from, and serves in a consulting/advisory role for, Bristol Myers Squibb, Celgene, Kite (a Gilead Company), Miltenyi Biotech, Novartis, and Roche; received research funding from Kite, Roche, Takeda, and Celgene; and received travel support from Kite, Miltenyi Biotech, Novartis, and Roche. S.M. serves, via his institution, on the speakers' bureau of Celgene/BMS, Novartis, and Janssen; has received travel support from, and serves on the expert panels of, Gilead/Kite; serves on the data safety monitoring boards of Miltenyi and Immunicon/Mendes; is the founder of, and serves in a leadership role in, SWECARNET; and is the spouse of the founder of ScientifyResearch. F.O. declares having had travel, accommodation, and expenses paid for by Takeda, Kyowa Kirin International, and Medac. C.R. received honoraria from BMS, Kite/Gilead, Novartis, and Amgen. F.S.G. received research support from Novartis and Gilead; and honoraria from Novartis, Gilead, Pfizer, and BMS-Celgene. D.S. serves on the advisory board of Autolus. M.L.S. serves on the advisory boards of, and declares consultancy with, Novartis, Kite-Gilead, Takeda, and Janssen; and received honoraria from Kite/Gilead. J.S. received speaker fees from Gilead; and reports consultancy with Kiadis, Medac, and Vertex. C.T. serves on the board of Novartis, Kite/Gilead, BMS, Roche, Takeda, Incyte, and Beigene. M.T. serves on the advisory boards of KITE/Gilead, BMS, Janssen, Novartis, Roche, Incyte, and AstraZeneca; and received research funding from KITE/Gilead, Takeda, Incyte, Roche, and Regeneron. P.L.Z. reports consultancy for MSD, Eusapharma, and Novartis; serves on the advisory boards of Securia Bio, Celltrion, Kite/Gilead, Janssen, BMS/Celgene, Servier, Sandoz, MSD, AstraZeneca, Takeda, Roche, Eusapharma, Kyowa Kirin, Novartis, ADC Therapeutics, Incyte, and Belgene; serves on the speakers' bureau of Celltrion, Kite/Gilead, Janssen, BMS/Celgene, Servier, MSD, AstraZeneca, Takeda, Roche, Eusapharma, Kyowa Kirin, Novartis, Incyte, and Belgene. J.G.G. received honoraria from Amgen, BMS Gilead/Kite, Janssen, and Novartis; and received research funding from AZ, BMS, and Janssen. C.B. is an inventor of

different patents on cancer immunotherapy and genetic engineering; is a member of advisory boards of, and consultant for, Intellia, Kite/Gilead, Miltenyi, Kiadis, QuellTx, Janssen, Chroma, Genyo, Pancancer-T, and Alia; and received research support from Intellia Therapeutics. A.S. received honoraria from Takeda, Bristol Myers Squibb, Merck Sharp & Dohme, Celgene, Janssen, Sanofi, Roche, Novartis, Gilead Sciences, and Janssen-Cilag; reports a consulting or advisory role for Takeda, Bristol Myers Squibb, Gilead Sciences, Celgene, Janssen, and Novartis; serves on the speakers' bureau of Takeda; received travel, accommodations, and expenses funds from Kite/Gilead. I.Y.-A. received honoraria from Novartis, Gilead/Kite, BMS, and Janssen. None of the mentioned conflicts of interest were related to financing or the content of this manuscript. The remaining authors declare no competing financial interests.

ORCID profiles: K.R., [0000-0003-3905-0251](#); M.S., [0000-0003-3905-0251](#); E.B., [0000-0003-2694-7510](#); A.B., [0000-0002-5879-0610](#); B.B., [0000-0002-7299-6770](#); M.G.C., [0000-0001-7015-4839](#); C.C., [0000-0002-3755-4889](#); P.C., [0000-0002-9186-1353](#); J.D., [0000-0002-5157-4376](#); R.D.B., [0000-0001-9001-573X](#); R.G., [0000-0002-6117-5328](#); R.H., [0000-0003-1729-8213](#); G.I., [0000-0003-0805-9288](#); U.J., [0000-0001-9826-1062](#); S.M., [0000-0002-8325-9215](#); C.R., [0000-0002-4901-5858](#); F.S.-G., [0000-0002-7076-7739](#); I.S.-O., [0009-0005-5519-9154](#); J.A.S.,

[0000-0001-6819-3476](#); P.L.Z., [0000-0002-2112-2651](#); J.G.G., [0000-0002-8505-7430](#); I.Y.-A., [0000-0003-4524-8782](#).

Correspondence: Marion Subklewe, Department of Medicine III–Hematology/Oncology, LMU–Klinikum der Universität München, Marchioninistr 15, 81377 Munich, Germany; email: marion.subklewe@med.uni-muenchen.de; and Ibrahim Yakoub-Agha, UAM allogreffes de CSH, CHRU, F-59037 Lille CEDEX, France; email: ibrahim.yakoubagha@chu-lille.fr.

Footnotes

Submitted 27 March 2023; accepted 23 May 2023; prepublished online on *Blood First Edition* 10 June 2023. <https://doi.org/10.1182/blood.2023020578>.

*K.R. and M.S. contributed equally to this study.

The online version of this article contains a data supplement.

There is a [Blood Commentary](#) on this article in this issue.

REFERENCES

- Locke FL, Miklos DB, Jacobson CA, et al. Axicabtagene ciloleucel as second-line therapy for large B-cell lymphoma. *N Engl J Med*. 2022;386(7):640-654.
- Abramson JS, Solomon SR, Arason JE, et al. Lisocabtagene maraleucel as second-line therapy for large B-cell lymphoma: primary analysis of phase 3 TRANSFORM study. *Blood*. 2023;141(14):1675-1684.
- Shah BD, Ghobadi A, Oluwole OO, et al. KTE-X19 for relapsed or refractory adult B-cell acute lymphoblastic leukaemia: phase 2 results of the single-arm, open-label, multicentre ZUMA-3 study. *Lancet*. 2021;398(10299):491-502.
- Wang M, Munoz J, Goy A, et al. KTE-X19 CAR T-cell therapy in relapsed or refractory mantle-cell lymphoma. *N Engl J Med*. 2020;382(14):1331-1342.
- Raje N, Berdeja J, Lin Y, et al. Anti-BCMA CAR T-cell therapy bb2121 in relapsed or refractory multiple myeloma. *N Engl J Med*. 2019;380(18):1726-1737.
- Snowden JA, Sanchez-Ortega I, Corbacioglu S, et al. Indications for haematopoietic cell transplantation for haematological diseases, solid tumours and immune disorders: current practice in Europe, 2022. *Bone Marrow Transplant*. 2022;57(8):1217-1239.
- Passweg JR, Baldomero H, Chabannon C, et al. Hematopoietic cell transplantation and cellular therapy survey of the EBMT: monitoring of activities and trends over 30 years. *Bone Marrow Transplant*. 2021;56(7):1651-1664.
- Shimabukuro-Vornhagen A, Godel P, Subklewe M, et al. Cytokine release syndrome. *J Immunother Cancer*. 2018;6(1):56.
- Karschnia P, Jordan JT, Forst DA, et al. Clinical presentation, management, and biomarkers of neurotoxicity after adoptive immunotherapy with CAR T cells. *Blood*. 2019;133(20):2212-2221.
- Sandler RD, Tattersall RS, Schoemans H, et al. Diagnosis and management of secondary HLH/MAS following HSCT and CAR-T cell therapy in adults; a review of the literature and a survey of practice within EBMT centres on behalf of the Autoimmune Diseases Working Party (ADWP) and Transplant Complications Working Party (TCWP). *Front Immunol*. 2020;11:524.
- Hines MR, Knight TE, McNerney KO, et al. Immune effector cell associated hemophagocytic lymphohistiocytosis-like syndrome (IEC-HS). *Transplant Cell Ther*. 2023;29(7):438.e1-438.e16.
- Fried S, Avigdor A, Bielora B, et al. Early and late hematologic toxicity following CD19 CAR-T cells. *Bone Marrow Transplant*. 2019;54(10):1643-1650.
- Rejeski K, Perez A, Sesques P, et al. CAR-HEMATOTOX: a model for CAR T-cell-related hematologic toxicity in relapsed/refractory large B-cell lymphoma. *Blood*. 2021;138(24):2499-2513.
- Wudhikarn K, Pennisi M, Garcia-Recio M, et al. DLBCL patients treated with CD19 CAR T cells experience a high burden of organ toxicities but low nonrelapse mortality. *Blood Adv*. 2020;4(13):3024-3033.
- Dickinson MJ, Carlo-Stella C, Morschhauser F, et al. Glofitamab for relapsed or refractory diffuse large B-cell lymphoma. *N Engl J Med*. 2022;387(24):2220-2231.
- Thieblemont C, Phillips T, Ghesquieres H, et al. Epcoritamab, a novel, subcutaneous CD3xCD20 bispecific T-cell-engaging antibody, in relapsed or refractory large B-cell lymphoma: dose expansion in a phase I/II trial. *J Clin Oncol*. 2023;41(12):2238-2247.
- Charl A, Minnema MC, Berdeja JG, et al. Talquetamab, a T-cell-redirecting GPRC5D bispecific antibody for multiple myeloma. *N Engl J Med*. 2022;387(24):2232-2244.
- Moreau P, Girgis S, Goldberg JD. Teclistamab in relapsed or refractory multiple myeloma. Reply. *N Engl J Med*. 2022;387(18):1722-1723.
- Goebeler ME, Knop S, Viardot A, et al. Bispecific T-cell engager (BiTE) antibody construct blinatumomab for the treatment of patients with relapsed/refractory non-Hodgkin lymphoma: final results from a phase I study. *J Clin Oncol*. 2016;34(10):1104-1111.
- Hill JA, Seo SK. How I prevent infections in patients receiving CD19-targeted chimeric antigen receptor T cells for B-cell malignancies. *Blood*. 2020;136(8):925-935.
- Nastoupil LJ, Jain MD, Feng L, et al. Standard-of-care axicabtagene ciloleucel for relapsed or refractory large B-cell lymphoma: results from the US Lymphoma CAR T Consortium. *J Clin Oncol*. 2020;38(27):3119-3128.
- Rejeski K, Perez A, Iacoboni G, et al. The CAR-HEMATOTOX risk-stratifies patients for severe infections and disease progression after CD19 CAR-T in R/R LBCL. *J Immunother Cancer*. 2022;10(5):e004475.
- Bethge WA, Martus P, Schmitt M, et al. GLA/DRST real-world outcome analysis of CAR-T cell therapies for large B-cell lymphoma in Germany. *Blood*. 2022;140(4):349-358.
- Berdeja JG, Madduri D, Usmani SZ, et al. Ciltacabtagene autoleucel, a B-cell maturation antigen-directed chimeric antigen receptor T-cell therapy in patients with relapsed or refractory multiple myeloma (CARTITUDE-1): a phase 1b/2 open-label study. *Lancet*. 2021;398(10297):314-324.
- Munshi NC, Anderson LD Jr, Shah N, et al. Idecabtagene vicleucel in relapsed and refractory multiple myeloma. *N Engl J Med*. 2021;384(8):705-716.
- Jacobson CA, Chavez JC, Sehgal AR, et al. Axicabtagene ciloleucel in relapsed or refractory indolent non-Hodgkin lymphoma (ZUMA-5): a single-arm,

- multicentre, phase 2 trial. *Lancet Oncol*. 2022;23(1):91-103.
27. Iacoboni G, Rejeski K, Villacampa G, et al. Real-world evidence of brexucabtagene autoleucel for the treatment of relapsed or refractory mantle cell lymphoma. *Blood Adv*. 2022;6(12):3606-3610.
 28. Maude SL, Laetsch TW, Buechner J, et al. Tisagenlecleucel in children and young adults with B-cell lymphoblastic leukemia. *N Engl J Med*. 2018;378(5):439-448.
 29. Lichtenstein DA, Schischlik F, Shao L, et al. Characterization of HLH-like manifestations as a CRS variant in patients receiving CD22 CAR T cells. *Blood*. 2021;138(24):2469-2484.
 30. Cordeiro A, Bezerra ED, Hirayama AV, et al. Late events after treatment with CD19-targeted chimeric antigen receptor modified T cells. *Biol Blood Marrow Transplant*. 2020;26(1):26-33.
 31. Rejeski K, Kunz WG, Rudelius M, et al. Severe *Candida glabrata* pancolitis and fatal *Aspergillus fumigatus* pulmonary infection in the setting of bone marrow aplasia after CD19-directed CAR T-cell therapy - a case report. *BMC Infect Dis*. 2021;21(1):121.
 32. Jain T, Olson TS, Locke FL. How I treat cytopenias after CAR T-cell therapy. *Blood*. 2023;141(20):2460-2469.
 33. Juluri KR, Wu V, Voutsinas JM, et al. Severe cytokine release syndrome is associated with hematologic toxicity following CD19 CAR T-cell therapy. *Blood Adv*. 2022;6(7):2055-2068.
 34. Jain T, Knezevic A, Pennisi M, et al. Hematopoietic recovery in patients receiving chimeric antigen receptor T-cell therapy for hematologic malignancies. *Blood Adv*. 2020;4(15):3776-3787.
 35. Rejeski K, Wu Z, Blumenberg V, et al. Oligoclonal T-cell expansion in a patient with bone marrow failure after CD19 CAR-T for Richter transformed DLBCL. *Blood*. 2022;140(20):2175-2179.
 36. Rejeski K, Greco R, Onida F, et al. An international survey on grading, diagnosis, and management of immune effector cell-associated hematotoxicity (ICAHT) following CAR T-cell therapy on behalf of the EBMT and EHA. *Hemasphere*. 2023;7(5):e889.
 37. Taplit RA, Kennedy EB, Bow EJ, et al. Antimicrobial prophylaxis for adult patients with cancer-related immunosuppression: ASCO and IDSA Clinical Practice Guideline update. *J Clin Oncol*. 2018;36(30):3043-3054.
 38. Yakoub-Agha I, Greco R, Onida F, et al. Practice harmonization workshops of EBMT: an expert-based approach to generate practical and contemporary guidelines within the arena of hematopoietic cell transplantation and cellular therapy. *Bone Marrow Transplant*. 2023;58(6):696-700.
 39. Xia Y, Zhang J, Li J, et al. Cytopenias following anti-CD19 chimeric antigen receptor (CAR) T cell therapy: a systematic analysis for contributing factors. *Ann Med*. 2022;54(1):2951-2965.
 40. Brudno JN, Natrakul D, Lam N, Dulau-Florea A, Yuan CM, Kochenderfer JN. Acute and delayed cytopenias following CAR T-cell therapy: an investigation of risk factors and mechanisms. *Leuk Lymphoma*. 2022;63(8):1849-1860.
 41. Roddie C, Neill L, Osborne W, et al. Effective bridging therapy can improve CD19 CAR-T outcomes while maintaining safety in patients with large B-cell lymphoma. *Blood Adv*. 2023;7(12):2872-2883.
 42. Rejeski K, Perez Perez A, Iacoboni G, et al. Biphasic neutrophil recovery after CD19 CART in R/R LBCL is associated with superior PFS/OS, robust CAR T-cell expansion in relation to baseline tumor volume, and a decrease of systemic inflammation over time [abstract]. *Blood*. 2022;140(suppl 1):4549-4551.
 43. Rejeski K, Wang Y, Albany O, et al. The CAR-hematotox score identifies patients at high risk for hematological toxicity, infections and poor clinical outcomes following brexucabtagene autoleucel in relapsed/refractory mantle cell lymphoma [abstract]. *Blood*. 2022;140(suppl 1):651-653.
 44. Rejeski K, Hansen DK, Bansal R, et al. The CAR-hematotox score as a prognostic model of toxicity and response in patients receiving BCMA-directed CAR-T for relapsed/refractory multiple myeloma [abstract]. *Blood*. 2022;140(suppl 1):7506-7508.
 45. Liu Y, Derkach A, Lewis N, et al. Clonal hematopoiesis in diffuse large B-cell lymphoma: clinical impact and genetic relatedness to lymphoma and therapy-related myeloid neoplasm. *Haematologica*. 2023;108(3):917-922.
 46. Saini NY, Swoboda DM, Greenbaum U, et al. Clonal hematopoiesis is associated with increased risk of severe neurotoxicity in axicabtagene ciloleucel therapy of large B-cell lymphoma. *Blood Cancer Discov*. 2022;3(5):385-393.
 47. Miller PG, Sperling AS, Brea EJ, et al. Clonal hematopoiesis in patients receiving chimeric antigen receptor T-cell therapy. *Blood Adv*. 2021;5(15):2982-2986.
 48. Jain MD, Zhao H, Wang X, et al. Tumor interferon signaling and suppressive myeloid cells are associated with CAR T-cell failure in large B-cell lymphoma. *Blood*. 2021;137(19):2621-2633.
 49. Bachy E, Le Gouill S, Di Blasi R, et al. A real-world comparison of tisagenlecleucel and axicabtagene ciloleucel CAR T cells in relapsed or refractory diffuse large B cell lymphoma. *Nat Med*. 2022;28(10):2145-2154.
 50. Luo W, Li C, Zhang Y, et al. Adverse effects in hematologic malignancies treated with chimeric antigen receptor (CAR) T cell therapy: a systematic review and meta-analysis. *BMC Cancer*. 2022;22(1):98.
 51. Kawalekar OU, O' Connor RS, Fraietta JA, et al. Distinct signaling of coreceptors regulates specific metabolism pathways and impacts memory development in CAR T cells. *Immunity*. 2016;44(3):712.
 52. Zhao Z, Condomines M, van der Stegen SJC, et al. Structural design of engineered costimulation determines tumor rejection kinetics and persistence of CAR T cells. *Cancer Cell*. 2015;28(4):415-428.
 53. Wang Y, Song Z, Geng Y, et al. The risk factors and early predictive model of hematotoxicity after CD19 chimeric antigen receptor T cell therapy. *Front Oncol*. 2022;12:987965.
 54. Li X, Deng Q, Henderson J, et al. Targetable cellular etiology of prolonged cytopenia following CD19 CAR T-cell therapy [abstract]. *Blood*. 2022;140(suppl 1):4502-4503.
 55. de Bruin AM, Demirel O, Hooibrink B, Brandts CH, Nolte MA. Interferon-gamma impairs proliferation of hematopoietic stem cells in mice. *Blood*. 2013;121(18):3578-3585.
 56. Morales-Mantilla DE, King KY. The role of interferon-gamma in hematopoietic stem cell development, homeostasis, and disease. *Curr Stem Cell Rep*. 2018;4(3):264-271.
 57. Rejeski K, Blumenberg V, Iacoboni G, et al. Identifying early infections in the setting of CRS with routine and exploratory serum proteomics and the HT10 score following CD19 CAR-T for relapsed/refractory B-NHL. *Hemasphere*. 2023;7(4):e858.
 58. Neelapu SS, Tummala S, Kebriaei P, et al. Chimeric antigen receptor T-cell therapy - assessment and management of toxicities. *Nat Rev Clin Oncol*. 2018;15(1):47-62.
 59. Hayden PJ, Roddie C, Bader P, et al. Management of adults and children receiving CAR T-cell therapy: 2021 best practice recommendations of the European Society for Blood and Marrow Transplantation (EBMT) and the Joint Accreditation Committee of ISCT and EBMT (JACIE) and the European Haematology Association (EHA). *Ann Oncol*. 2022;33(3):259-275.
 60. Cutini I, Puccini B, Fabbri A, et al. Late haemophagocytic lymphohistiocytosis in a patient treated with axicabtagene ciloleucel. *Transpl Immunol*. 2022;75:101719.
 61. Henter JL, Horne A, Arico M, et al. HLH-2004: diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer*. 2007;48(2):124-131.
 62. Fardet L, Galicier L, Lambotte O, et al. Development and validation of the HScore, a score for the diagnosis of reactive hemophagocytic syndrome. *Arthritis Rheumatol*. 2014;66(9):2613-2620.

63. Zoref-Lorenz A, Murakami J, Hofstetter L, et al. An improved index for diagnosis and mortality prediction in malignancy-associated hemophagocytic lymphohistiocytosis. *Blood*. 2022;139(7):1098-1110.
64. McNerney KO, DiNofia AM, Teachey DT, Grupp SA, Maude SL. Potential role of IFN γ inhibition in refractory cytokine release syndrome associated with CAR T-cell therapy. *Blood Cancer Discov*. 2022;3(2):90-94.
65. Rainone M, Ngo D, Baird JH, et al. Interferon-gamma blockade in CAR T-cell therapy-associated macrophage activation syndrome/hemophagocytic lymphohistiocytosis. *Blood Adv*. 2023;7(4):533-536.
66. La Rosee P, Horne A, Hines M, et al. Recommendations for the management of hemophagocytic lymphohistiocytosis in adults. *Blood*. 2019;133(23):2465-2477.
67. Kopolovic I, Ostro J, Tsubota H, et al. A systematic review of transfusion-associated graft-versus-host disease. *Blood*. 2015;126(3):406-414.
68. Foukaneli T, Kerr P, Bolton-Maggs PHB, et al. Guidelines on the use of irradiated blood components. *Br J Haematol*. 2020;191(5):704-724.
69. Giavridis T, van der Stegen SJC, Eyquem J, Hamieh M, Piersigilli A, Sadelain M. CAR T cell-induced cytokine release syndrome is mediated by macrophages and abated by IL-1 blockade. *Nat Med*. 2018;24(6):731-738.
70. Sterner RM, Sakemura R, Cox MJ, et al. GM-CSF inhibition reduces cytokine release syndrome and neuroinflammation but enhances CAR-T cell function in xenografts. *Blood*. 2019;133(7):697-709.
71. Barreto JN, Bansal R, Hathcock MA, et al. The impact of granulocyte colony stimulating factor on patients receiving chimeric antigen receptor T-cell therapy. *Am J Hematol*. 2021;96(10):E399-E402.
72. Galli E, Allain V, Di Blasi R, et al. G-CSF does not worsen toxicities and efficacy of CAR-T cells in refractory/relapsed B-cell lymphoma. *Bone Marrow Transplant*. 2020;55(12):2347-2349.
73. Lievin R, Di Blasi R, Morin F, et al. Effect of early granulocyte-colony-stimulating factor administration in the prevention of febrile neutropenia and impact on toxicity and efficacy of anti-CD19 CAR-T in patients with relapsed/refractory B-cell lymphoma. *Bone Marrow Transplant*. 2022;57(3):431-439.
74. Miller KC, Johnson PC, Abramson JS, et al. Effect of granulocyte colony-stimulating factor on toxicities after CAR T cell therapy for lymphoma and myeloma. *Blood Cancer J*. 2022;12(10):146.
75. Ma S, Li H, Zhou D, et al. Associations of granulocyte colony-stimulating factor with toxicities and efficacy of chimeric antigen receptor T-cell therapy in relapsed or refractory multiple myeloma. *Cytotherapy*. 2023;25(6):653-658.
76. Baur R, Jitschin R, Kharboulit S, et al. Thrombopoietin receptor agonists for acquired thrombocytopenia following anti-CD19 CAR-T-cell therapy: a case report. *J Immunother Cancer*. 2021;9(7):e002721.
77. Beyar-Katz O, Perry C, On YB, et al. Thrombopoietin receptor agonist for treating bone marrow aplasia following anti-CD19 CAR-T cells-single-center experience. *Ann Hematol*. 2022;101(8):1769-1776.
78. Drillet G, Lhomme F, De Guibert S, Manson G, Houot R. Prolonged thrombocytopenia after CAR T-cell therapy: the role of thrombopoietin receptor agonists. *Blood Adv*. 2023;7(4):537-540.
79. Drexler B, Passweg J. Current evidence and the emerging role of eltrombopag in severe aplastic anemia. *Ther Adv Hematol*. 2021;12:2040620721998126.
80. Peffault de Latour R, Kulasekararaj A, Iacobelli S, et al. Eltrombopag added to immunosuppression in severe aplastic anemia. *N Engl J Med*. 2022;386(1):11-23.
81. Bento L, Bastida JM, Garcia-Cadenas I, et al. Thrombopoietin receptor agonists for severe thrombocytopenia after allogeneic stem cell transplantation: experience of the Spanish Group of Hematopoietic Stem Cell Transplant. *Biol Blood Marrow Transplant*. 2019;25(9):1825-1831.
82. Ball S, Brown TJ, Das A, Khera R, Khanna S, Gupta A. Effect of neutropenic diet on infection rates in cancer patients with neutropenia: a meta-analysis of randomized controlled trials. *Am J Clin Oncol*. 2019;42(3):270-274.
83. Sonbol MB, Jain T, Firwana B, et al. Neutropenic diets to prevent cancer infections: updated systematic review and meta-analysis. *BMJ Support Palliat Care*. 2019;9(4):425-433.
84. Stella F, Marasco V, Levati GV, et al. Non-restrictive diet does not increase infections in patients with neutropenia after stem cell transplantation: final analysis of the Neutrodiet Multicenter, Randomized trial [abstract]. *Blood*. 2022;140(suppl 1):417-419.
85. Teipel R, Kroschinsky F, Kramer M, et al. Prevalence and variation of CHIP in patients with aggressive lymphomas undergoing CD19-directed CAR T-cell treatment. *Blood Adv*. 2022;6(6):1941-1946.
86. Spellberg B, Doi Y. The rise of fluoroquinolone-resistant *Escherichia coli* in the community: scarier than we thought. *J Infect Dis*. 2015;212(12):1853-1855.
87. Lautenbach E, Metlay JP, Bilker WB, Edelstein PH, Fishman NO. Association between fluoroquinolone resistance and mortality in *Escherichia coli* and *Klebsiella pneumoniae* infections: the role of inadequate empirical antimicrobial therapy. *Clin Infect Dis*. 2005;41(7):923-929.
88. Trecarichi EM, Tumbarello M, Spanu T, et al. Incidence and clinical impact of extended-spectrum-beta-lactamase (ESBL) production and fluoroquinolone resistance in bloodstream infections caused by *Escherichia coli* in patients with hematological malignancies. *J Infect*. 2009;58(4):299-307.
89. Bow EJ. Fluoroquinolones, antimicrobial resistance and neutropenic cancer patients. *Curr Opin Infect Dis*. 2011;24(6):545-553.
90. Schubert ML, Rohrbach R, Schmitt M, Stein-Thoeringer CK. The potential role of the intestinal micromilieu and individual microbes in the immunobiology of chimeric antigen receptor T-cell therapy. *Front Immunol*. 2021;12:670286.
91. Blumenberg V, Busch G, Baumann S, et al. High bacterial abundances of *Dorea* and *Pediococcus* in the gut microbiome linked to expansion, immune checkpoint expression and efficacy of CD19-directed CAR T-cells in patients with r/r DLBCL [abstract]. *Blood*. 2021;138(suppl 1):2792.
92. Smith M, Dai A, Ghilardi G, et al. Gut microbiome correlates of response and toxicity following anti-CD19 CAR T cell therapy. *Nat Med*. 2022;28(4):713-723.
93. Stein-Thoeringer CK, Saini NY, Zamir E, et al. A non-antibiotic-disrupted gut microbiome is associated with clinical responses to CD19-CAR-T cell cancer immunotherapy. *Nat Med*. 2023;29(4):906-916.
94. Little JS, Aleissa MM, Beluch K, et al. Low incidence of invasive fungal disease following CD19 chimeric antigen receptor T-cell therapy for non-Hodgkin lymphoma. *Blood Adv*. 2022;6(16):4821-4830.
95. Haidar G, Dorritie K, Farah R, Bogdanovich T, Nguyen MH, Samanta P. Invasive mold infections after chimeric antigen receptor-modified t-cell therapy: a case series, review of the literature, and implications for prophylaxis. *Clin Infect Dis*. 2020;71(3):672-676.
96. Rejeski K, Burchert A, Iacoboni G, et al. Safety and feasibility of stem cell boost as a salvage therapy for severe hematotoxicity after CD19 CAR T-cell therapy. *Blood Adv*. 2022;6(16):4719-4725.
97. Mullanfiroze K, Lazareva A, Chu J, et al. CD34+-selected stem cell boost can safely improve cytopenias following CAR T-cell therapy. *Blood Adv*. 2022;6(16):4715-4718.
98. Gagelmann N, Wulf GG, Duell J, et al. Hematopoietic stem cell boost for persistent neutropenia after CAR T-cell therapy: a GLA/DRST study. *Blood Adv*. 2023;7(4):555-559.
99. Lipsitt A, Beattie L, Harstead E, et al. Allogeneic CD34(+) selected hematopoietic stem cell boost following CAR T-cell therapy in a patient with prolonged cytopenia and active infection. *Pediatr Blood Cancer*. 2023;70(3):e30166.

100. Chhabra S, Thapa B, Szabo A, et al. Utilization and cost implications of hematopoietic progenitor cells stored for a future salvage autologous transplantation or stem cell boost in myeloma patients. *Biol Blood Marrow Transplant*. 2020;26(11):2011-2017.
101. Liang EC, Muffy LS, Shiraz P, et al. Use of backup stem cells for stem cell boost and second transplant in patients with multiple myeloma undergoing autologous stem cell transplantation. *Transplant Cell Ther*. 2021; 27(5):405.e1-405.e6.
102. Logue JM, Peres LC, Hashmi H, et al. Early cytopenias and infections after standard of care idecabtagene vicleucel in relapsed or refractory multiple myeloma. *Blood Adv*. 2022;6(24):6109-6119.
103. Logue JM, Zucchetti E, Bachmeier CA, et al. Immune reconstitution and associated infections following axicabtagene ciloleucel in relapsed or refractory large B-cell lymphoma. *Haematologica*. 2020;106(4):978-986.
104. Alizadeh D, Wong RA, Yang X, et al. IL15 enhances CAR-T cell antitumor activity by reducing mTORC1 activity and preserving their stem cell memory phenotype. *Cancer Immunol Res*. 2019;7(5):759-772.
105. Pascutti MF, Erkelens MN, Nolte MA. Impact of viral infections on hematopoiesis: from beneficial to detrimental effects on bone marrow output. *Front Immunol*. 2016; 7:364.
106. Porter TJ, Lazarevic A, Ziggas JE, et al. Hyperinflammatory syndrome resembling haemophagocytic lymphohistiocytosis following axicabtagene ciloleucel and brexucabtagene autoleucel. *Br J Haematol*. 2022;199(5):720-727.

© 2023 by The American Society of Hematology