

The growing importance of genetics in human reproduction and development

Ježek, Davor

Source / Izvornik: **Croatian Medical Journal, 2023, 64, 305 - 306**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.3325/cmj.2023.64.305>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:222463>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International/Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-01-27**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



The growing importance of genetics in human reproduction and development

Davor Ježek^{1,2,3}

¹Department of Histology and Embryology, University of Zagreb, School of Medicine, Zagreb, Croatia

²Department for Transfusion Medicine and Transplantation Biology, Zagreb University Hospital, Zagreb, Croatia

³Scientific Centre of Excellence for Reproductive and Regenerative Medicine, University of Zagreb, School of Medicine, Zagreb, Croatia

davor.jezek@mef.hr

Human reproduction and growth depend on normal developmental processes such as the production of competent gametes, fertilization, embryonic development, pregnancy, and birth. The characteristics of a newborn are determined by specific genes on maternally and paternally inherited chromosomes. Forty-six chromosomes in humans bear approximately 23 000 genes. If genes are located on the same chromosome, they are usually inherited together and are known as linked genes. Human fertility or infertility is determined by several biological processes, including organ formation and development, neuroendocrine regulation, hormone production, meiosis, and mitosis (1-3).

If one takes the example of neuroendocrine regulation, under normal physiological conditions, a precise arrangement of hypothalamic kisspeptin neurons and gonadotropin-releasing hormone neurons induces the secretion of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) from the anterior pituitary. Adequate secretion of gonadotropins initiates gonadal steroid production, which regulates testicular or ovarian development. Hormones produced by the testes or ovaries then exert feedback effects on the hypothalamus and pituitary to adjust the concentrations of FSH and LH. Defects at one or more levels of the hypothalamic-pituitary-gonadal axis can cause hypogonadism, which manifests as a reduction in sex hormones, absent or delayed puberty, and abnormal function of the gonads. Hypogonadotropic hypogonadism refers to an insufficient release of gonadotropin-releasing hormone or gonadotropin caused by dysfunction of the

pituitary gland or the hypothalamus. More than 40 genes have been associated with the pathogenesis of congenital hypogonadotropic hypogonadism (4,5).

Chromosomal abnormalities, which can be numerical or structural, are frequent causes of birth defects and spontaneous abortions. Even in healthy young couples, 50% of conceptions end in a spontaneous abortion, and 50% of these have significant chromosomal abnormalities. The most common chromosomal abnormalities are Turner syndrome, triploidy, and trisomy 16. Chromosomal abnormalities account for 10% of major birth defects, and gene mutations account for an additional 8%. Microdeletions spanning only a few adjoining genes may result in a microdeletion or contiguous gene syndrome. Identifying genetic pathways controlling gamete and somatic cell differentiation is vital for future diagnostics, especially concerning azoospermia. Owing to emerging genetic panels, patients could now be spared from unnecessary conventional or microtesticular sperm extraction procedures in the near future. In addition, novel marker genes identified in the frame of single-cell sequencing (scRNA-seq) studies may be included in genetic screening in the expanding field of reproductive genetics. This could lead to new infertility treatments, including stem cell differentiation, *in vitro* spermatogenesis/ovary, and gene therapy (6,7).

Several articles in this issue of the *Croatian Medical Journal* indicate the importance of genetic diagnosis and screening. The article by Sansović et al (8) highlights

the importance of clinical evaluation in subjects with familial hearing loss and the research on family segregation in assessing the pathogenicity of variants in the α -tectorin gene. Odak et al (9) focused on the role of the *KMT5B* gene in the pathogenesis of neurodevelopmental disorders. The *KMT5B* gene, discovered a few years ago, is related to abnormal activities of enzymes that regulate histone activity and gene expression during brain development. A comprehensive clinical and molecular approach to patients with *KMT5B* variants is necessary to optimize their treatment. Another case report, authored by Morožin-Pohovski et al (10), describes a boy with microcephaly, trigonocephaly, and dysmorphic features. This is very likely the first report in the literature describing a patient who inherited 16p11.2 duplications from both parents. The article by Fučić et al (11) deals with the early environmental exposure of the mother and the child to pesticides. The authors found that interleukin-2 levels, hypomethylation of interleukin-2 gene site 1, and the mother's rural residence, probably due to pesticide exposure, were predictive biomarkers for preterm birth. Finally, the article by Hat et al (12) focused on aging and sexual dimorphism of the human lacrimal gland.

The diagnostic and therapeutic procedures in human reproduction and development will soon include more advanced genetic testing, multi-omics analysis, microbiota treatment, germline nuclear transfer, *in vitro* gametogenesis, gene therapy, and increased application of artificial intelligence.

References

- 1 Verpoest W, Okutman Ö, Van Der Kelen A, Sermon K, Viville S. Genetics of infertility: a paradigm shift for medically assisted reproduction. *Hum Reprod*. 2023. [Medline:37801292](#) [doi:10.1093/humrep/dead199](#)
- 2 The Guideline Group on Unexplained Infertility, Romualdi D, Ata B, Bhattacharya S, Bosch E, Costello M, et al. Evidence-based guideline: unexplained infertility. *Hum Reprod*. 2023;38:1881-90. [Medline:37599566](#) [doi:10.1093/humrep/dead150](#)
- 3 Wei Y, Wang J, Qu R, Zhang W, Tan Y, Sha Y, et al. Genetic mechanisms of fertilization failure and early embryonic arrest: a comprehensive review. *Human Reproduction Update*. 2023; [dmad026](#).
- 4 Vallet-Buisan M, Mecca R, Jones C, Coward K, Yeste M. Contribution of semen to early embryo development: fertilization and beyond. *Hum Reprod Update*. 2023;29:395-433. [Medline:36882116](#) [doi:10.1093/humupd/dmad006](#)
- 5 Liang Z, Wen Q, Li J, Zeng D, Huang P. A systematic review and meta-analysis: clinical outcomes of recurrent pregnancy failure resulting from preimplantation genetic testing for aneuploidy. *Front Endocrinol (Lausanne)*. 2023;14:1178294. [Medline:37850092](#) [doi:10.3389/fendo.2023.1178294](#)
- 6 Schaub AM, Gonzalez TL, Dorfman AE, Novoa AG, Hussaini RA, Harakuni PM, et al. A systematic review of genome wide analyses of methylation changes associated with assisted reproductive technologies in various tissues. *Fertil Steril*. 2023 Oct 10:S0015-0282(23)01922-2.
- 7 McCoy RC, Summers MC, McCollin A, Ottolini CS, Ahuja K, Handyside AH. Meiotic and mitotic aneuploidies drive arrest of in vitro fertilized human preimplantation embryos. *Genome Med*. 2023;15:77. [Medline:37779206](#) [doi:10.1186/s13073-023-01231-1](#)
- 8 Sansović I, Meašić AM, Odak LJ, Kero M. Clinical relevance of the TECTA c.6183G>T variant identified in a family with autosomal dominant hearing loss: a case report. *Croat Med J*. 2023;64:329-33.
- 9 Odak LJ, Vulin K, Meašić AM, Šamadan L, Tripalo Batoš A. Neurodevelopmental disorder caused by an inherited novel *KMT5B* variant: case report. *Croat Med J*. 2023;64:334-8.
- 10 Morožin-Pohovski L, Sansović I, Vulin K, Odak LJ. The first case report of distal 16p12.1p11.2 trisomy and proximal 16p11.2 tetrasomy inherited from both parents. *Croat Med J*. 2023;64:339-43.
- 11 Fučić A, Knežević J, Krsić J, Polančec D, Sinčić N, Sindičić Dessardo N, et al. Interleukin-2 gene methylation levels and interleukin-2 levels associated with environmental exposure as risk biomarkers for preterm birth. *Croat Med J*. 2023;64:320-8.
- 12 Hat K, Kaštelan S, Planinić A, Muller D, Ježek D. Pathohistological features of the aging human lacrimal gland. *Croat Med J*. 2023;64:307-19.