

Application of microsatellite loci on the chromosome X for rapid prenatal detection of the chromosome X numerical abnormalities

Crkvenac Gornik, Kristina; Grubić, Zorana; Štingl, Katarina; Tonković Đurišević, Ivana; Begović, Davor

Source / Izvornik: **Croatian Medical Journal, 2011, 52, 392 - 395**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.3325/cmj.2011.52.392>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:863810>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-01-22**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



Application of microsatellite loci on the chromosome X for rapid prenatal detection of the chromosome X numerical abnormalities

Kristina Crkvenac Gornik¹,
Zorana Grubić², Katarina
Štingl², Ivana Tonković
Đurišević¹, Davor Begović¹

¹Division of Genetics, Department
of Pediatrics University Hospital
Center, Zagreb, Croatia

²Tissue Typing Center, University
Hospital Center, Zagreb, Croatia

Aim To determine the value of short-tandem repeat markers on the chromosome X (X-STR) for prenatal diagnostics of the chromosome X numerical disorders.

Methods We investigated the genetic variability of 5 X-markers (DXS9895, DXS6810, DXS6803, GATA172D05, and HPRTB) in 183 healthy Croatian individuals (90 men and 93 women). We also tested 13 patients with X chromosome disorders (Turner syndrome – 6 cases; Klinefelter syndrome – 5 cases, and Triple X syndrome – 2 cases). The analysis was performed using polymerase chain reaction amplification with specific primers and electrophoresis on a polyacrylamide gel. The study was performed in 2010.

Results Our sample showed no significant differences in allelic frequencies of the investigated X-markers from other European populations. A set of 5 X-STR markers was sufficiently informative for a successful determination of the chromosome X numerical abnormalities.

Conclusion Since no false positive or negative results were observed, diagnostic value of the investigated X-STR loci for prenatal detection of chromosome X numerical disorders was confirmed. Our study represents an important step toward an improved prenatal diagnostics in Croatia.

Received: March 31, 2011

Accepted: May 9, 2011

Correspondence to:

Kristina Crkvenac Gornik
Division of Genetics, Department of
Pediatrics, University Hospital Center
Kišpatičeva 12
10000 Zagreb, Croatia
kcrkven@kbc-zagreb.hr

The analysis of short tandem repeat (STR) markers using polymerase chain reaction (PCR) method has become a widely applied technique in forensic individual identification, rapid detection of chromosome aneuploidies in prenatal and postnatal diagnosis, as well as paternity testing (1-5). Until now, a large number of autosomal and Y-chromosomal markers has been forensically evaluated and used for various purposes. Although X-chromosomal markers have been increasingly applied in both forensic and medical field, their role has not been as extensively investigated as that of autosomal and Y-chromosomal markers. Several investigations have documented the accuracy of fluorescent PCR using STR loci for the rapid prenatal diagnoses of numerical disorders affecting the chromosomes 21, 18, and 13 (6-8). However, the low polymorphism of the most chromosome X and Y markers has hampered the use of the PCR-STR approach for the detection of numerical disorders of sex chromosomes, such as the Turner (45, X) or Klinefelter (47, XXY) syndromes (1,7). A few years ago, a group of authors reported on the application of PCR-STR method in the detection of X-chromosome abnormalities (9,10). One of the biggest challenges is the Turner syndrome, in which a sufficient number of STR loci has to be included to be sure that the individual has only a single X chromosome. The aims of the present study were to investigate the diagnostic informativeness of 5 X-linked STR markers: DXS9895 (Xpter-Xp22.1), GATA172D05 (Xq26.1), DXS6810 (Xq12-Xq21.33), DXS6803 (Xq24-Xq27), and HPRTB (Xq27.3) in the Croatian population and to evaluate the diagnostic value of these 5 loci.

PARTICIPANTS AND METHODS

We used blood samples from 183 unrelated healthy individuals (90 men and 93 women) from the Croatian population (citizens of Zagreb), available in the Tissue Typing Center of the University Hospital Center. All participants gave a written informed consent and filled out a questionnaire on their demographic characteristics and medical background. Their anonymity was ensured by giving each participant a code number. Hundred and fifty samples had been previously tested by conventional cytogenetic analysis. In this group, there were also 13 samples with X chromosome disorders (2 with 47, XXX; 6 with 45, X; and 5 with 47, XXY). The analysis of all participants was performed by PCR-STR method. Genomic DNA was isolated from peripheral blood by NucleoSpin Blood isolation kit (Machery-Nagel, Duran, Germany) (11). The study was performed in 2010. Ethical approval was received from the Ethics Committee of the University of Zagreb School of Medicine.

Amplification was performed in a total volume of 11.5 μ L in Mastercycler gradient thermocycler (Eppendorf, Duran, Germany) as previously described (12,13). After amplification, the PCR products (1.5 μ L) were mixed with 3 μ L of loading buffer and 1 μ L of each of 2 internal size markers and applied on gel. On each gel, we also included one external, commercial size marker. Electrophoresis was performed using a 6% standard denaturing polyacrylamide gel in an automated laser fluorescence DNA sequencer (ALFexpress, Pharmacia Biotech, Uppsala, Sweden). The amplification products were analyzed and their relative fluorescent intensities calculated using AlleleLocator software (Pharmacia Biotech). The assignment of alleles was performed using allelic ladders.

The Hardy-Weinberg equilibrium test was performed by χ^2 test. Allele and genotype frequencies for each X-STR locus were determined by direct counting. The power of exclusion was calculated as described by Crow (14), while polymorphism information content (PIC) value was obtained as suggested by Hearne (15). The average power of discrimination (PD) was estimated as proposed by Desmarais (16).

RESULTS AND DISCUSSION

Allele frequencies of 5 X-linked microsatellites in the Croatian population were calculated separately for men and women, and for all participants together (Table 1), and no sex-related differences were found. DXS6803 locus was most polymorphic with 15 different alleles, while DXS6810 locus was least polymorphic with only 6 different alleles. The population study ($n = 183$) confirmed that all 5 STR loci were informative, which is in concordance with the data from other populations (17-19).

The observed statistical parameters of the examined X-linked markers are shown in Table 2. PIC was calculated from the combined data for men and women, whereas observed heterozygosity (Hobs), genotypes, matching probability, and power of exclusion were obtained only from data for women. No deviation from Hardy-Weinberg equilibrium was found. Four loci (DXS9895, GATA172D05, DXS6803, and HPRTB) showed high PIC value and high Hobs value. The PIC value of the examined markers ranged from 0.623 (DXS6810) to 0.901 (DXS6803). The DXS6810 showed a Hobs value lower than 0.60, which corresponds to the high frequency of the allele 5 in our sample.

Among the successfully processed 150 samples, sex chromosomal aneuploidies were diagnosed by stan-

TABLE 1. Distribution of allele frequencies at DXS6810, HPRTB, DXS9895, DXS6803, and GATA172D05 loci in the Croatian population (n = 183)

DXS6810		HPRTB		DXS9895		DXS6803		GATA172D05	
alleles	frequency	alleles	frequency	alleles	frequency	alleles	frequency	alleles	frequency
1	–	6	–	11	0.0254	9	0.0072	6	0.1413
2	0.0072	7	–	12	0.0797	10	0.0326	7	0.0290
3	0.0543	8	0.0072	13	0.2065	11	0.1993	8	0.1232
4	0.3261	9	0.0435	14	0.2174	11.3	0.0652	9	0.0870
5	0.4130	10	0.0543	15	0.2029	12	0.1884	10	0.2030
6	0.1812	11	0.1159	16	0.2174	12.3	0.1522	11	0.2500
7	0.0181	12	0.239	17	0.0435	13	0.1014	12	0.1449
		13	0.2790	18	0.0072	13.3	0.1703	13	0.0217
		14	0.1413			14	0.0036		
		15	0.1087			14.3	0.0580		
		16	0.0109			15.3	0.0181		
		17	–			16.3	0.0036		

TABLE 2. Statistical parameters for short tandem repeat markers HPRTB, DXS6810, DXS6803, GATA172D05, and DXS9895 obtained from the population study (n = 183).

	HPRTB	DXS6810	DXS6803	GATA172D05	DXS9895
Observed heterozygosity in women	0.656	0.591	0.849	0.731	0.656
Expected heterozygosity in men	0.797	0.609	0.895	0.850	0.781
Polymorphism information content	0.805	0.623	0.901	0.857	0.795
Power of discrimination in men	0.799	0.708	0.894	0.871	0.816
Average power of discrimination in women	0.925	0.740	0.979	0.958	0.918
Expected probability of exclusion in women	0.944	0.836	0.982	0.967	0.935
Matching probability	0.056	0.164	0.017	0.033	0.065

TABLE 3. Thirteen samples from patients with X chromosome disorders tested by 5 X-short tandem repeat markers

Sample	HPRTB	DXS6810	DXS6803	GATA172D05	DXS9895
Karyotype 45, X					
1.	12	5	12	12	13
2.	14	4	13.3	11	16
3.	12	5	11	11	16
4.	11	4	12.3	10	16
5.	13	5	12	11	14
6.	13	5	11	10	15
Karyotype 47, XXY					
7.	13	5	10, 12	10	13, 16
8.	12, 13	6	11, 12	8, 11	14, 17
9.	12	4	12.3, 13.3	10, 12	14, 16
10.	12, 13	3, 6	11, 12	10, 11	15, 16
11.	12	4, 5	11, 13.3	11	13, 14
Karyotype 47, XXX					
12.	12*, 13	4*, 5	11, 12*	10*, 12	15, 16, 17
13.	14	4, 5*	11.3, 12, 13	10, 11*	14, 15, 18

*Trisomic diallelic.

standard cytogenetic methods in 13 cases (Table 3). For each sample with X chromosome disorders, we also performed a PCR-STR analysis using these 5 STR loci. All 6 samples from patients with Turner syndrome had only one fluorescent peak (Table 2). At the same time, all 5 patients with karyogram 47, XXY had 2 alleles at DXS9895 and DXS6803, and 2 or 1 allele at the other 3 STR loci. It is important to mention that for all samples from this subgroup at least 2 tested X-microsatellites showed 2 different alleles. Both samples of Triple X syndrome demonstrated 3 alleles at DXS9895 and a double dose of 1 or 2 alleles at the remaining 4 loci (two times 1 or 2 alleles at the other 4 STR).

Our results are in agreement with the results of other studies demonstrating the usefulness of PCR-STR method for detection of X chromosomal disorders (9,10,20). The method is fast, sensitive, and suitable for screening of a large number of samples. In conclusion, data on X-chromosome STRs among Croatians would be useful for paternity testing and forensic purposes but also for X-chromosome

mapping studies. The set of 5 STR loci tested in the present study showed to be sufficient for the diagnosis of Turner, Klinefelter, and Triple X syndrome.

Funding From the project of the Ministry of Science, Education and Sport of the Republic of Croatia (No. 0108027).

Ethical approval Received from the Ethics Committee of the University of Zagreb School of Medicine.

Declaration of authorship KCG wrote the article and participated in all investigations. ZG participated in all investigations. KŠ participated in all investigations. ITĐ participated in all investigations. DB participated in all investigations.

Competing interests: All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- Mansfield ES. Diagnosis of Down syndrome and other aneuploidies using quantitative polymerase chain reaction and small tandem repeat polymorphisms. *Hum Mol Genet.* 1993;2:43-50. [Medline:8490622](#) [doi:10.1093/hmg/2.1.43](#)
- Pertl B, Yau SC, Sherlock J, Davies AF, Mathew CG, Adinolfi M. Rapid molecular method for prenatal detection of Down's syndrome. *Lancet.* 1994;343:1197-8. [Medline:7909872](#) [doi:10.1016/S0140-6736\(94\)92404-X](#)
- Pertl B, Weitgasser U, Kopp S, Kroisel PM, Sherlock J, Adinolfi M. Rapid detection of trisomies 21 and 18 and sexing by quantitative fluorescent multiplex PCR. *Hum Genet.* 1996;98:55-9. [Medline:8682507](#) [doi:10.1007/s004390050159](#)
- Alford RL, Hammond HA, Coto I, Caskey CT. Rapid and efficient resolution of parentage by amplification of short tandem repeats. *Am J Hum Genet.* 1994;55:190-5. [Medline:8023845](#)
- Hammond HA, Jin L, Zhong Y, Caskey CT, Chakraborty R. Evaluation of 13 short tandem repeat loci for use in personal identification applications. *Am J Hum Genet.* 1994;55:175-89. [Medline:7912887](#)
- Pertl B, Kopp S, Kroisel PM, Häusler M, Sherlock J, Winter R, et al. Quantitative fluorescence polymerase chain reaction for the rapid prenatal detection of common aneuploidies and fetal sex. *Am J Obstet Gynecol.* 1997;177:899-906. [Medline:9369843](#) [doi:10.1016/S0002-9378\(97\)70292-8](#)
- Adinolfi M, Cirigliano V. Detection of fetal cells in transcervical samples using X22 marker. *J Med Genet.* 2000;37:E1. [Medline:10807700](#) [doi:10.1136/jmg.37.5.e1](#)
- Mann K, Fox SP, Abbs SJ, Yau SC, Scriven PN, Docherty Z, et al. Development and implementation of a new rapid aneuploidy diagnostic service within the UK National Health Service and implications for the future of prenatal diagnosis. *Lancet.* 2001;358:1057-61. [Medline:11589937](#) [doi:10.1016/S0140-6736\(01\)06183-9](#)
- Cirigliano V, Ejarque M, Fuster C, Adinolfi M. X chromosome dosage by quantitative fluorescent PCR and rapid prenatal diagnosis of sex chromosome aneuploidies. *Mol Hum Reprod.* 2002;8:1042-5. [Medline:12397218](#) [doi:10.1093/molehr/8.11.1042](#)
- Donaghue C, Roberts A, Mann K, Ogilvie CM. Development and targeted application of a rapid QF-PCR test for sex chromosome imbalance. *Prenat Diagn.* 2003;23:201-10. [Medline:12627420](#) [doi:10.1002/pd.569](#)
- Vogelstein B, Gillespie D. Preparative and analytical purification of DNA from agarose. *Proc Natl Acad Sci U S A.* 1979;76:615-9. [Medline:284385](#) [doi:10.1073/pnas.76.2.615](#)
- Huang D, Yang Q, Yu C, Yang R. Development of the X-linked tetrameric microsatellite markers HumDXS6803 and HumDXS9895 for forensic purpose. *Forensic Sci Int.* 2003;133:246-9. [Medline:12787659](#) [doi:10.1016/S0379-0738\(03\)00077-X](#)
- Shin SH, Yu JS, Park SW, Min GS, Chung KW. Genetic analysis of 18 X-linked short tandem repeat markers in Korean population. *Forensic Sci Int.* 2005;147:35-41. [Medline:15541590](#) [doi:10.1016/j.forsciint.2004.04.012](#)
- Crow JF, Kimura M. An introduction to population genetics theory. New York, Evanston and London: Harper and Row; 1970.
- Hearne CM, Ghosh S, Todd JA. Microsatellites for linkage analysis of genetic traits. *Trends Genet.* 1992;8:288-94. [Medline:1509520](#)
- Desmarais D, Zhong Y, Chakraborty R, Perreault C, Busque L. Development of a highly polymorphic STR marker for identity testing purposes at the human androgen receptor gene (HUMARA). *J Forensic Sci.* 1998;43:1046-9. [Medline:9729823](#)
- Edelmann J, Szibor R. The X-linked STRs DXS7130 and DXS6803. *Forensic Sci Int.* 2003;136:73-5. [Medline:12969622](#) [doi:10.1016/S0379-0738\(03\)00257-3](#)
- Ribeiro Rodrigues EM, Leite FP, Hutz MH, Palha Tde J, Ribeiro dos Santos AK, dos Santos SE. A multiplex PCR for 11 X chromosome STR markers and population data from a Brazilian Amazon Region. *Forensic Sci Int Genet.* 2008;2:154-8. [Medline:19083811](#) [doi:10.1016/j.fsigen.2007.10.179](#)
- Chr X-STR. org 2.0. Available from: <http://www.chrx-str.org/>. Accessed: May 12, 2011.
- Pertl B, Pieber D, Lercher-Hartlieb A, Orescovic I, Häusler M, Winter R, et al. Rapid prenatal diagnosis of aneuploidy by quantitative fluorescent PCR on fetal samples from mothers at high risk for chromosome disorders. *Mol Hum Reprod.* 1999;5:1176-9. [Medline:10587374](#) [doi:10.1093/molehr/5.12.1176](#)