

Interaction of genetic risk factors confers increased risk for metabolic syndrome: the role of peroxisome proliferator-activated receptor γ

Božina, Tamara; Sertić, Jadranka; Lovrić, Jasna; Jelaković, Bojan; Šimić, Iveta; Reiner, Željko

Source / Izvornik: **Genetic Testing and Molecular Biomarkers**, 2014, 18, 32 - 40

Journal article, Accepted version

Rad u časopisu, Završna verzija rukopisa prihvaćena za objavljivanje (postprint)

<https://doi.org/10.1089/gtmb.2013.0344>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:478688>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-05-21**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine](#)
[Digital Repository](#)





Središnja medicinska knjižnica

Božina T., Sertić J., Lovrić J., Jelaković B., Šimić I., Reiner Ž. (2014)
Interaction of genetic risk factors confers increased risk for metabolic syndrome: the role of peroxisome proliferator-activated receptor γ .
Genetic Testing and Molecular Biomarkers, 18(1). pp. 32-40. ISSN 1945-0265

<http://online.liebertpub.com/gte>

<http://dx.doi.org/10.1089/gtmb.2013.0344>

<http://medlib.mef.hr/2256>

University of Zagreb Medical School Repository

<http://medlib.mef.hr/>

Interaction of Genetic Risk Factors Confers Increased Risk for Metabolic Syndrome: The Role of PPAR γ

Božina T^a, Sertić J^{a,b}, Lovrić J^a, Jelaković B^c, Šimić I^d, Reiner Ž^d

^a *Department of Medical Chemistry, Biochemistry and Clinical Chemistry, School of Medicine, University of Zagreb, Šalata 3, 10 000 Zagreb, Croatia*

^b *Department of Laboratory Diagnostics, University Hospital Center, Kišpatićeva 12, 10000 Zagreb, Croatia*

^c *Department of Internal Medicine, Division of Nephrology and Arterial Hypertension; University Hospital Center, Kišpatićeva 12, 10000 Zagreb, Croatia*

^d *Department of Internal Medicine, Division of Metabolic Diseases, University Hospital Center, Kišpatićeva 12, 10000 Zagreb, Croatia*

Corresponding author

Professor Jadranka Sertić PhD

Department of Laboratory Diagnostics, University Hospital Center Zagreb,

Kišpatićeva 12, 10000 Zagreb, Croatia

Phone: +385 1 2367248

Fax: +385 1 2367395

Email: jadranka.sertic@kbc-zagreb.hr

Running title

Genetic Risk Factors and Metabolic Syndrome

Abstract

The aim of the study was to estimate the influence of interactions between peroxisome proliferator-activated receptor γ (*PPAR* γ) and target genes lipoprotein lipase (*LPL*), interleukin 6 (*IL6*), angiotensin converting enzyme (*ACE*), and angiotensin II type 1 receptor (*AT1R*) on metabolic syndrome (MetSy) and its traits.

Methods: The study included 527 participants (263 with MetSy and 264 controls). Genotyping of *PPAR* γ Pro12Ala, *LPL* PvuII (-/+), *IL6* -174G>C, *ACE* I/D and *AT1R* 1166A>C was performed using polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) based methods.

Results: Interaction between *PPAR* γ Pro12Ala and *LPL* (Pvu-/+) improved prediction of MetSy over and above prediction based on a model containing no interactions ($\chi^2=7.22$; df=1; p=0.007). In the group of participants with *PPAR* γ Pro12Ala or Ala12Ala genotypes, those with *LPL* Pvu (-/+) or (+/+) genotype had greater odds for MetSy (OR=5.98; 95% CI: 1.46-24.47, p=0.013). Interaction between *PPAR* γ Pro12Ala and *IL6* -174G>C improved prediction of high fasting blood glucose ($\chi^2=13.99$; df=1; p<0.001). *PPAR* γ Ala12 variant was found protective in patients with *IL6* -174GG genotype (OR=0.10; 95% CI: 0.02-0.57, p=0.01), while in the case of *IL6* -174C allele carriers, for *PPAR* γ Ala12 carriers larger odds for high glucose levels compared with Pro12 variant were observed (OR=2.39; 95% CI: 1.11-5.17, p=0.026). Interactions of *PPAR* γ and *ACE* were significant for BMI. In the group with *ACE* DD genotype, those with *PPAR* γ Pro12Ala or Ala12Ala genotype have greater odds for obesity (OR=9.98; 95% CI: 1.18-84.14, p=0.034). Conclusions: *PPAR* γ gene variants can, in interaction with some of its target genes, modulate physiological processes leading to the development of MetSy.

Key words: Metabolic syndrome, Genetic polymorphisms, Peroxisome proliferator-activated receptor γ gene, Lipoprotein lipase, Interleukin 6, Angiotensin converting enzyme, Angiotensin II type 1 receptor

INTRODUCTION

The pathogenesis of metabolic syndrome (MetSy) has been associated with the effect of genetic predisposition in combination with environmental factors (Lusis et al. 2008; Lin et al., 2005). Whereas the importance of genetic factors for the development of MetSy has been widely recognized, the contribution of genes has not yet been fully clarified. Many association studies have been conducted with inconsistent results (Monda et al., 2010; Park et al. 2009). Considering the central role of adipose tissue in MetSy, different adipocyte-related genes have been studied as possible candidates in MetSy, including peroxisome proliferator-activated receptor γ (*PPAR γ*), lipoprotein lipase (*LPL*), renin-angiotensin system (RAS)-related genes - angiotensin converting enzyme (*ACE*) and angiotensin II type 1 receptor (*AT1R*), and interleukin 6 (*IL6*), which exhibit considerable variability.

PPARs are transcription factors implicated in different biological pathways ranging from lipid and glucose homeostasis and insulin sensitization, to control of cell proliferation/differentiation, inflammation, and immunity (Sharma and Staels, 2007; Olefsky, 2000). Endogenous ligands are thought to bind *PPAR γ* and promote downstream gene target transcription (Ide et al., 2003; Bell-Parikh et al., 2003). Because of alternative mRNA splicing, two protein isoforms occur: *PPAR γ 1* and *PPAR γ 2*. While *PPAR γ 1* is found ubiquitously in the body, *PPAR γ 2* is largely found in adipose tissue. Studies have demonstrated that *PPAR γ* has a critical role in regulating adipocyte differentiation and lipid accumulation (Rangwala and Lazar, 2009). The *PPAR γ* Pro12Ala variant (rs1801282) has been found to modulate transcriptional activity and has a reduced affinity for the response element in target genes, which leads to the less efficient stimulation of *PPAR γ* target genes (He, 2009). The frequency of the 12Ala allele ranges from 2% to 18% in healthy people (Paracchini et al., 2005). *PPAR γ* function is a key factor in mediating conditions such as dyslipidemia, obesity, and insulin resistance (Huang et al., 2011; Gouda et al., 2010), and has also been implicated in the development of cardiovascular diseases (Azhar, 2010).

LPL plays the major role in the metabolism and transport of lipids. Distinct physiological activities of *LPL* together regulate the supply of fatty acids to various tissues for either storage or oxidation. Insulin has a major effect on *LPL* activity in adipose tissue during adipocyte differentiation by increasing *LPL* gene transcription (Semenkovich et al., 1989). Glucose also increases adipose tissue *LPL* activity. Several mutation loci have been detected in *LPL* gene and investigated for their associations with plasma lipid level and the development of cardiovascular diseases (Angelakopoulou et al., 2012). Polymorphism *LPL* *PvuII* (-/+) (rs285) is caused by the absence (-) or presence (+) of a C>T transition at position 497 in intron 6, and may interfere with correct splicing of mRNA, diminishing the enzyme activity.

Interleukin 6 (*IL6*) is a pleiotropic inflammatory cytokine derived from diverse tissues. In chronic inflammation it has rather proinflammatory properties. There is evidence that variations in *IL6* gene are associated with cytokine and

metabolic modulation, leading to impaired glucose and lipid homeostasis and increasing cardiometabolic risk (Stephens et al., 2004). Polymorphism -174G>C (rs1800795) is most prevalent and of significant biological importance since it affects the IL6 transcription (Curti et al., 2011). The frequencies of -174G>C variants were estimated to be 0.57 in Caucasians and 0.93 in Afro-Americans (Huang et al., 2007). Higher circulating IL6 levels have been associated with obesity and visceral fat deposition (Qi et al., 2007; Stephens et al., 2007), increased risk of impaired glucose tolerance, type 2 diabetes mellitus (T2DM) (Sattar et al., 2003), and high blood pressure (Fernandez-Real et al., 2001).

PPARs modulate RAS by transcriptional control of all its components (Roszer and Ricote, 2010). Polymorphism in *ACE* and *AT1R* gene has been found significant for variability in many pathophysiological processes connected with RAS, including hypertension (Tiret et al., 1998), diabetes, and cardiovascular disease (Pujia et al., 1994). RAS might be involved in the pathophysiology of obesity (Engeli et al., 2000). The most common *ACE* gene polymorphism is the insertion/deletion (I/D) (rs4646994) of 287-bp *Alu* repeats located in intron 16.

A nucleotide substitution (1166A>C) in 3'-UTR of the *AT1R* gene (rs1272176) results in increased expression of the receptor gene (Bonnardeaux et al., 1994). The increased frequency of the *AT1R* 1166C allele has been associated with essential hypertension (Tiret et al., 1998), cardiac hypertrophy (Pujia et al., 1994), myocardial infarction (Engeli et al., 2000). But opposite findings have also been published where *AT1R* 1166CC genotype predisposes to favorable anthropometric and metabolic traits relative to cardiovascular risk (Möller et al., 2008).

The influence of gene variants on phenotype expression seems to be population specific because of possible differences in environmental factors and genetic background (Groop, 2000).

In this paper we consider the combined effects of *PPARγ*, *LPL*, *IL6*, *ACE* and *AT1R* that might confer a higher risk for MetSy or for some of its traits than individual gene variants in Croatian population.

MATERIAL AND METHODS

Subjects

A total of 527 subjects (343 female, 184 male) of Croatian origin participated in the study, including 265 patients with MetSy and 262 controls (without MetSy criteria). Participants with MetSy were recruited among patients admitted to the University Hospital Center Zagreb, and control subjects were selected among hospital staff and other Zagreb citizens appointed for routine check up. The study was performed in the period between March 2010 and October 2012. MetSy was defined using the criteria established in the third report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment in Adults (NCEP-ATPIII, 2001). An individual with a combination of any three or more of the following risk factors was classified as having MetSy: waist circumference (WC), male >102 cm, female >88 cm; TG ≥1.7 mmol/L; HDL-C <1.0 in men and <1.3 mmol/L in women; systolic

blood pressure (SBP) or diastolic blood pressure (DBP) $\geq 130/85$ mm Hg, or use of anti-hypertensive medications; and fasting blood glucose ≥ 6.1 mmol/L or use of antidiabetic medication with the age of diagnosis of T2DM ≥ 40 years. In the case of participants' use of medications, corrections of medicated traits were made (Kraja et al., 2006). For control group we selected healthy subjects without MetSy and without any other serious illness. Data were collected on clinical variables including age, height, weight, and WC. Body mass index (BMI) was calculated as weight (kg)/height (m^2). Participants with BMI between 25-30 kg/m^2 were considered overweight, and those with BMI > 30 kg/m^2 are obese. Blood samples for biochemical analyses (total cholesterol, TG, LDL-C, HDL-C and glucose) were collected after overnight fasting and were analyzed by using routine laboratory methods. All participants signed informed consent forms, and the study protocol was approved by Ethics Committee of the University Hospital Center Zagreb.

Genotyping

Genomic DNA was extracted from leukocytes using the salting out procedure (Miller et al., 1988), and genotyping of *PPAR γ* Pro12Ala, *LPL* PvuII (-/+), *IL6* -174G>C, *ACE* I/D and *AT1R* 1166A>C was performed according to previously published methods based on PCR or PCR-RFLP procedures (Oh et al., 2000; Xu et al., 2008; Jamie et al., 2005; Rigat et al., 1992; Hilgers et al., 1999, respectively). PCR amplifications were performed in 25 μL final volume in GeneAmp PCR System 9600 (Applied Biosystems, USA).

Statistical analysis

The level of statistical significance was set to 5% ($p < 0.05$), and in all instances two-tailed tests of statistical significance were used. Means and standard deviations were used as measures of central tendency and variability for continuous variables, and independent samples t-test was used for their comparison. If variances were heterogeneous, t-test for unequal variances with corrected degrees of freedom was used. Univariate and multivariate prediction of MetSy, its traits and BMI was carried out by means of logistic regression, and odds ratios with 95% confidence intervals were given for each variable. To examine gene-gene interactions, total predictive model was built using hierarchical backward elimination approach (Kleinbaum and Klein, 2002), with initial model containing all variables and all possible pairs of gene-gene interactions. Interaction with the highest level of statistical significance from the full model was considered first so that logistic regression model without that interaction was compared to the model containing that interaction. If the difference between two models was not statistically significant, interaction was dropped out of the model and the procedure was repeated for the least significant interaction from the reduced model. If gene-gene interaction proved to be statistically significant, regression coefficients for each polymorphism at each of the categories

of the other one were examined by redefining their reference groups, since coefficients of variables in interaction are conditioned to the reference group of the other variable from that interaction (Jaccard, 2001). The analyses were carried out using SPSS 17.0 (SPSS Inc., Chicago, IL, USA) statistical software package.

RESULTS

Baseline characteristics

Baseline characteristics of study participants are given in Table 1. Patients had significantly higher BMI, WC, higher levels of TG, total cholesterol, LDL-C, glucose and significantly higher blood pressure and lower levels of HDL-C compared to controls ($p < 0.001$). MetSy and control group did not differ significantly with respect to age and gender.

Table 1. Baseline characteristics of study participants

	<i>Patients</i>	<i>Controls</i>	<i>P</i>
	<i>Mean +/- SD</i>	<i>Mean +/- SD</i>	
Age (yrs)	53.3 +/- 10.51	53.1 +/- 10.72	0.874
Gender (N of male participants/All)	99/265	85/262	0.238
Body mass index (kg/m ²)	32.1 +/- 4.77	24.6 +/- 3.44	<0.001
Waist circumference (cm)	107.3 +/- 11.15	85.1 +/- 11.42	<0.001
Triglycerides (mmol/L)	2.82 +/- 2.21	1.21 +/- 0.73	<0.001
Cholesterol (mmol/L)	6.0 +/- 1.11	5.5 +/- 1.02	<0.001
High-density lipoprotein cholesterol (mmol/L)	1.22 +/- 0.33	1.69 +/- 0.41	<0.001
Low-density lipoprotein cholesterol (mmol/L)	3.7 +/- 1.14	3.3 +/- 0.88	<0.001
Glucose (mmol/L)	6.24 +/- 2.21	5.0 +/- 0.77	<0.001
High Blood pressure (N of subjects with high BP/All)	235/260	71/255	<0.001
Systolic blood pressure	152.4 +/- 26.0	124.8 +/- 14.24	<0.001
Diastolic blood pressure	94.4 +/- 15.87	79.5 +/- 8.60	<0.001

Differences between groups were evaluated by t-test, Mann-Whitney test or chi-square test depending on distribution normality. Significant values are indicated in bold. $P < 0.05$ was considered statistically significant.

Associations of *PPAR* γ , *LPL*, *IL6*, *ACE* and *AT1R* gene variants with MetSy or its trait

We found no differences in *PPAR* γ Pro12Ala, *LPL* PvuII (-/+), *IL6* 174G>C, *ACE* I/D and *AT1R* 1166A>C genotype distributions between MetSy cases and controls (Table 2). We found no departure from the Hardy-Weinberg equilibrium.

Table 2. Genotype frequencies in case and control group

Gene/Genotype	Patients n(%)	Controls n(%)	OR (95% CI)
<i>ACE</i>			
DD	72 (27.2)	81 (30.9)	1
ID	132 (49.8)	130 (49.6)	1.14 (0.77-1.70)
II	61 (23.0)	51 (19.5)	1.35 (0.83-2.19)
<i>PPAR</i>			
Pro12Pro	200 (75.5)	199 (76.0)	1
Pro12Ala	62 (23.4)	60 (22.9)	1.03 (0.69-1.53)
Ala12Ala	3 (1.1)	3 (1.1)	1.03 (0.69-1.53)
<i>IL6</i>			
GG	85 (32.1)	91 (34.7)	1
GC	130 (49.1)	137 (52.3)	1.02 (0.69-1.49)
CC	50 (18.9)	34 (13.0)	1.57 (0.93-2.67)
<i>AT1R</i>			
AA	136 (51.3)	134 (51.1)	1
AC	114 (43.0)	110 (42.0)	1.02 (0.72-1.46)
CC	15 (5.7)	18 (6.9)	0.82 (0.40-1.70)
<i>LPL</i>			
-/-	62 (23.4)	71 (27.1)	1
-/+	124 (46.8)	132 (50.4)	1.08 (0.71-1.64)
+/+	79 (29.8)	59 (22.5)	1.53 (0.95-2.48)

ABBREVIATIONS: OR = univariate odds ratio; 95% CI = 95% confidence interval for odds ratio

When tested for the influence of interactions between *PPAR γ* and its target genes *LPL*, *IL6*, *ACE* and *AT1R* on MetSy and its traits, separately in the group with and without MetSy, several statistically significant interactions were observed (Table 3).

Table 3. Hierarchical backward elimination of gene-gene interactions

	Least significant interaction / feature	P (interaction)	P (difference between models)
Full model*			
Reduced model**			
1	<i>PPARγ</i> x <i>LPL</i> PvuII / MetSy	0.008	0.007
2	<i>PPARγ</i> x <i>ACE</i> I/D / BMI	0.010	0.004
3	<i>PPARγ</i> x <i>IL6</i> -174G>C / Glucose	0.001	<0.001

ABBREVIATIONS: MetSy- metabolic syndrome, BMI-body mass index, p (interaction) = level of statistical significance for interaction, results of Wald χ^2 test; p (difference between models) = level of statistical significance, comparison of model without interaction and model with interaction

*Full model with all possible pairs of interactions included

**Model from which interaction with highest level of statistical significance from previous model was excluded

Interaction of PPAR γ and LPL and MetSy

Interaction between *PPAR γ* Pro12Ala and *LPL* PvuII (-/+) significantly improved prediction of MetSy, over and above prediction based on model containing no interactions ($\chi^2=7.22$; df=1; p=0.008) (Table 4).

In the group of participants with *LPL* PvuII (-/-) genotype, participants with *PPAR γ* Pro12Ala or Ala12Ala genotype had smaller odds for MetSy compared to those with Pro12Pro genotype (OR=0.12; 95% CI: 0.03-0.52, p=0.005). In the group of those participants with *PPAR γ* Pro12Ala or Ala12Ala genotypes, those with *LPL* PvuII (-/+) or PvuII (+/+) genotype had greater odds for MetSy (OR=5.98; 95% CI: 1.46-24.47, p=0.013).

In multivariate prediction, when adjusted for all other variables, age (p=0.012) and BMI (p<0.001) were also statistically significantly associated with MetSy. With each year increase in age, odds for MetSy decrease by 3% (OR = 0.97; 95% CI = 1.54 – 1.85) and with each unit increase in BMI the odds for MetSy increase more than 1.5 times (OR = 1.69; 95% CI = 1.54 – 1.85).

Table 4. Total predictive model for MetSy with all polymorphisms

	Patients n(%)	Controls n(%)	OR _{mv} (95% CI)	P
Age*	53.3 (10.51)	53.1 (10.72)	0.97 (0.94-0.99)	0.012
Gender				
female	166 (62.6)	177 (67.6)	1	
male	99 (37.4)	85 (32.4)	1.27 (0.70-2.29)	0.427
total	265 (100.0)	262 (100.0)		
ACE				
DD	72 (27.2)	81 (30.9)	1	
ID or II	193 (72.8)	181 (69.1)	1.04 (0.56-1.92)	0.909
total	265 (100.0)	262 (100.0)		
<i>PPARγ</i> if <i>LPL</i> Pvu (-/-)				
Pro12Pro	50 (80.6)	52 (73.2)	1	
Pro12Ala or Ala12Ala	12 (19.4)	19 (26.8)	0.12 (0.03-0.52)	0.005
total	62 (100.0)	71 (100.0)		
<i>PPARγ</i> if <i>LPL</i> (-/+) or (+/+)				
Pro12Pro	150 (73.9)	147 (77.0)	1	
Pro12Ala or Ala12Ala	53 (26.1)	44 (23.0)	1.11 (0.54-2.25)	0.781
total	203 (100.0)	191 (100.0)		
<i>IL6</i>				
GG	85 (32.1)	91 (34.7)	1	
GC or CC	180 (67.9)	171 (65.3)	0.80 (0.45-1.44)	0.803
total	265 (100.0)	262 (100.0)		
<i>AT1R</i>				
AA	136 (51.3)	134 (51.1)	1	
AC or CC	129 (48.7)	128 (48.9)	0.69 (0.39-1.20)	0.185
total	265 (100.0)	262 (100.0)		
<i>LPL</i> PvuII if <i>PPARγ</i> Pro12Pro				

(-/-)	50 (25.0)	52 (26.1)	1	
(-/+) or (+/+)	150 (75.0)	147 (73.9)	0.67 (0.32-1.41)	0.290
total	200 (100.0)	199 (100.0)		
<i>LPL</i> PvuII if <i>PPAR</i> γ Pro12Ala or Ala12Ala				
(-/-)	12 (18.5)	19 (30.2)	1	
(-/+) or (+/+)	53 (81.5)	44 (69.8)	5.98 (1.46-24.47)	0.013
total	65 (100.0)	63 (100.0)		
BMI*	32.1 (4.77)	24.6 (3.44)	1.69 (1.54-1.85)	<0.001
Total cholesterol*	6.0 (1.11)	5.5 (1.02)	1.32 (0.82-1.12)	0.248
LDL cholesterol*	3.7 (1.14)	3.3 (0.88)	1.41 (0.86-2.31)	0.170
<i>PPAR</i> γ x <i>LPL</i>			8.92 (1.77-44.88)	0.008

ABBREVIATIONS: OR_{mv} = multivariate odds ratio; 95% CI = 95% confidence interval for odds ratio; *Mean (standard deviation) Significant values are indicated in bold. P<0.05 was considered statistically significant.

Interaction of PPAR γ and IL6 and fasting blood glucose

Interaction of *PPAR* γ Pro12Ala and *IL6* -174G>C statistically significantly improved prediction of high fasting blood glucose in the group of MetSy patients, over and above model containing no interactions ($\chi^2=13.99$; df=1; p=0.001). Inclusion of this interaction to the model containing no interactions increased Nagelkerke R square from 0.21 to 0.27 (Table 5). In the group of patients with *IL6* -174GG genotype, *PPAR* γ Pro12Ala or Ala12Ala genotype carriers (taken together) had smaller odds of having high glucose compared to Pro12Pro genotype carriers (OR=0.10; 95% CI: 0.02-0.57, p=0.01). In the group of patients with *IL6* -174GC or CC genotype, *PPAR* γ Pro12Ala or Ala12Ala genotype carriers (taken together) had larger odds of having high glucose compared to Pro12Pro genotype carriers (OR=2.39; 95% CI: 1.11-5.17, p=0.026). In the group of patients with *PPAR* γ Pro12Pro genotype, *IL6* -174GC or CC genotype carriers (taken together) had smaller odds of having high glucose compared to *IL6* -174GG genotype carriers (OR=0.41; 95% CI: 0.20-0.83, p=0.013). In multivariate prediction, age (p=0.001), gender (p=0.045), and TGs (p=0.001) were also associated with high fasting blood glucose. With each year increase in age, odds for high glucose increase by 6% (OR = 1.06; 95% CI = 1.02 – 1.09), and male patients had more than two times larger odds of having high glucose (OR = 2.08; 95 % CI = 1.02 – 4.26). Each unit increase in TGs increases the odds for high glucose 1.3 times (OR = 1.3; 95% CI = 1.11 – 1.53).

Table 5. Multivariate prediction of high fasting blood glucose in MetSy group

	High n(%)	Normal n(%)	OR _{mv} (95% CI)	P
Age*	56.3 (9.25)	51.6 (10.87)	1.06 (1.02-1.09)	0.001
Gender				
female	52 (53.6)	113 (68.5)	1	
male	45 (46.4)	52 (31.5)	2.08 (1.02-4.26)	0.045
total	97 (100.0)	165 (100.0)		
ACE				
DD	24 (24.7)	47 (28.5)	1	
ID or II	73 (75.3)	118 (71.5)	1.92 (0.95-3.89)	0.071
total	97 (100.0)	165 (100.0)		
PPAR γ if IL6 -174GG				
Pro12Pro	33 (94.3)	35 (70.0)	1	
Pro12Ala or Ala12Ala	2 (5.7)	15 (30.0)	0.10 (0.02-0.57)	0.010
total	35 (100.0)	50 (100.0)		
PPAR γ if IL6 -174GC/CC				
Pro12Pro	39 (62.9)	90 (78.3)	1	
Pro12Ala or Ala12Ala	23 (37.1)	25 (21.7)	2.39 (1.11-5.17)	0.026
total	62 (100.0)	115 (100.0)		
IL6 if PPAR γ Pro12Pro				
GG	33 (45.8)	35 (28.0)	1	
GC or CC	39 (54.2)	90 (72.0)	0.41 (0.20-0.83)	0.013
total	72 (100.0)	125 (100.0)		
IL6 if PPAR γ ProAla/AlaAla				
GG	2 (8.0)	15 (37.5)	1	
GC or CC	23 (92.0)	25 (62.5)	10.13 (1.66-61.89)	0.012
total	25 (100.0)	40 (100.0)		
AT1R A1166C				
AA	49 (50.5)	86 (52.1)	1	
AC or CC	48 (49.5)	79 (47.9)	0.96 (0.53-1.74)	0.893
total	97 (100.0)	165 (100.0)		
LPL PvuII				
(-/-)	21 (21.6)	39 (23.6)	1	
(-/+) or (+/+)	76 (78.4)	126 (76.4)	1.01 (0.49-2.05)	0.989
total	97 (100.0)	165 (100.0)		
BMI	32.3 (5.06)	32.1 (4.65)	1.01 (0.92-1.12)	0.781
Total cholesterol	5.9 (1.05)	6.0 (1.14)	0.73 (0.56-1.12)	0.214
LDL cholesterol	3.7 (1.01)	3.8 (1.22)	1.14 (0.84-1.55)	0.400
Waist circumference	108.7 (11.11)	106.4 (11.07)	1.01 (0.97-1.06)	0.615
Triglycerides	3.1 (2.97)	2.7 (1.60)	1.30 (1.11-1.53)	0.001
HDL cholesterol	1.3 (0.37)	1.2 (0.31)	3.60 (0.91-2.23)	0.101
Systolic blood pressure	156.5 (28.40)	150.3 (24.32)	1.02 (1.00-1.03)	0.120
Diastolic blood pressure	95.1 (16.01)	94.3 (15.65)	0.98 (0.95-1.02)	0.330
PPAR γ x IL6			24.81 (3.49-176.38)	0.001

ABBREVIATIONS: OR_{mv} = multivariate odds ratio; 95% CI = 95% confidence interval for odds ratio; *Mean (standard deviation)
Significant values are indicated in bold. P<0.05 was considered statistically significant.

Interaction of PPAR γ and ACE and obesity

PPAR γ Pro12Ala and ACE I/D gene interaction statistically significantly improved prediction of obesity in the group of MetSy patients, over and above prediction based on model containing no interactions ($\chi^2=8.08$; $df=1$; $p=0.004$). Inclusion of this interaction to the model containing no interactions increased Nagelkerke R square from 0.51 to 0.53 (Table 6). In the group of patients with PPAR γ Pro12Ala or Ala12Ala genotype, patients with ACE ID or II genotype had 89% smaller odds for obesity compared to those with DD genotype (OR=0.11; 95% CI: 0.01-0.88, $p=0.038$). In the group of those patients with ACE DD genotype, those with PPAR γ Pro12Ala or Ala12Ala genotype had about ten times greater odds for obesity (OR=9.98; 95% CI: 1.18-84.14, $p=0.034$).

In multivariate prediction, gender ($p=0.009$), WC ($p<0.001$), and HDL-C ($p=0.043$) were also associated with obesity. Male patients had smaller odds for obesity (OR = 0.30; 95% CI = 0.12 – 0.74). With each unit increase in WC the odds for obesity increase 1.26 times (OR = 1.26; 95% CI = 1.18 – 1.34), and with each unit increase in HDL-C the odds for obesity decrease 76% (OR = 0.24; 95% = 0.06 – 0.96).

Table 6. Multivariate prediction of obesity in MetSy group

	Yes	No	OR _{mv} (95% CI)	P
Age*	53.1 (9.97)	53.7 (11.57)	0.99 (0.96-1.03)	0.582
Gender				
female	104 (59.8)	61 (67.8)	1	
male	70 (40.2)	29 (32.2)	0.30 (0.12-0.74)	0.009
total	174 (100.0)	90 (100.0)		
ACE if PPAR γ Pro12Pro				
DD	32 (24.6)	20 (29.0)	1	
ID or II	98 (75.4)	49 (71.0)	2.36 (0.93-6.01)	0.072
total	130 (100.0)	69 (100.0)		
ACE if PPAR γ Pro12Ala/Ala12Ala				
DD	18 (40.9)	2 (9.5)	1	
ID or II	26 (59.1)	19 (90.5)	0.11 (0.01-0.88)	0.038
total	44 (100.0)	21 (100.0)		
PPAR γ if ACE DD				
Pro12Pro	32 (64.0)	20 (90.9)	1	
Pro12Ala or Ala12Ala	18 (36.0)	2 (9.1)	9.98 (1.18-84.14)	0.034
total	50 (100.0)	22 (100.0)		
PPAR γ if ACE ID/DD				
Pro12Pro	98 (79.0)	49 (72.1)	1	
Pro12Ala or Ala12Ala	26 (21.0)	19 (27.9)	0.44 (0.17-1.15)	0.094
total	124 (100.0)	68 (100.0)		
IL6				
GG	51 (29.3)	33 (36.7)	1	
GC or CC	123 (70.7)	57 (63.3)	1.16 (0.55-2.46)	0.698

total	174 (100.0)	90 (100.0)		
<i>ATIR</i>				
AA	90 (51.7)	45 (50.0)	1	
AC or CC	84 (48.3)	45 (50.0)	1.14 (0.56-2.31)	0.722
total	174 (100.0)	90 (100.0)		
<i>LPL</i> PvuII				
(-/-)	43 (24.7)	18 (20.0)	1	
(-/+) or (+/+)	131 (75.3)	72 (80.0)	0.77 (0.33-1.81)	0.550
total	174 (100.0)	90 (100.0)		
Total cholesterol*	5.9 (1.11)	6.1 (1.12)	0.91 (0.49-1.70)	0.777
LDL cholesterol*	3.7 (1.15)	3.8 (1.13)	0.93 (0.50-1.74)	0.813
Waist circumference*	111.7 (10.44)	98.9 (6.83)	1.26 (1.18-1.34)	<0.001
Triglycerides*	2.91 (2.27)	2.6 (2.12)	1.04 (0.83-1.30)	0.730
HDL cholesterol*	1.2 (0.32)	1.3 (0.36)	0.24 (0.06-0.96)	0.043
Systolic blood pressure*	153.7 (26.50)	149.7 (25.13)	1.02 (0.99-1.04)	0.197
Diastolic blood pressure*	94.8 (15.09)	93.2 (17.17)	0.99 (0.96-1.03)	0.719
Fasting blood glucose*	6.3 (2.28)	6.1 (2.09)	0.90 (0.73-1.11)	0.309
<i>PPARγ</i> x <i>ACE</i> I/D			0.04 (0.004-0.47)	0.010

ABBREVIATIONS: OR_{mv} = multivariate odds ratio; 95% CI = 95% confidence interval for odds ratio; *Mean (standard deviation)
Significant values are indicated in bold. P<0.05 was considered statistically significant.

Single gene correlations in the control group

1. In multivariate prediction *PPARγ* Pro12Ala or Ala12Ala genotype carriers had almost four times larger odds of having high WC compared to those with *PPARγ* Pro12Pro genotype carriers (OR=3.86; 95% CI: 1.24-12.06; p = 0.020) (Table 7).
2. In multivariate prediction *ATIR* 1166AC or CC genotype carriers had about eight times larger odds of having high TGs compared to *ATIR* AA genotype carriers (OR=8.05; 95% CI: 1.85-35.0; p = 0.005).

Table 7. Associations of single gene variants in the control group

Feature	Genotype	High n (%)	Normal n (%)	OR _{mv} (95% CI)	P
Waist circumference	<i>PPARγ</i> Pro12Ala				
	Pro12Pro	27 (65.9)	142 (77.2)	1	
	Pro12Ala/Ala12Ala	14 (34.1)	42 (22.8)	3.86 (1.24-12.06)	0.020
Triglycerides	<i>ATIR</i> 1166A>C				
	AA	8 (26.7)	112 (53.1)	1	
	AC or CC	22 (73.3)	99 (46.9)	8.05 (1.85-35.0)	0.005
	total	30 (100.0)	211 (100.0)		

ABBREVIATIONS: OR_{mv} = multivariate odds ratio; 95% CI = 95% confidence interval for odds ratio
Significant values are indicated in bold. P<0.05 was considered statistically significant.

DISCUSSION

Assuming a modest role of single polymorphisms as markers of complex traits and diseases, the study of variant gene combinations could provide complementary information that could be clinically important. The obtained results confirmed the significance of the interacting effects of the *PPAR* γ polymorphisms with some of its target genes.

Although *PPAR* γ Pro12Ala variant in this study showed no influence on MetSy when tested separately, its interaction with *LPL* PvuII (-/+) variants seems to be relevant. Carriers of low activity alleles, *PPAR* γ Ala and *LPL* PvuII (+), had six times greater odds for MetSy. The PPARs have been implicated in the regulation of adipocyte differentiation and lipid and fatty acid metabolism. As both central obesity and serum lipids were found to be related with LPL activity, it is reasonable to assume that those two gene interactions contribute to the development of MetSy. Considering transcriptional regulation, the activity of the LPL promoter has been studied extensively. The 5' regulatory region contains a large number of specific *cis*-acting elements, among others the PPAR response element (Schoonjans et al., 1996). Because the *LPL* promoter was shown to be ~ 30% less efficiently transactivated in the presence of the *PPAR* γ Ala allele in a study in vitro, Schneider group proved that the Pro12Ala substitution may lead to decreased LPL activity in vivo (Schneider et al, 2002).

Our results further pointed to the importance of the interacting role of the *PPAR* γ with target gene *IL6* influencing the glucose level. The role of PPARs has been confirmed in glucose homeostasis but with different results. When analyzed as a single marker for T2DM risk, *PPAR* γ Ala variant was considered protective in different studies (Gouda et al., 2010). However, for association between *PPAR* γ and T2DM, environmental factors as dietary lipids (Scacchi et al., 2007) and even intrauterine condition (de Rooij et al., 2006) were pointed to be relevant in addition to gene variants. In our study *PPAR* γ Ala variant was found protective in patients with *IL6* -174GG genotype, while in the case of *IL6* -174C allele carriers, *PPAR* γ Ala12 variant lost its protective role, and Ala12 carriers had even larger odds for high glucose levels compared to Pro12 variant. In Asian Indian population *PPAR* γ Ala12Ala genotype was associated with obesity and insulin resistance (Bhatt et al., 2012), indicating that ethnicity with different genetic and environmental background can have dominant role. For Croatian population, multivariate prediction revealed that cholesterol and TGs were also associated with high fasting blood glucose, where each unit increase in TGs increased the odds for high glucose 1.3 times. The relationship between insulin resistance and metabolic risk factors is complex and abdominal obesity with dysfunctional adipose tissue might be one of risk predispositions. Variants of *IL6* gene -174G>C are associated with cytokine and metabolic modulation that can lead to impaired glucose and lipid homeostasis and increased metabolic risk. Association of the C allele with obesity has been documented (Berthier et al., 2003). Due to lower energy expenditure and insulin sensitivity, carriers of the CC genotype could be more prone to insulin resistance

and obesity (Fernandez-Real et al., 2000; Kubaszek et al., 2003). Our results are in accordance to such published data. Again, other opposite results can be found in the literature. It has been found that *IL6* -174C allele carriers had significantly lower fasting glucose (Hutch et al., 2009). In our study, association between *IL6* -174CC genotype and higher WC was observed in the control group, which is in accordance with published data and points to the association between C allele and obesity (Berthier et al., 2003). Further, we found interactions of *PPARγ* and *ACE* genotypes significant for BMI, specifically in the group of patients with *ACE* DD genotype where *PPARγ* Ala allele carriers had greater odds for obesity, which is in accordance with the data from previous studies (Passaro et al., 2011) in which have also been reported *ACE* DD / *PPARγ* Ala female carriers having a higher BMI and fat distribution, but not MetSy. Association of the *ACE* DD genotype with BMI can be explained by the observation that *ACE* DD subjects have increased levels of plasma ACE (Rigat et al., 1990; Tiret et al., 1992), and therefore potentially increased level of angiotensin II which is considered a trophic factor in the differentiation of preadipocytes to mature adipocytes (Saint-Marc et al., 2001). Previously published data also suggest that the RAS might be involved in the pathophysiology of obesity (Engeli et al., 2000). The data on the influence of *PPARγ* 12Ala variant on BMI are inconsistent, being associated with decreased BMI (Deeb et al., 1998; Damcott et al., 2004) or increased BMI (Masud and Ye, 2003), and increased risk of obesity (Ochoa et al., 2004). In our study, it was observed for the control group in multivariate prediction, that *PPARγ* 12Ala allele carriers had larger odds of having high WC compared to those with *PPARγ* Pro12 allele carriers.

Presented data confirmed that *PPARγ* gene variants, interacting with some of its polymorphic target genes, can modulate pathophysiological processes in the development of MetSy in Croatian population. We have also confirmed that between-study heterogeneity can be at least partly attributed to ethnicity and gene-gene interactions.

Study Limitations

The size of our population might be a limitation. Another limitation might be that the effects of environmental factors and controlling the risk of obesity and MetSy were not evaluated in this study.

Although further studies are needed to confirm our results, the interplay between genes seems to be significant.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgements

This study was supported by grant of Croatian Ministry of Science, Education and Sports as part of Project No. 108-1080134-0136 'Functional Genomics and Proteomics of Risk Factors for Atherosclerosis'.

References

- Angelakopoulou A, Shah T, Sofat R, et al. (2012) Comparative analysis of genome-wide association studies signals for lipids, diabetes, and coronary heart disease: Cardiovascular Biomarker Genetics Collaboration. *Eur Heart J* 33(3):393-407.
- Azhar S. (2010) Peroxisome proliferator-activated receptors, metabolic syndrome and cardiovascular disease. *Future Cardiol* 6(5):657-691.
- Bell-Parikh LC, Ide T, Lawson JA, et al. (2003) Biosynthesis of 15-deoxydelta12,14-PGJ2 and the ligation of PPARgamma. *J Clin Invest* 112(6):945-955.
- Berthier MT, Paradis AM, Tchernof A, et al. (2003) The interleukin 6 -174G/C polymorphism is associated with indices of obesity in men. *J Hum Genet* 48(1):14-19.
- Bhatt SP, Misra A, Sharma M, et al. (2012) Ala/Ala genotype of Pro12Ala polymorphism in the peroxisome proliferator-activated receptor- γ 2 gene is associated with obesity and insulin resistance in Asian Indians. *Diabetes Technol Ther* 14(9):828-834.
- Bonnardeaux A, Davies E, Jeunemaitre X, et al. (1994) Angiotensin II type 1 receptor gene polymorphisms in human essential hypertension. *Hypertension* 24(1):63-69.
- Damcott CM, Moffett SP, Feingold E, et al. (2004) Genetic variation in fatty acid-binding protein-4 and peroxisome proliferator-activated receptor gamma interactively influence insulin sensitivity and body composition in males. *Metabolism* 53(3):303-309.
- de Rooij SR, Painter RC, Phillips DI, et al. (2006) The effects of the Pro12Ala polymorphism of the peroxisome proliferator-activated receptor-gamma2 gene on glucose/insulin metabolism interact with prenatal exposure to famine. *Diabetes Care* 29(5):1052-1057.
- Deeb SS, Fajas L, Nemoto M, et al. (1998) A Pro12Ala substitution in PPAR γ 2 associated with decreased receptor activity, lower body mass index and improved insulin sensitivity. *Nat Genet* 20(3):284-287.
- Engeli S, Negrel R, Sharma AM. (2000) Physiology and pathophysiology of the adipose tissue renin angiotensin system. *Hypertension* 35(6):1270-1277.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. (2001) Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 285(19):2486-2497.

- Fernandez-Real JM, Broch M, Vendrell J, et al. (2000) Interleukin-6 gene polymorphism and lipid abnormalities in healthy subjects. *J Clin Endocrinol Metab* 85(3):1334-1339.
- Fernandez-Real JM, Vayreda M, Richart C, et al. (2001) Circulating interleukin 6 levels, blood pressure, and insulin sensitivity in apparently healthy men and women. *J Clin Endocrinol Metab* 86(3):1154-1159.
- Gouda HN, Sagoo GS, Harding AH, et al. (2010) The association between the peroxisome proliferator-activated receptor-gamma2 (PPARG2) Pro12Ala gene variant and type 2 diabetes mellitus: a HuGE review and meta-analysis. *Am J Epidemiol* 171(6):645-655.
- Groop L. (2000) Genetics of the metabolic syndrome. *Br J Nutr* 83(Suppl 1):S39-48.
- He W. (2009) PPARGgamma2 Polymorphism and Human Health. *PPAR Res* 2009:849538.
- Hilgers KF, Langenfeld MRW, Schlaich M, et al. (1999) 1166 A/C polymorphism of the angiotensin II type 1 receptor gene and the response to short-term infusion of angiotensin II. *Circulation* 100(13):1394-1399.
- Huang HY, Thuita L, Strickland P, et al. (2007) Frequencies of single nucleotide polymorphisms in genes regulating inflammatory responses in a community-based population. *BMC Genet* 8:7.
- Huang X, Zhao J, Zhao T. (2011) Effects of peroxisome proliferator activated receptor-gamma 2 gene Pro12Ala polymorphism on fasting blood lipids: a meta-analysis. *Atherosclerosis* 215(1):136-144.
- Hutch C, Illig T, Herder C, et al. (2009) Joint analysis of individual participants' data from 17 studies on the association of the IL-6 variant -174G>C with circulating glucose levels, interleukin-6 levels, and body mass index. *Ann Med* 41(2):128-138.
- Ide T, Egan K, Bell-Parikh LC, et al. (2003) Activation of nuclear receptors by prostaglandins. *Thromb Res* 110(5-6):311-315.
- Jaccard J. (2001) Interaction effects in logistic regression. *Sage University Papers Series on Quantitative Applications in the Social Sciences*. Thousand Oaks, CA: SAGE, pp 07-135.
- Jamie WE, Edwards RK, Ferguson RJ, et al. (2005) The interleukin-6--174 single nucleotide polymorphism: cervical protein production and the risk of preterm delivery. *Am J Obstet Gynecol* 192(4):1023-1027.
- Kleinbaum DG and Klein M. (2002) *Logistic regression: A self learning text* (2nd ed.). New York: Springer.
- Kraja AT, Borecki IB, North K, et al. (2006) Longitudinal and age trends of metabolic syndrome and its risk factors: the Family Heart Study. *Nutr Metab (Lond)* 3:41.
- Kubaszek A, Pihlajamaki J, Punnonen K, et al. (2003) The C-174G promoter polymorphism of the IL-6 gene affects energy expenditure and insulin sensitivity. *Diabetes* 52(2):558-561.
- Curti ML, Jacob P, Borges MC, et al. (2011) Studies of gene variants related to inflammation, oxidative stress, dyslipidemia, and obesity: implications for a nutrigenetic approach. *J Obes* 2011:497401.

- Lin HF, Boden-Albala B, Juo SH, et al. (2005) Heritabilities of the metabolic syndrome and its components in the Northern Manhattan Family Study. *Diabetologia* 48(10):2006-2012.
- Lusis AJ, Attie AD, Reue K. (2008) Metabolic syndrome: from epidemiology to systems biology. *Nat Rev Genet* 9(11):819-830.
- Masud S, Ye S, SAS Group. (2003) Effect of the peroxisome proliferator activated receptor-gamma gene Pro12Ala variant on body mass index: a meta-analysis. *J Med Genet* 40(10):773-780.
- Miller SA, Dykes DD, Polesky HF. (1988) A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 16(3):1215.
- Möllersten A, Stegmayr B, Wiklund PG. (2008) Genetic polymorphisms in the renin-angiotensin system confer increased risk of stroke independently of blood pressure: a nested case-control study. *J Hypertens* 26(7):1367-1372.
- Monda KL, North KE, Hunt SC, et al. (2010) The genetics of obesity and the metabolic syndrome. *Endocr Metab Immune Disord Drug Targets* 10(2):86-108.
- Ochoa MC, Marti A, Azcona C, et al. (2004) Gene-gene interaction between PPAR gamma 2 and ADR beta 3 increases obesity risk in children and adolescents. *Int J Obes Relat Metab Disord* 28(Suppl 3):S37-S41.
- Oh EY, Min KM, Chung JH, et al. (2000) Significance of Pro12Ala mutation in peroxisome proliferator-activated receptor-gamma2 in Korean diabetic and obese subjects. *J Clin Endocrinol Metab* 85(5):1801-1804.
- Olefsky JM. (2000) Treatment of insulin resistance with peroxisome proliferator-activated receptor gamma agonists. *J Clin Invest* 106:467-472.
- Paracchini V, Pedotti P, Taioli E. (2005) Genetics of leptin and obesity: a HuGE review. *Am J Epidemiol* 162(2):101-114.
- Park YM, Province MA, Gao X, et al. (2009) Longitudinal trends in the association of metabolic syndrome with 550 k single-nucleotide polymorphisms in the Framingham Heart Study. *BMC Proc* 3(Suppl 7):S116.
- Passaro A, Dalla Nora E, Marcello C, et al. (2011) PPAR γ Pro12Ala and ACE ID polymorphisms are associated with BMI and fat distribution, but not metabolic syndrome. *Cardiovasc Diabetol* 10:112.
- Pujia A, Gnasso A, Irace C, et al. (1994) Association between ACE-D/D polymorphism and hypertension in type II diabetic subjects. *J Hum Hypertens* 8(9):687-691.
- Qi L, Zhang C, van Dam RM, et al. (2007) Interleukin-6 genetic variability and adiposity: associations in two prospective cohorts and systematic review in 26,944 individuals. *J Clin Endocrinol Metab* 92(9):3618-3625.
- Rangwala SM and Lazar MA. (2000) Transcriptional control of adipogenesis. *Annu Rev Nutr* 20:535-559.
- Rigat B, Hubert C, Alhenc-Gelas F, et al. (1990) An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *J Clin Invest* 86(4):1343-1346.

- Rigat B, Hubert C, Corvol P, et al. (1992) PCR detection of the insertion/deletion polymorphism of the human angiotensin converting enzyme gene (DCP1) (dipeptidyl carboxypeptidase 1). *Nucleic Acids Res* 20(6):1433.
- Roszer T and Ricote M. (2010) PPARs in the renal regulation of systemic blood pressure. *PPAR Res* 2010:698730.
- Saint-Marc P, Kozak LP, Ailhaud G, et al. (2001) Angiotensin II as a trophic factor of white adipose tissue: stimulation of adipose cell formation. *Endocrinology* 142(1):487-492.
- Sattar N, Perry CG, Petrie JR. (2003) Type 2 diabetes as an inflammatory disorder. *Br J Diabetes Vasc Dis* 3(1):36-41.
- Scacchi R, Pinto A, Rickards O, et al. (2007) An analysis of peroxisome proliferator-activated receptor gamma (PPAR-gamma 2) Pro12Ala polymorphism distribution and prevalence of type 2 diabetes mellitus (T2DM) in world populations in relation to dietary habits. *Nutr Metab Cardiovasc Dis* 17(9):632-41.
- Schneider J, Kreuzer J, Hamann A, et al. (2002) The proline12alanine substitution in the peroxisome proliferator-activated receptor- γ 2 gene is associated with lower lipoprotein lipase activity in vivo. *Diabetes* 51(3):867-870.
- Schoonjans K, Peinado-Onsurbe J, Lefebvre AM, et al. (1996) PPARalpha and PPARgamma activators direct a distinct tissue-specific transcriptional response via a PPRE in the lipoprotein lipase gene. *EMBO J* 15(19):5336-5348.
- Semenkovich CF, Wims M, Noe L, et al. (1989) Insulin regulation of lipoprotein lipase activity in 3T3-L1 adipocytes is mediated at posttranscriptional and posttranslational levels. *J Biol Chem* 264(15):9030-9038.
- Sharma AM and Staels B. (2007) Review: Peroxisome proliferator-activated receptor gamma and adipose tissue--understanding obesity-related changes in regulation of lipid and glucose metabolism. *J Clin Endocrinol Metab* 92(2):386-395.
- Stephens JW, Hurel SJ, Cooper JA, et al. (2004) A common functional variant in the interleukin-6 gene is associated with increased body mass index in subjects with type 2 diabetes mellitus. *Mol Genet Metab* 82(2):180-186.
- Stephens JW, Hurel SJ, Lowe GDO, et al. (2007) Association between plasma IL-6, the IL6 -174G>C gene variant and the metabolic syndrome in type 2 diabetes mellitus. *Mol Genet Metab* 90(4):422-428.
- Tiret L, Blanc H, Ruidavets JB, et al. (1998) Gene polymorphisms of the renin-angiotensin system in relation to hypertension and parental history of myocardial infarction and stroke: the PEGASE study. *Projet d'Etude des Genes de l'Hypertension Arterielle Severe a moderee Essentielle*. *J Hypertens* 16(1):37-44.
- Tiret L, Rigat B, Visvikis S, et al. (1992) Evidence, from combined segregation and linkage analysis, that a variant of the angiotensin I-converting enzyme (ACE) gene controls plasma ACE levels. *Am J Hum Genet* 51(1):197-205.
- Xu E, Li W, Zhan L, et al. (2008) Polymorphisms of the lipoprotein lipase gene are associated with atherosclerotic cerebral infarction in the Chinese. *Neuroscience* 155(2):403-408.