

Electroencephalographic response to different odors in healthy individuals: a promising tool for objective assessment of olfactory disorders

Krbot Skorić, Magdalena; Adamec, Ivan; Jerbić, Ana Branka; Gabelić, Tereza; Hajnšek, Sanja; Habek, Mario

Source / Izvornik: **Clinical EEG and Neuroscience, 2015, 46, 370 - 376**

Journal article, Accepted version

Rad u časopisu, Završna verzija rukopisa prihvaćena za objavljivanje (postprint)

<https://doi.org/10.1177/1550059414545649>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:120320>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-09-08**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine](#)
[Digital Repository](#)





Središnja medicinska knjižnica

Krbot Skorić M., Adamec I., Jerbić A. B., Gabelić T., Hajnšek S., Habek M. (2015) *Electroencephalographic response to different odors in healthy individuals: a promising tool for objective assessment of olfactory disorders*. *Clinical EEG and Neuroscience*, 46 (4). pp. 370-6. ISSN 1550-0594

<http://eeg.sagepub.com>

<http://dx.doi.org/10.1177/1550059414545649>

<http://medlib.mef.hr/2398>

University of Zagreb Medical School Repository

<http://medlib.mef.hr/>

EEG response to different odors in healthy individuals: a promising tool for objective assessment of olfactory disorders

Magdalena Krbot Skorić¹, Ivan Adamec¹, Ana Branka Jerbić², Tereza Gabelić¹, Sanja Hajnšek^{1,3}, Mario Habek^{1,3}

¹University Hospital Center Zagreb, Department of Neurology, Referral Center for Demyelinating Diseases of the Central Nervous System, Zagreb, Croatia

²University of Zagreb

³University of Zagreb, School of Medicine

Corresponding author:

Mario Habek, MD, PhD

University Department of Neurology, University Hospital Center Zagreb

Kišpatićeva 12

HR-10000 Zagreb

Croatia

Phone/Fax: +38512376033; e-mail: mhabek@mef.hr

Word count: 2825

Number of references: 18

Number of tables: 4

Number of figures: 3

Authors' contributions:

Study concept and design: Krbot Skorić, Habek. Acquisition of data: Krbot Skorić, Adamec, Jerbić, Gabelić, Hajnšek, Habek. Analysis and interpretation of data: Krbot Skorić, Adamec, Jerbić, Gabelić, Hajnšek, Habek. Drafting of the manuscript: Krbot Skorić, Habek. Critical revision of the manuscript for important intellectual content: Krbot Skorić, Adamec, Jerbić, Gabelić, Hajnšek, Habek. Administrative, technical, and material support: Krbot Skorić, Adamec, Jerbić, Gabelić, Hajnšek, Habek.

Conflict of interest statement:

Mrs. Krbot Skorić reports no disclosures.

Dr. Adamec reports no disclosures.

Mrs. Jerbić reports no disclosures.

Dr. Gabelić reports no disclosures.

Dr. Hajnšek reports no disclosures.

Dr. Habek reports no disclosures.

Source of funding: None other than the author's own institution.

Abstract

The aim of the present study was to examine human CNS response to three different types of odor: lemon, vanilla and peppermint. Electrophysiological activity was recorded in the baseline state and for three odors, lemon, peppermint and vanilla in 16 healthy participants. For further analysis, electrodes were separated into groups according to the spatial position on the head. Fast Fourier Transformation analysis was performed on every set and mean value of activity in theta interval was exported. As the theta region showed statistically significant results, further analysis was based only on the theta frequency band. On electrodes FP1, F3, Fz, F4, F8, T7, C3, Cz, C4, T8, TP9, CP5, CP1, CP2, CP6, P7, P3, Pz, P4, P8, PO9 and PO10 there was statistically significant difference in the electrical activity of the brain between four conditions. For peppermint and lemon there was statistically significant difference in activity between different regions ((F (1.576, 23.637) = 16.030, p = 0.000) and (F (1.362, 20.425) = 4.54, p = 0.035), respectively), where the activity in the central area was significantly reduced compared to the activity in the other four areas and in the left and right anterior and left posterior area, respectively. There was no statistically significant difference for vanilla in activity between specific areas (F(1.217, 18.257) = 1.155, p = 0.309). The results obtained in this study indicate that olfactory stimuli can affect the frequency characteristics of the electrical activity of the brain.

Key words: odor, EEG response, theta band

Introduction

Although many studies have shown that olfactory function is affected in many neurological diseases, olfactory disorders are often neglected in neurological examination and even rarer rated in the clinical setting. Nevertheless, their evaluation can be useful for diagnosis of many neurological disorders like Parkinson's disease, Alzheimer's disease, multiple sclerosis, Huntington's disease and motor neuron disease. (1) In all of these disorders, basic ability of sense of odors can be reduced, as well as the ability to distinguish different odors or determining thresholds / odor detection. The malfunctioning of the olfactory system can also be an indicator of disease progression and correlate with cognitive deterioration in patients with dementia. (2) According to the results of the meta-analysis, the ability to sense and distinguishing odors could be used as one method of detecting patients with Alzheimer's disease, and the method that examines the threshold stimulus for certain odors could be used in the detection of patients with Parkinson's disease. (3) In patients with mild cognitive impairment (MCI) it was observed that weakened olfactory function, and unconsciousness of the deficit, could also be one of the indicators of progression of mild cognitive impairment to Alzheimer's disease.(4) It is important to mention that the attenuation of function of the olfactory system is also observed during natural aging. (5)

Methods used in the examination of the olfactory system are psychophysical tests, olfactory evoked potentials and functional magnetic resonance imaging (fMRI). The mostly used psychophysical tests are: the University of Pennsylvania Smell Identification Test (UPSIT or SIT) , Connecticut Chemosensory Clinical Research Center Test (CCCRC test) and Sniffin' Sticks test.(1) UPSIT test consists of 40 different scents and is one of the most commonly used tests to determine the functional state of the olfactory system. CCCRC test consists of ten different scents, of which seven are used to test the olfactory system, while three fragrances are used for testing the trigeminal system. Sniffin' Sticks test has three levels and is

used to test thresholds for different odors, the ability of sense of odors and ability to distinguish between different odors. Lack of psychophysical tests is their dependence on the active participation of patients and their subjective impact. Olfactory evoked potentials are responses of the olfactory system related to specific olfactory stimulus. Their advantage is the independence of the cooperation of participants and the ability to use in situations in which the active cooperation is very difficult to implement. The disadvantage of this method is the need for very precise synchronization of olfactory stimulus with electrical brain activity that has been recorded in a given moment and a need for a complete ventilation of the chamber after each presentation of stimuli, so that each stimulus presentation has the same initial, i.e. neutral conditions as the previous presentation. fMRI on the other hand gives an insight into the functional neuroanatomy of the olfactory system.(1,6-8)

On the other hand several studies have shown that electrical brain activity (electroencephalography – EEG) in humans can be influenced by various odors. (9-12) Despite these promising results, the widespread use and adoption of the method is still missing.

Recording of the electrical brain activity (EEG) is a noninvasive method that could serve as an objective method for evaluating the function of the olfactory system. The method is available in almost every clinical facility, and it is possible to achieve results in a relative short period of time and the method does not require active cooperation of participants.

The aim of the present study was to examine human CNS response to three different types of odor: lemon, vanilla and peppermint.

Materials and methods

Sixteen healthy participants participated in this study, 7 females and 9 males (mean age 29+/- 3). They had no known neurological disorders, and according to subjective claims they all have functional sense of smell. Participants were informed about all aspects of the study and

they all signed informed consent approved by Ethical Committee of University Hospital Center Zagreb.

Ability to distinguish odors was determined using a simple test in which participants had to recognize different odors that will be used later in the study. Each odor was presented twice in arbitrary order. All participants recognized presented odors with complete accuracy.

During the experiment participants were placed in sound- and light-insulated chamber with its own ventilation system. They sat in a comfortable armchair and were instructed to relax and to minimize blinking in order to reduce internal artifacts. During the experiment participants had to keep their eyes open in order to avoid unwanted alpha activity, which has very large amplitude and occurs in a state in which a person is at rest with eyes closed. Due to the very small amplitude of the signal that is important for this study, the presence of alpha rhythm would mean an unwanted noise. The activity of the alpha rhythm has frequency of 10 Hz, and is better to avoid this activity during the recording of the experiment, so after, during signal analysis, no important parts of signal would be removed with filtering. Participants were asked to minimize their cognitive activity in order to put their brain in “idling” state.

Electrophysiological activity was recorded with EEG cap with 31 active electrodes [BrainProducts GmbH, Germany] positioned according to International 10-20 System. Active electrodes based on quality Ag/AgCl sensors have integrated circuits for noise reduction. For further analysis, electrodes were separated into groups according to the spatial position on the head: left anterior - LA (FP1, F3, F7, Fc1, Fc5, T7, C3), right anterior - RA (FP2, F4, F8, FC2, FC6, T8 C4), central - C (Fz, Cz, Pz), left posterior - LP (TP9, CP1, CP5, P3, P7, O1, PO9), right posterior - RP (TP10, CP2, CP6, P4, P8, O2, PO10) as presented in Figure 1.

Areas under each electrode were cleaned with abrasive paste in order to reduce impedance and conductive paste was applied to each area in order to achieve adequate conductivity for recording very small signal (order of magnitude $\sim\mu\text{V}$). Vertical oculogram was recorded

below the right eye to detect vertical ocular movements for more precise treatment of ocular artifacts.

Recordings were performed with BrainAmp amplifier and recording software Brain Vision Recorder [BrainProducts GmbH, Germany]. The recording scheme is presented on Figure 2. Recorded signals were filtered with band pass filter from 0.1 Hz to 250 Hz. Sampling frequency was 1000 Hz. Data analysis was performed with software Brain Vision Analyzer [BrainProducts GmbH, Germany].

Each session consisted of five sets, two baseline sets with no odor, and three sets with three different odors (lemon, vanilla, peppermint). During recording, the subject was instructed to breathe evenly through the nose. In the first two sets of recordings no odors were presented to participants; goal of the first set was the preparation for the experiment, so participants could relax and get used to the conditions in the chamber, while in the second set the basic, spontaneous electrical activity of the brain was recorded, and that activity was later used as the basis for comparison with the electrical activity of the brain in experimental conditions.

Each participant was his own control, comparing the conditions of spontaneous brain activity with conditions in which the participant was influenced with presented olfactory stimulus.

After the initial phase, three different odors in random order were presented to each participant. Odors used in this study were essential oils, recognized by each participant in the initial part of the experiment. Essential oils used in this study were lemon, vanilla and peppermint and were selected according to previously conducted studies. (10,11) Odors were prepared immediately before each set of recordings and presented on clean, unused perfume test strips fixed 10 cm in front of the participant's nose.

Each set, the baseline set and the odor set, lasted two minutes. Between each set there was two minutes break with ventilation system on in order to reduce influence of previously presented odor to odor that will be presented in the next set.

For each set of data (basic activity, lemon, vanilla, peppermint) a fast Fourier transform (FFT) was performed in order to achieve the separation of the frequency components that are of interest for further analysis. The frequency resolution used in this analysis was 0.002 Hz. The result of the frequency analysis is expressed as a power, i.e. squared amplitude value of activity (μV^2). Hanning window is used in analysis, in order to reduce the boundary conditions. Mean activity in a particular frequency band was used for further statistical analysis.

Statistical analysis was performed using software IBM SPSS 20.0 (Chicago, IL). The collected data had a normal distribution and was mutually dependent, because there are several successive measurements at the same participant, and suitable statistical analysis was repeated measures ANOVA, with Bonferroni correction for post hoc analysis. P values less than 0.05 were considered statistically significant.

Results

All EEG results were interpreted by regarding each subject as his/her own control (comparison of conditions in which there is an olfactory stimulus with the condition in which there is only a basic activity). The initial analysis was conducted in all frequency bands (alpha, beta, delta, theta) (Table 1), but only the theta region showed statistically significant results and therefore further analysis was based only on the theta frequency band.

In the first step of the analysis, we examined whether on each electrode there is a difference between the four conditions (peppermint, lemon, vanilla, basic activity).

The analysis was conducted by repeated measures ANOVA. When the rule of sphericity was violated, the Greenhouse-Geisser correction was conducted. The analysis showed that on electrodes FP1, F3, Fz, F4, F8, T7, C3, Cz, C4, T8, TP9, CP5, CP1, CP2, CP6, P7, P3, Pz, P4,

P8, PO9 and PO10 there was statistically significant difference in the electrical activity of the brain between four conditions. Results of the analysis are presented in Table 2.

After the initial statistical analysis, post hoc analysis with Bonferroni correction was performed in order to determine exactly between which pairs of conditions there is a statistically significant difference. The results of this analysis are shown in Table 3. In all cases the mean intensity of the electrical activity in the theta band excited with presented olfactory stimulus (lemon, peppermint, and vanilla) was reduced relative to the mean intensity value of primary, spontaneous brain activity. The table shows that the electrical activity of the brain induced with the odor of lemon showed a statistically significant reduction in relation to the baseline spontaneous brain activity on the greatest number of electrodes (15 of 31, 48%), followed by brain activity induced with the odor of peppermint (10 of 31, 32%) and the fewest number of electrodes was activated with the odor of vanilla (3 of 31, 10%).

In the second part of analysis, the values obtained at specific electrodes are grouped according to the spatial position in one of the five areas (LA, DA, C, LP, DP) and mean activity value was averaged for each region for each of the three scents (peppermint, vanilla, lemon). For each of the odors, repeated measures ANOVA with Greenhouse-Geisser correction was performed in order to examine whether there was a difference in activity between different areas.

To determine between which areas there was a statistically significant difference, the post hoc analysis with Bonferroni correction was performed, and the results are presented in Table 4.

For peppermint there was statistically significant difference in activity between different regions ($F(1.576, 23.637) = 16.030, p = 0.000$), where the activity in the central area was significantly reduced compared to the activity in the other four areas.

There was no statistically significant difference for vanilla in activity between specific areas ($F(1.217, 18.257) = 1.155, p = 0.309$).

For lemon there was statistically significant difference in activity between different regions ($F(1.362, 20.425) = 4.54, p = 0.035$), where the activity in the central area was significantly reduced compared to the activity in the left and right anterior and left posterior area.

Figure 3 presents the spatial distribution of mean values of the electrical activity of the brain for peppermint, vanilla and lemon.

Discussion

This study has shown that stimulation with selected odors is associated with a significant reduction in theta band. Furthermore we have shown that lemon and peppermint beside reduction of theta band have the most pronounced effect of the reduction of theta band in the central region (electrodes).

However, it has to be emphasized that previously published studies related to the impact of olfactory stimulus to changes in the intensity of the brain electrical activity were performed with different methodologies. According to some authors, there is a statistically significant difference in the alpha frequency range between conditions in which there is olfactory stimulus and the baseline condition with no stimulus. (13,14) Other studies report a statistically significant difference in the theta frequency range. (9,10,11) The methodology used in the present study was based on the methodology described in the paper by Neil Martin, (11) and the results obtained are consistent with the results obtained in that study, indicating the reproducibility of the results when using a similar methodology.

The analysis showed that on electrodes FP1, F3, Fz, F4, F8, T7, C3, Cz, C4, T8, TP9, CP5, CP1, CP2, CP6, P7, P3, Pz, P4, P8, PO9 and PO10 there was statistically significant difference in the theta electrical activity of the brain induced with odors of peppermint, vanilla and lemon and between the baseline activity. Between activities induced by different olfactory stimulus, there was no statistically significant difference. For all statistically significant

results, the intensity of activity in the theta frequency range induced by certain odor was reduced in comparison with the intensity of activity in the theta frequency range in situations where there was no olfactory stimulus. These results indicate that there is a difference in the electrical activity of the brain evoked by olfactory stimulation related to the baseline electrical brain activity with no presentation of odors.

Lemon and peppermint elicited reduction of intensity of theta activity on larger number of electrodes than vanilla. In initial tests, where participants had to recognize presented odors, most participants characterized scent of vanilla as very mild odor, although it was dosed in the same amount as the other two scents. It is possible that in the case of vanilla arousal of the olfactory system was not the same as the arousal for lemon and peppermint, which is why the influence of the vanilla odor to intensity of brain activity is reduced.

Analysis of the activities of groups of electrodes showed that peppermint and lemon elicited statistically significant difference in the intensity of activity between different regions, while with the odor of vanilla, this difference was not statistically proven. If we compare this with previously presented results where the vanilla had the slightest impact on the intensity of activity of theta band and the fact that the subjects who participated in the survey characterized vanilla as very mild odor, then we can assume that the odor of vanilla did not have the same effect on olfactory system as the other two odors. All odors that were used in this study were rated by participants as pleasant, which is extremely important because the degree of pleasantness is an important factor that has influence on the intensity of activity, and it is also possible to perform a classification of EEG signals to determine whether the odor is pleasant or unpleasant smell for subjects. (15)

For odors of lemon and peppermint, induced activity in the central area was significantly reduced in comparison to the induced activity in the remaining areas. In a study by Cherninskii et al, the changes of activity in given frequency bands were mostly pronounced in

the left temporal and parietal regions. (16) Also, there was a clear difference between the excitation of the olfactory and trigeminal systems, where the response to trigeminal stimulation (CO₂) occurs in areas of the cortex associated with locomotion of sniffing / smelling, while the response to olfactory stimuli (H₂S) occurs to a greater extent in the primary olfactory cortex. (17) Also, activation of the olfactory area is manifested as increased intensity of the electrode Pz (parietal area), while the activation of the trigeminal nerve is manifested as increased intensity of the electrode Cz (central area). (18)

The amygdala is also a part of olfactory system and its role in emotions and memory gives a complexity to functioning of olfactory system and may be one possible explanation for the different frequency and spatial distribution of research results related to the olfactory system. (15)

Conclusion

The results obtained in this study indicate that olfactory stimuli can affect the frequency characteristics of the electrical activity of the brain. The method itself is non-invasive and easily enforceable and shows promising contribution in testing the functionality of the olfactory system in humans. It is necessary to conduct research on a larger number of participants and in conditions were not only healthy participants are included, but also participants with various neurological diseases.

References:

1. Barresi M, Ciurleo R, Giacoppo S, Foti Cuzzola V, Celi D, Bramanti P, Marino S. Evaluation of olfactory dysfunction in neurodegenerative diseases. *J Neurol Sci* 2012;323:16-24.
2. Velayudhan L, Pritchard M, Powell JF, Proitsi P, Lovestone S. Smell identification function as a severity and progression marker in Alzheimer's disease. *Int Psychogeriatr* 2013;25:1157-66.
3. Rahayel S, Frasnelli J, Joubert S. The effect of Alzheimer's disease and Parkinson's disease on olfaction: a meta-analysis. *Behav Brain Res.* 2012;231:60-74.
4. Devanand DP, Michaels-Marston KS, Liu X, Pelton GH, Padilla M, Marder K, Bell K, Stern Y, Mayeux R. Olfactory deficits in patients with mild cognitive impairment predict Alzheimer's disease at follow-up. *Am J Psychiatry* 2000;157:1399-405.
5. Kovács T. Mechanisms of olfactory dysfunction in aging and neurodegenerative disorders. *Ageing Res Rev* 2004;3:215–232.
6. Poellinger A, Thomas R, Lio P, Lee A, Makris N, Rosen BR, Kwong KK. Activation and habituation in olfaction—an fMRI study. *Neuroimage* 2001;13:547–60.
7. Murata Y, Okutani F, Nakahira M, Ushida T, Ikemoto T, Yokoe I, Takeda T, Kaba H, Tani T, Ogawa Y. Effects of olfactory stimulation with isovaleric acid on brain activation in informed and naïve conditions: a functional MRI study. *Auris Nasus Larynx* 2007;34:465-9.
8. Cerf-Ducastel B, Murphy C. FMRI brain activation in response to odors is reduced in primary olfactory areas of elderly subjects. *Brain Res* 2003;986:39-53.
9. Lorig TS, Schwartz GE. Brain and odor: I. Alteration of human EEG by odor administration. *Psychobiology* 1988;16:281-284.
10. Klemm WR, Lutes SD, Hendrix DV, Warrenberg S. Topographical EEG maps of human responses to odors. *Chem Senses* 1992;17, 347-361.

11. Martin GN. Human electroencephalographic (EEG) response to olfactory stimulation: two experiments using the aroma of food. *Int J Psychophysiol* 1998;30:287-302.
12. Brauchli P, Rüegg PB, Etzweiler F, Zeier H. Electrocortical and autonomic alteration by administration of a pleasant and an unpleasant odor. *Chem Senses* 1995;20:505-15.
13. Lorig TS, Huffman E, DeMartino A, DeMarco J. The effects of low concentration odors on EEG activity and behavior. *J Psychophysiol* 1991;5:69-77.
14. Van Toller S, Behan J, Howells P, Kendal-Reed M, Richardson A. An analysis of spontaneous human cortical EEG activity to odors. *Chem Senses* 1993;18:1-16.
15. Yazdani A, Kroupi E, Vesin J, Ebrahimi T. Electroencephalogram alterations during perception of pleasant and unpleasant odors. *Quality of Multimedia Experience (QoMEX), 2012 Fourth International Workshop on* 2012:271-277; doi: 10.1109/QoMEX.2012.6263860
16. Cherninskii AA, Zima IG, Makarchouk NYe, Piskorskaya NG, Kryzhanovskii SA. Modifications of EEG Related to Directed Perception and Analysis of Olfactory Information in Humans. *Neurophysiol* 2009;41:63-70.
17. Iannilli E, Wiens S, Arshamian A, Seo HS. A spatiotemporal comparison between olfactory and trigeminal event-related potentials. *Neuroimage* 2013;77:254-61.
18. Livermore A, Hummel T, Kobal G. Chemosensory event-related potentials in the investigation of interactions between the olfactory and the somatosensory (trigeminal) systems. *Electroencephalogr Clin Neurophysiol* 1992;83:201-10.

Tables

Table 1. A) Results of repeated measures ANOVA that show on which electrodes there was a statistically significant difference between alpha activity in condition with olfactory stimulation and baseline activity

		df	F	Sig.
Fp1	Greenhouse-Geisser	1.609	0.097	0.868
Fp2	Greenhouse-Geisser	1.250	0.528	0.516
F7	Greenhouse-Geisser	1.974	0.098	0.904
F3	Greenhouse-Geisser	2.392	0.695	0.530
Fz	Greenhouse-Geisser	1.916	0.580	0.559
F4	Greenhouse-Geisser	1.897	1.067	0.354
F8	Greenhouse-Geisser	2.457	1.187	0.323
FC5	Greenhouse-Geisser	2.079	0.519	0.607
FC1	Greenhouse-Geisser	2.249	0.068	0.949
FC2	Greenhouse-Geisser	2.157	0.785	0.473
FC6	Greenhouse-Geisser	2.118	1.280	0.293
T7	Greenhouse-Geisser	2.295	0.980	0.395
C3	Greenhouse-Geisser	2.633	1.062	0.370
Cz	Greenhouse-Geisser	1.723	0.728	0.473
C4	Greenhouse-Geisser	2.240	1.123	0.342
T8	Greenhouse-Geisser	2.087	0.215	0.817
TP9	Greenhouse-Geisser	1.691	0.244	0.748
CP5	Greenhouse-Geisser	1.875	0.617	0.536
CP1	Greenhouse-Geisser	1.740	0.546	0.562
CP2	Greenhouse-Geisser	1.727	0.668	0.500
CP6	Greenhouse-Geisser	1.599	0.332	0.673
TP10	Greenhouse-Geisser	2.157	0.400	0.689
P7	Greenhouse-Geisser	1.380	0.311	0.655
P3	Greenhouse-Geisser	1.374	0.513	0.539
Pz	Greenhouse-Geisser	1.629	0.596	0.526
P4	Greenhouse-Geisser	1.475	0.360	0.638
P8	Greenhouse-Geisser	1.405	0.144	0.791
PO9	Greenhouse-Geisser	1.502	0.506	0.558
O1	Greenhouse-Geisser	1.369	0.394	0.602
PO10	Greenhouse-Geisser	1.769	0.358	0.676
O2	Greenhouse-Geisser	1.277	0.709	0.444

B) Results of repeated measures ANOVA that show on which electrodes there was a statistically significant difference between beta activity in condition with olfactory stimulation and baseline activity.

		df	F	Sig.
Fp1	Greenhouse-Geisser	2.208	1.999	0.148
Fp2	Greenhouse-Geisser	1.821	0.928	0.399
F7	Greenhouse-Geisser	1.737	0.864	0.419
F3	Greenhouse-Geisser	2.281	0.375	0.717
Fz	Greenhouse-Geisser	2.243	0.557	0.598
F4	Greenhouse-Geisser	1.893	0.809	0.449
F8	Greenhouse-Geisser	2.346	0.442	0.678
FC5	Greenhouse-Geisser	1.879	0.897	0.413
FC1	Greenhouse-Geisser	2.069	0.503	0.616
FC2	Greenhouse-Geisser	1.304	0.384	0.598
FC6	Greenhouse-Geisser	1.572	1.743	0.200
T7	Greenhouse-Geisser	2.047	0.579	0.570
C3	Greenhouse-Geisser	2.188	1.757	0.186
Cz	Greenhouse-Geisser	2.184	0.347	0.728
C4	Greenhouse-Geisser	2.523	1.452	0.246
T8	Greenhouse-Geisser	2.286	0.956	0.405
TP9	Greenhouse-Geisser	2.123	0.542	0.597
CP5	Greenhouse-Geisser	2.066	0.859	0.437
CP1	Greenhouse-Geisser	2.315	1.002	0.387
CP2	Greenhouse-Geisser	1.711	0.715	0.478
CP6	Greenhouse-Geisser	1.947	0.612	0.545
TP10	Greenhouse-Geisser	1.986	2.260	0.122
P7	Greenhouse-Geisser	1.615	0.265	0.723
P3	Greenhouse-Geisser	2.166	0.473	0.643
Pz	Greenhouse-Geisser	2.332	0.868	0.443
P4	Greenhouse-Geisser	2.208	0.234	0.814
P8	Greenhouse-Geisser	1.696	0.170	0.810
PO9	Greenhouse-Geisser	1.908	0.337	0.706
O1	Greenhouse-Geisser	1.076	0.973	0.345
PO10	Greenhouse-Geisser	2.544	2.623	0.073
O2	Greenhouse-Geisser	1.799	1.689	0.205

C) Results of repeated measures ANOVA that show on which electrodes there was a statistically significant difference between delta activity in condition with olfactory stimulation and baseline activity.

		df	F	Sig.
Fp1	Greenhouse-Geisser	1.203	0.723	0.431
Fp2	Greenhouse-Geisser	1.640	1.082	0.343
F7	Greenhouse-Geisser	1.360	0.632	0.483
F3	Greenhouse-Geisser	2.284	0.430	0.680
Fz	Greenhouse-Geisser	2.094	0.872	0.433
F4	Greenhouse-Geisser	1.067	2.200	0.157
F8	Greenhouse-Geisser	2.205	0.935	0.411
FC5	Greenhouse-Geisser	2.125	0.849	0.444
FC1	Greenhouse-Geisser	1.370	0.336	0.638
FC2	Greenhouse-Geisser	1.797	0.589	0.544
FC6	Greenhouse-Geisser	2.280	0.265	0.797
T7	Greenhouse-Geisser	1.021	0.966	0.343
C3	Greenhouse-Geisser	2.460	0.178	0.877
Cz	Greenhouse-Geisser	1.626	0.368	0.653
C4	Greenhouse-Geisser	2.699	0.630	0.584
T8	Greenhouse-Geisser	1.111	2.655	0.120
TP9	Greenhouse-Geisser	2.321	0.792	0.478
CP5	Greenhouse-Geisser	2.362	0.236	0.825
CP1	Greenhouse-Geisser	1.798	0.910	0.405
CP2	Greenhouse-Geisser	2.377	0.740	0.506
CP6	Greenhouse-Geisser	1.789	0.833	0.434
TP10	Greenhouse-Geisser	1.458	0.277	0.691
P7	Greenhouse-Geisser	2.408	0.182	0.870
P3	Greenhouse-Geisser	2.082	0.719	0.501
Pz	Greenhouse-Geisser	1.559	1.005	0.363
P4	Greenhouse-Geisser	1.591	1.340	0.276
P8	Greenhouse-Geisser	2.063	0.674	0.521
PO9	Greenhouse-Geisser	1.804	0.200	0.798
O1	Greenhouse-Geisser	1.064	1.109	0.313
PO10	Greenhouse-Geisser	1.184	0.859	0.385
O2	Greenhouse-Geisser	1.000	0.995	0.334

Table 2. Results of repeated measures ANOVA that show on which electrodes there was a statistically significant difference between theta activity in condition with olfactory stimulation and baseline activity.

		df	F	Sig.
Fp1	Sphericity Assumed	3	2.842	0.048
Fp2	Greenhouse-Geisser	1.931	3.239	0.055
F7	Sphericity Assumed	3	1.815	0.158
F3	Greenhouse-Geisser	1.893	3.825	0.036
Fz	Sphericity Assumed	3	3.500	0.023
F4	Sphericity Assumed	3	3.563	0.021
F8	Sphericity Assumed	3	2.828	0.049
FC5	Greenhouse-Geisser	1.609	1.526	0.237
FC1	Greenhouse-Geisser	1.237	0.626	0.472
FC2	Sphericity Assumed	3	0.506	0.680
FC6	Sphericity Assumed	3	2.698	0.057
T7	Sphericity Assumed	3	4.462	0.008
C3	Sphericity Assumed	3	4.565	0.007
Cz	Sphericity Assumed	3	4.659	0.006
C4	Greenhouse-Geisser	1.904	3.985	0.031
T8	Sphericity Assumed	3	4.858	0.005
TP9	Sphericity Assumed	3	6.221	0.001
CP5	Sphericity Assumed	3	4.836	0.005
CP1	Sphericity Assumed	3	4.637	0.007
CP2	Sphericity Assumed	3	4.559	0.007
CP6	Sphericity Assumed	3	6.056	0.001
TP10	Greenhouse-Geisser	1.929	1.271	0.295
P7	Sphericity Assumed	3	6.669	0.001
P3	Sphericity Assumed	3	4.990	0.005
Pz	Sphericity Assumed	3	4.460	0.008
P4	Sphericity Assumed	3	5.740	0.002
P8	Sphericity Assumed	3	8.074	0.000
PO9	Sphericity Assumed	3	5.155	0.004
O1	Greenhouse-Geisser	1.000	0.974	0.339
PO10	Sphericity Assumed	3	6.311	0.001
O2	Greenhouse-Geisser	1.042	0.797	0.391

Table 3. Results of repeated measures ANOVA after Bonferroni correction; a statistically significant difference between odors and baseline; comparison of mean values of the activity [μV^2]

	Vanilla vs. baseline	Peppermint vs. baseline	Lemon vs. baseline
F3			0.004450 vs.0.006025
T7		0.008963 vs. 0.010156	0.008788 vs. 0.010156
C4		0.003069 vs. 0.003494	0.003031 vs. 0.003494
T8		0.009706 vs. 0.010763	0.009181 vs. 0.010763
TP9	0.014400 vs. 0.017056	0.014344 vs. 0.017056	0.014019 vs. 0.017056
CP5			0.006769 vs. 0.008075
CP1			0.003763 vs. 0.004675
CP2			0.003706 vs. 0.004631
CP6		0.006838 vs. 0.008025	0.006631 vs. 0.008025
P7		0.011000 vs. 0.013231	0.010831 vs. 0.013231
P3			0.006856 vs. 0.008363
P4		0.006869 vs. 0.008344	0.006650 vs. 0.008344
P8	0.010731 vs. 0.013200	0.010769 vs. 0.013200	0.010356 vs. 0.013200
PO9		0.013938 vs. 0.016406	0.013556 vs. 0.016406
PO10	0.013963 vs. 0.016881	0.014219 vs. 0.016881	0.013825 vs. 0.016881

Table 4. Results of repeated measures ANOVA after Bonferroni correction; a statistically significant difference between different areas; comparison of mean values of the activity [μV^2]

Peppermint		
LA vs. C	0.0111 vs. 0.0032	0.001
RA vs. C	0.0113 vs. 0.0032	0.000
LP vs. C	0.0099 vs. 0.0032	0.000
RP vs. C	0.0096 vs. 0.0032	0.000
Lemon		
LA vs. C	0.0115 vs. 0.0032	0.002
RA vs. C	0.0123 vs. 0.0032	0.005
LP vs. C	0.0112 vs. 0.0032	0.000

R - Right, L - left, C - central, A - anterior, P - posterior

Figures

Figure 1. The distribution of electrodes according to spatial position.




Figure 2. Recording scheme.




Figure 3. Spatial distribution of mean values of the electrical activity of the brain for peppermint (A). vanilla (B) and lemon (C).

