

# What is the acceptable rate of negative appendectomy? Comment on "Prospective evaluation of the added value of imaging within the Dutch National Diagnostic Appendicitis Guideline--do we forget our clinical eye"?

---

Papeš, Dino; Sršen Medančić, Suzana; Antabak, Anko; Sjekavica, Ivica; Luetić, Tomislav

Source / Izvornik: **Digestive Surgery, 2015, 32, 181 - 182**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1159/000380772>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:081200>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-07-13**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine Digital Repository](#)



## What Is the Acceptable Rate of Negative Appendectomy? Comment on Prospective Evaluation of the Added Value of Imaging within the Dutch National Diagnostic Appendicitis Guideline – Do We Forget Our Clinical Eye?

Dino Papeš<sup>a</sup> Suzana Sršen Medančić<sup>a</sup> Anko Antabak<sup>a</sup> Ivica Sjekavica<sup>b</sup> Tomislav Luetić<sup>a</sup>Departments of <sup>a</sup>Surgery and <sup>b</sup>Radiology, Clinical Hospital Center Zagreb, Zagreb, Croatia

We read with interest the article of Schok et al. [1]. In 2014 we did an audit of our results on pediatric appendectomy (0–18 years) in the form of a cross-sectional retrospective study that was conducted using hospital database that reviewed all appendectomies in our department in two two-year periods (1999–2000 and 2012–2013) to see the changes after introduction of ultrasound and CT scan into practice. There were a total of 380 appendectomies performed: 154 as laparoscopic and 226 as open procedures. Regarding the pathological outcome of the operation, negative appendectomy rate (NAR) was 12% (47 patients) and perforation rate (PR) was 18% (69 patients). In children younger than 5 years of age (26 patients) PR was 30% and NAR 8%. Regarding imaging, 83 patients (22%) received preoperative ultrasound examination, and 2 received a CT scan (0.5%). The NAR and PR among children who received preoperative imaging were 8.4% (7/83) and 15.7% (13/83), respectively, which did not differ significantly ( $p > 0.05$ ) from children who did not receive any imaging.

To evaluate our results we searched Pubmed and found that NAR and PR in our series were comparable to other similar reports, including the report from Schok et al. As mentioned by the authors, accord-

ing to Dutch guidelines, such rates may be considered unacceptable. So what is the acceptable NAR? When searching through the literature, one may find reports with considerably low NAR, that is, below 10%. However, there are several factors that apparently decrease NAR.

The three largest series on pediatric appendectomy within the last two years are by Bachur that included 55,227 appendectomies (NAR 3.6%), Oyetunji that included 250,783 appendectomies (NAR 6.7%), and Cheong that included 78,625 children from US and 41,492 children from Canada (NAR 6.3 and 4.3%) [2–4]. Large database analyses report only the discharge letter diagnosis or intraoperative appearance of the appendix without analyzing the histology report. This is present in all three previously mentioned reports. Had we used this criterion, the rate of negative appendectomy in our series would be 3% since of 47 negative appendectomies, 32 (68%) were initially during operation diagnosed as inflamed appendicitis and were later on histological examination found to be non-inflamed. Further, even if histology reports are used, the histological definition of appendicitis also differs. Histological definition of inflamed appendix is not described in many reports that have low negative appendec-

tomy rates [5], and one study has clearly shown that the change in histological criteria for diagnosing inflamed appendix raised the negative appendectomy by 3–6% [6]. Finally, as noted in previous reports, published figures are often without adjustment for age and sex as many hospitals admit proportionally fewer teenage girls or young infants, which are two high-risk groups [7].

For example, one large analysis of 30 pediatric hospitals in the United States found the reported NAR ranging from 0 to 17%, and PR ranging from 20 to 76%. Such differences are most likely the consequence of the aforementioned factors [8].

Regarding the usage of imaging in Europe, this method traditionally plays a modest role in the evaluation of suspected appendicitis when compared to the ultrasound (US) [7]. Although in some studies preoperative US and CT scan did lower the rates of negative appendectomy, both our results and those of Schok et al. led to the conclusion that diagnosis of appendicitis should still mainly be based on history, clinical and laboratory data. Only the development of fast, inexpensive, accurate and non-ionizing radiation imaging modality will influence negative appendectomy rates in a significant way.

## References

- 1 Schok T, Simons PC, Janssen-Heijnen ML, Peters NA, Konsten JL: Prospective evaluation of the added value of imaging within the Dutch national diagnostic appendicitis guideline – do we forget our clinical eye? *Dig Surg* 2014;31:436–443.
- 2 Oyetunji TA, Ong’uti SK, Bolorunduro OB, Cornwell EE 3rd, Nwomeh BC: Pediatric negative appendectomy rate: trend, predictors, and differentials. *J Surg Res* 2012;173:16–20.
- 3 Bachur RG, Hennelly K, Callahan MJ, Chen C, Monuteaux MC: Diagnostic imaging and negative appendectomy rates in children: effects of age and gender. *Pediatrics* 2012;129:877–884.
- 4 Cheong LH, Emil S: Outcomes of pediatric appendicitis: an international comparison of the United States and Canada. *JAMA Surg* 2014;149:50–55.
- 5 Saito JM, Yan Y, Evashwick TW, Warner BW, Tarr PI: Use and accuracy of diagnostic imaging by hospital type in pediatric appendicitis. *Pediatrics* 2013;131:e37–e44.
- 6 Mariadason JG, Wang WN, Wallack MK, Belmonte A, Matari H: Negative appendectomy rate as a quality metric in the management of appendicitis: impact of computed tomography, Alvarado score and the definition of negative appendectomy. *Ann R Coll Surg Engl* 2012;94:395–401.
- 7 Lander A: The role of imaging in children with suspected appendicitis: the UK perspective. *Pediatr Radiol* 2007;37:5–9.
- 8 Newman K, Ponsky T, Kittle K, Dyk L, Throop C, Gieseker K, Sills M, Gilbert J: Appendicitis 2000: variability in practice, outcomes, and resource utilization at thirty pediatric hospitals. *J Pediatr Surg* 2003;38:372–379; discussion 372–379.