

The surgical treatment of patients with colorectal cancer and liver metastases in the setting of the "liver first" approach

Patrlj, Leonardo; Kopljar, Mario; Kliček, Robert; Hrelec Patrlj, Maša; Kolovrat, Marijan; Rakić, Mislav; Đuzel, Antonija

Source / Izvornik: **Hepatobiliary Surgery and Nutrition, 2014, 3, 324 - 329**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.3978/j.issn.2304-3881.2014.09.12>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:105:702507>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-04**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine](#)
[Digital Repository](#)



The surgical treatment of patients with colorectal cancer and liver metastases in the setting of the “liver first” approach

Leonardo Patrlj¹, Mario Kopljar¹, Robert Kliček¹, Masa Hrelec Patrlj², Marijan Kolovrat¹, Mislav Rakić¹, Antonija Đuzel¹

¹Department of Abdominal Surgery, Surgical Clinic, Clinical Hospital Dubrava, Zagreb, Avenija Gojka Šuška 6, 10 000 Zagreb, Croatia; ²Department of Childrens' Surgery, Clinical Hospital for Childrens' Disease, Zagreb, Klaićeva, 10 000 Zagreb, Croatia

Correspondence to: Robert Kliček, MD, PhD. Department of Abdominal Surgery, Surgical Clinic, Clinical Hospital Dubrava, Zagreb, Avenija Gojka Šuška 6, 10 000 Zagreb, Croatia. Email: rklicek@gmail.com.

Abstract: A surgical resection is the only curative method in the therapy of colorectal carcinoma and liver metastases. Along with the development of interventional radiological techniques the indications for surgery widen. The number of metastases and patients age should not present a contraindication for surgical resection. However, there are still some doubts concerns what to resect first in cases of synchronous colorectal carcinoma and liver metastases and how to ensure the proper remnant liver volume in order to avoid postoperative liver failure and achieve the best results. Through this review the surgical therapy of colorectal carcinoma and liver metastases was revised in the setting of “liver-first” approach and the problem of ensuring of remnant liver volume.

Keywords: Colorectal liver metastases; liver resections; remnant volume; “liver-first” approach

Submitted Sep 08, 2014. Accepted for publication Sep 16, 2014.

doi: 10.3978/j.issn.2304-3881.2014.09.12

View this article at: <http://dx.doi.org/10.3978/j.issn.2304-3881.2014.09.12>

Introduction

Treatment of colorectal cancer and liver metastases are an extremely important clinical issue since that there are nearly a million newly diagnosed cases and nearly half of the million reported deaths worldwide (1). In large number of countries the incidence continue to rise (2), although the standardized prevention national programs of early detection have developed and brought to an earlier detection and diagnosed cases in early stage of tumor (3-5). In Asian countries, such as China, Japan, South Korea, and Singapore, a 2-4-fold increase in the incidence of colorectal cancer in the past few decades is experienced (6). In Western World the colorectal cancer is reported as the third the most frequent cancer and the most frequent cancer in population older than 75 years (7).

Approximately 25% of newly diagnosed patients with colorectal cancer will have liver metastases at the time of diagnosis, another 25% will develop liver metastases during the course of the disease and two-thirds of all patients with

liver metastases will die of them (8). The 10-year survival rate for patients with stage I disease is 90%, but for patients with inoperable stage IV disease, it is currently only 5% (9). For patients with liver metastases, the treatment strategy should be directed toward resectability (10).

The multidisciplinary therapeutic approach, consisting of new and more effective chemotherapeutic agents in single or combined therapy, an advanced role of interventional radiology with portal vein embolization (PVE) and tumor ablation and new strategies and techniques for hepatic resections, brought improved resectability rate of metastases to 20-30% of cases and has resulted in 5-year survival of 35-50% for selected cases (11-13). A need has been recognized for a new staging system that acknowledges the improvements in surgical techniques for resectable metastases and the impact of modern chemotherapy on rendering initially unresectable liver metastases from colorectal carcinoma resectable while distinguishing between patients with a chance for cure at presentation and those for whom only palliative treatment is possible (14).

There have been presented the predictive factors for survival and local recurrence (15,16). Traditionally, a staged approach (colorectal first) has been used in the management of patients with synchronous colorectal cancer and liver metastases. This involves the initial extirpation of the primary tumor. Systemic chemotherapy followed the operation, after which liver-directed operation was performed. The last 2 decades have brought an increased understanding of the biology of colorectal liver metastases, resulting in more effective targeted therapies in addition to decreased mortality after liver-directed operations (17,18).

The goal of this review is to focus onto the doubts concerning the operators all around the world in the context of reassuring the proper remnant liver volume and especially what to resect first in the cases of synchronous liver metastases of colorectal carcinoma.

The preoperative imaging and planning the surgical resection

The R0 resection is the ultimate goal of the surgical therapy. However the proper indication is essential in order to achieve adequate result of resection. Resectability depends onto the multiple factors: the number and location of metastases, the remnant liver volume and quality of the liver tissue that is not infiltrated by tumor. All lesions identified at the initial imaging records (CT or MRI) before any therapy is performed have to be accounted during planning the liver resection in order to predict the total risk and the outcome of surgical procedure. It is recognized that chemotherapy can induce toxic injury of liver tissue, primarily steatohepatitis and sinusoidal injury. Non-contrast CT and MRI could be used to assess steatosis (19-21), but steatohepatitis cannot be diagnosed with imaging. Sinusoidal injury can be judged by indirect signs of portal hypertension, particularly spleen size (22), or by using the liver-specific MRI contrast agent gadoxetic acid (23). The essential three points that are ultimate for complete resection are preservation of liver vascularity, the adequate remnant liver volume with reference to body weight and total liver volume, and that the quality of the remnant liver parenchyma is acceptable (24). The ultrasound (US), especially contrast enhanced ultrasound (CEUS) presents a unique imaging method for intraoperative assessment of unrevealed metastases, and the relation between tumor and vascular and biliar structures (25), sometimes even significantly more sensitive than CT and/or MRI preoperative imaging records (26). For the detection of extra hepatic metastases and local recurrence at the site of the initial colorectal surgery, apart

from CT the use of FDG-PET is widespread. A high quality CT can detect the majority of extrahepatic disease, however the FDG-PET may reveal additional signs of disease as high metabolic activity. Although some studies showed a change in management in 10-20% of patients according to record of FDG-PET (27,28), some reports lower percentage and even seem to be more suspicious in its cost-effective role (29), especially in the context of FDG-records following the preoperative chemotherapy which reduces its sensitivity.

The surgical resection—what to resect first in synchronous metastases?

Surgical treatment of colorectal liver metastases remains the only treatment associated with a long survival time in patients with liver metastases from colorectal carcinoma, with a 40% survival at 5 years and almost 25% postoperative survival up to 10 years in specialized centers (30). The very important issue that the liver surgeon has to deal with is to proceed decide what to resect first liver or colon and/or when to undertake simultaneous surgical resections of both. The perfect solution seems to be a single stage colon and liver operation. The advantage of the one stage procedure could be less psychological stress for the patient, lower financial cost and shorter hospitalization time. On the other hand the advantages of the staged procedure are that there is no accumulation of the risks of liver and bowel resections at the same time. Neoadjuvant chemotherapy may be given before liver resection, and an extended hepatectomy or demanding bowel resection could be performed with the full attention of the surgical team focused on the liver or bowel disease, although, the key point for decision-making is the patient's safety (1). According to the reported initial experience with simultaneous versus staged resections, a French multicenter study showed an operative mortality of 7% for simultaneous 2% for staged surgery (31), while in a single center US study the mortality was 12% for simultaneous and 4% for staged resections (32). Several studies reported simultaneous operations performed without mortality, however patients were selected by experienced hepatobiliary surgeons and the major hepatectomies were avoided in elderly patients the same as in those with demanding colorectal surgery (33-36). In addition, since the surgical mortality rate is significantly higher when surgery of extensive hepatic resections is combined with colorectal resection (37), this approach should be only performed in carefully selected patients.

The standard staged operative treatment recommendations

in the literature suggest resection of the primary tumor followed by chemotherapy for 3-6 months and second stage of surgical treatment that includes liver surgery. The problem with this approach lies in the fact that liver metastases determine survival more intensive than the primary colorectal tumor. Chemotherapy can sometime not be performed after the surgical treatment of the primary tumor, especially when complicated by anastomotic leak or dehiscence, which occurs in 6-12% of patients (38,39). In cases of advanced rectal cancer usually a long term of radio-chemotherapy of 5 weeks is recommended and the second stage of operative treatment is planned 6-10 weeks following the neoadjuvant therapy. Therefore the patients do not receive a therapy of liver metastases for almost 15 weeks, which brings to the progress of liver metastatic disease (40). On the other hand some experimental studies have reported the rapid growth of metastases after removal of primary tumor (41,42). The underlying mechanism for those experimental results could be the loss of primary tumor-induced inhibition of angiogenesis in the metastases, which supports the founding of the increase of vascular density in humans after resection of primary tumor (43).

The reverse surgical approach onto the surgical treatment of colorectal liver metastases known as "liver-first" approach is reported as feasible and safe procedure with promising results, although it brings along the risk of bowel obstruction following the growth of primary tumor, which can be avoided by Hartmanns procedure (39,44). Results from the Liver Met Survey, involving 13,334 patients from 330 centers in 58 countries who underwent surgery for liver metastases, reported a better survival outcome in patients who undergo first resection of liver metastases than in those who do not (45). A recent systematic review of studies published in 1999-2010 confirmed these results and revealed 5-year survival rates for patients with liver metastases in the range of 16-74% (median, 38%) after liver resection (46).

The main idea of the "liver first" approach was to avoid the time loss between the operative therapy of primary tumor and the oncological therapy. Since the patients with rectal cancer often require a complex oncological therapy (chemotherapy, radiotherapy, and a complex pelvic operation), they could be the most proper candidates for such an approach (47). Despite liver-first patients usually have a greater hepatic disease burden and undergoing major resection more often, the reverse strategy was found safe and had long-term outcomes comparable to those of the other approaches (48).

How to achieve resectability without chemotherapy?

A large number of liver metastases should not be an absolute contraindication to surgery combined with chemotherapy provided that resection can be complete, with preservation of a functioning liver remnant of 25-30% (49). However, the problem is the loss of the proper functioning remnant volume of normal liver tissue, which presents an absolute contraindication for surgical resection. Advances in interventional radiology, particularly PVE in which the hypertrophy of normal liver tissue is provoked in order to ensure the proper remnant volume (50) and radiofrequency thermal ablation (RFA) widened the indications for surgical treatment of patients with colorectal cancer and liver metastases. In patients planned for major hepatectomies and with an otherwise normal liver, preoperative PVE is recommended when the ratio of the remnant liver to total liver volume is estimated to be less than 30%, whereas in patients with neoadjuvant chemotherapy this ratio is considered to be 40% (51,52). PVE is a safe procedure, but manipulation of the embolic material to the main portal vein or into branches that supply the future remnant liver remains a risk (1). RFA was initially anticipated for local treatment of hepatocellular carcinoma but has recently found application for the management of colorectal liver metastases, where its indications are still under doubt. Critical review of the results of RFA shows that it must be restricted in cases with a maximum of 3 lesions with the size of the biggest lesion less than 3 cm (53). Another limitation for the use of RFA in the management of colorectal liver metastases is the anatomic location of the lesion near big vessels, which increases the risk of incomplete ablation due to reduced heat effect that is used (54). A great indication of RFA is actually recurrence after resection, detected as small lesions, so it is possible not to interrupt chemotherapy (55).

A novel method in liver surgery that can solve the problem of remnant volume is the associating of liver partition and portal vein ligation (ALPPS) firstly reported 3 years ago (56). In ALPPS approach, the portal vein ligation associated with in situ splitting is able to induce enormously accelerated hypertrophy (57). The neovascularization and persistence of interlobar perfusion are prevented by performing parenchymal dissection and complete devascularization of segment IV (56). The nearly total parenchymal dissection induced a median hypertrophy of 74%, which is markedly above the range that can be achieved by portal vein ligation or PVE alone (58,59).

Conclusions

Surgical R0 resection still remains the only curative therapeutic tool in patients with colorectal cancer and liver metastases. The proper diagnostic algorithm is ultimate. The indications for surgical treatment are enlarged by the progress in neoadjuvant chemotherapy, diagnostic imaging, interventional radiology procedures especially the usage of PVE and radio frequent ablation. On the other hand the surgical techniques still develop producing the new pathways of treatment such as “liver-first approach” in the context of 2-stage operative therapy and ALPPS for the ensuring the remnant liver volume. Simultaneous liver and colorectal operations are feasible at carefully selected patients but should be avoided in cases of major hepatectomies, in elderly patients, and in patients with too complex intraoperative asset of colorectal tumor. The 2-stage hepatectomies as well as the “liver first” approach seem to become the new treatment strategies that improved the prognosis in patients in whom an R0 resection can be achieved with curative intention. The multidisciplinary treatment therapeutic approach in patients with colorectal cancer and liver metastases is essential to make the proper treatment plan and achieve the best results.

Acknowledgements

Disclosure: The authors declare no conflict of interest.

References

- Dimitroulis D, Nikiteas N, Troupis T, et al. Role of surgery in colorectal liver metastases: too early or too late? *World J Gastroenterol* 2010;16:3484-90.
- Ferlay J, Shin HR, Bray F, et al. GLOBOCAN 2008, Version 1.2, Cancer Incidence and Mortality Worldwide. IARC Cancer Base No.10. Available online: <http://globocan.iarc.fr>. accessed November 23, 2011.
- Jurisić I, Paradzik MT, Jurić D, et al. National program of colorectal carcinoma early detection in Brod-Posavina County (east Croatia). *Coll Antropol* 2013;37:1223-7.
- Milas J, Samardžić S, Miskulin M. Neoplasms (C00-D48) in Osijek-Baranja County from 2001 to 2006, Croatia. *Coll Antropol* 2013;37:1209-22.
- Samardžić S, Mihaljević S, Dmitrović B, et al. First six years of implementing colorectal cancer screening in the Osijek-Baranja County, Croatia--can we do better? *Coll Antropol* 2013;37:913-8.
- Sung JJ, Lau JY, Goh KL, et al. Increasing incidence of colorectal cancer in Asia: implications for screening. *Lancet Oncol* 2005;6:871-6.
- Boyle P, Leon ME. Epidemiology of colorectal cancer. *Br Med Bull* 2002;64:1-25.
- Khatri VP, Petrelli NJ, Belghiti J. Extending the frontiers of surgical therapy for hepatic colorectal metastases: is there a limit? *J Clin Oncol* 2005;23:8490-9.
- Van den Eynde M, Hendl isz A. Treatment of colorectal liver metastases: a review. *Rev Recent Clin Trials* 2009;4:56-62.
- Aloia TA, Adam R, Azoulay D, et al. Outcome following hepatic resection of metastatic renal tumors: the Paul Brousse Hospital experience. *HPB (Oxford)* 2006;8:100-5.
- Rees M, Tekkis PP, Welsh FK, et al. Evaluation of long-term survival after hepatic resection for metastatic colorectal cancer: a multifactorial model of 929 patients. *Ann Surg* 2008;247:125-35.
- Choti MA, Sitzmann JV, Tiburi MF, et al. Trends in long-term survival following liver resection for hepatic colorectal metastases. *Ann Surg* 2002;235:759-66.
- Krebs B, Kozelj M, Potrc S. Rectal cancer treatment and survival--comparison of two 5-year time intervals. *Coll Antropol* 2012;36:419-23.
- Poston GJ, Figueras J, Giuliani F, et al. Urgent need for a new staging system in advanced colorectal cancer. *J Clin Oncol* 2008;26:4828-33.
- Boras Z, Kondza G, Sisljagić V, et al. Prognostic factors of local recurrence and survival after curative rectal cancer surgery: a single institution experience. *Coll Antropol* 2012;36:1355-61.
- Fong Y, Fortner J, Sun RL, et al. Clinical score for predicting recurrence after hepatic resection for metastatic colorectal cancer: analysis of 1001 consecutive cases. *Ann Surg* 1999;230:309-18; discussion 318-21.
- Mayo SC, Heckman JE, Shore AD, et al. Shifting trends in liver-directed management of patients with colorectal liver metastasis: a population-based analysis. *Surgery* 2011;150:204-16.
- Asiyanbola B, Chang D, Gleisner AL, et al. Operative mortality after hepatic resection: are literature-based rates broadly applicable? *J Gastrointest Surg* 2008;12:842-51.
- Kodama Y, Ng CS, Wu TT, et al. Comparison of CT methods for determining the fat content of the liver. *AJR Am J Roentgenol* 2007;188:1307-12.
- Reeder SB, Sirlin CB. Quantification of liver fat with magnetic resonance imaging. *Magn Reson Imaging Clin N Am* 2010;18:337-57.

21. van Werven JR, Marsman HA, Nederveen AJ, et al. Assessment of hepatic steatosis in patients undergoing liver resection: comparison of US, CT, T1-weighted dual-echo MR imaging, and point-resolved 1H MR spectroscopy. *Radiology* 2010;256:159-68.
22. Overman MJ, Maru DM, Charnsangavej C, et al. Oxaliplatin-mediated increase in spleen size as a biomarker for the development of hepatic sinusoidal injury. *J Clin Oncol* 2010;28:2549-55.
23. Shin NY, Kim MJ, Lim JS, et al. Accuracy of gadoteric acid-enhanced magnetic resonance imaging for the diagnosis of sinusoidal obstruction syndrome in patients with chemotherapy-treated colorectal liver metastases. *Eur Radiol* 2012;22:864-71.
24. Zorzi D, Laurent A, Pawlik TM, et al. Chemotherapy-associated hepatotoxicity and surgery for colorectal liver metastases. *Br J Surg* 2007;94:274-86.
25. Chami L, Lassau N, Malka D, et al. Benefits of contrast-enhanced sonography for the detection of liver lesions: comparison with histologic findings. *AJR Am J Roentgenol* 2008;190:683-90.
26. Leen E, Ceccotti P, Moug SJ, et al. Potential value of contrast-enhanced intraoperative ultrasonography during partial hepatectomy for metastases: an essential investigation before resection? *Ann Surg* 2006;243:236-40.
27. Necib H, Garcia C, Wagner A, et al. Detection and characterization of tumor changes in 18F-FDG PET patient monitoring using parametric imaging. *J Nucl Med* 2011;52:354-61.
28. Hendlisz A, Golfopoulos V, Garcia C, et al. Serial FDG-PET/CT for early outcome prediction in patients with metastatic colorectal cancer undergoing chemotherapy. *Ann Oncol* 2012;23:1687-93.
29. Moulton C, Levine MN, Law C, et al. An Ontario Clinical Oncology Group (OCOG) randomized controlled trial (RCT) assessing FDG PET/CT in resectable liver colorectal adenocarcinoma metastases (CAM). *J Clin Oncol* 2011;29:abstr 3520.
30. Rees M, Tekkis PP, Welsh FK, et al. Evaluation of long-term survival after hepatic resection for metastatic colorectal cancer: a multifactorial model of 929 patients. *Ann Surg* 2008;247:125-35.
31. Nordlinger B, Guiguet M, Vaillant JC, et al. Surgical resection of colorectal carcinoma metastases to the liver. A prognostic scoring system to improve case selection, based on 1568 patients. *Association Française de Chirurgie. Cancer* 1996;77:1254-62.
32. Bolton JS, Fuhrman GM. Survival after resection of multiple bilobar hepatic metastases from colorectal carcinoma. *Ann Surg* 2000;231:743-51.
33. Weber JC, Bachellier P, Oussoultzoglou E, et al. Simultaneous resection of colorectal primary tumour and synchronous liver metastases. *Br J Surg* 2003;90:956-62.
34. Chua HK, Sondena K, Tsiotos GG, et al. Concurrent vs. staged colectomy and hepatectomy for primary colorectal cancer with synchronous hepatic metastases. *Dis Colon Rectum* 2004;47:1310-6.
35. Tanaka K, Shimada H, Matsuo K, et al. Outcome after simultaneous colorectal and hepatic resection for colorectal cancer with synchronous metastases. *Surgery* 2004;136:650-9.
36. Minagawa M, Yamamoto J, Miwa S, et al. Selection criteria for simultaneous resection in patients with synchronous liver metastasis. *Arch Surg* 2006;141:1006-12; discussion 1013.
37. Reddy SK, Pawlik TM, Zorzi D, et al. Simultaneous resections of colorectal cancer and synchronous liver metastases: a multi-institutional analysis. *Ann Surg Oncol* 2007;14:3481-91.
38. Chiappa A, Biffi R, Bertani E, et al. Surgical outcomes after total mesorectal excision for rectal cancer. *J Surg Oncol* 2006;94:182-93; discussion 181.
39. Tran CL, Udani S, Holt A, et al. Evaluation of safety of increased time interval between chemoradiation and resection for rectal cancer. *Am J Surg* 2006;192:873-7.
40. Verhoef C, van der Pool AE, Nuyttens JJ, et al. The "liver-first approach" for patients with locally advanced rectal cancer and synchronous liver metastases. *Dis Colon Rectum* 2009;52:23-30.
41. O'Reilly MS, Holmgren L, Shing Y, et al. Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma. *Cell* 1994;79:315-28.
42. O'Reilly MS, Holmgren L, Chen C, et al. Angiostatin induces and sustains dormancy of human primary tumors in mice. *Nat Med* 1996;2:689-92.
43. Peeters CF, de Waal RM, Wobbes T, et al. Outgrowth of human liver metastases after resection of the primary colorectal tumor: a shift in the balance between apoptosis and proliferation. *Int J Cancer* 2006;119:1249-53.
44. Mentha G, Roth AD, Terraz S, et al. 'Liver first' approach in the treatment of colorectal cancer with synchronous liver metastases. *Dig Surg* 2008;25:430-5.
45. Registry of patients operated for Colorectal Liver Metastasis. Available online: <http://www.livermetsurvey.org>. International livermetsurvey.org, accessed November 23, 2011.

46. Kanas GP, Taylor A, Primrose JN, et al. Survival after liver resection in metastatic colorectal cancer: review and meta-analysis of prognostic factors. *Clin Epidemiol* 2012;4:283-301.
47. Andres A, Toso C, Adam R, et al. A survival analysis of the liver-first reversed management of advanced simultaneous colorectal liver metastases: a LiverMetSurvey-based study. *Ann Surg* 2012;256:772-8; discussion 778-9.
48. Mayo SC, Pulitano C, Marques H, et al. Surgical Management of Patients with Synchronous Colorectal Liver Metastasis: A Multicenter International Analysis. *J Am Coll Surg* 2013;216: 707-16.
49. Adam R, De Gramont A, Figueras J, et al. The oncosurgery approach to managing liver metastases from colorectal cancer: a multidisciplinary international consensus. *Oncologist* 2012;17:1225-39.
50. Makuuchi M, Thai BL, Takayasu K, et al. Preoperative portal embolization to increase safety of major hepatectomy for hilar bile duct carcinoma: a preliminary report. *Surgery* 1990;107:521-7.
51. Abdalla EK, Barnett CC, Doherty D, et al. Extended hepatectomy in patients with hepatobiliary malignancies with and without preoperative portal vein embolization. *Arch Surg* 2002;137:675-80; discussion 680-1.
52. Vauthey JN, Chaoui A, Do KA, et al. Standardized measurement of the future liver remnant prior to extended liver resection: methodology and clinical associations. *Surgery* 2000;127:512-9.
53. Berber E, Pelley R, Siperstein AE. Predictors of survival after radiofrequency thermal ablation of colorectal cancer metastases to the liver: a prospective study. *J Clin Oncol* 2005;23:1358-64.
54. Terraz S, Constantin C, Majno PE, et al. Image-guided multipolar radiofrequency ablation of liver tumours: initial clinical results. *Eur Radiol* 2007;17:2253-61.
55. Mentha G, Majno P, Terraz S, et al. Treatment strategies for the management of advanced colorectal liver metastases detected synchronously with the primary tumour. *Eur J Surg Oncol* 2007;33 Suppl 2:S76-83.
56. Baumgart J, Lang S, Lang H. A new method for induction of liver hypertrophy prior to right trisectionectomy: A report of three cases HPB (Oxford, England) 2011;13:71-2.
57. Torres OJ, Moraes-Junior JM, Lima e Lima NC, et al. Associating liver partition and portal vein ligation for staged hepatectomy (ALPPS): a new approach in liver resections. *Arq Bras Cir Dig* 2012;25:290-2.
58. de Santibañes E, Alvarez FA, Ardiles V. How to avoid postoperative liver failure: a novel method. *World J Surg* 2012;36:125-8.
59. Schnitzbauer AA, Lang SA, Goessmann H, et al. Right portal vein ligation combined with in situ splitting induces rapid left lateral liver lobe hypertrophy enabling 2-staged extended right hepatic resection in small-for-size settings. *Ann Surg* 2012;255:405-14.

Cite this article as: Patrlj L, Kopljar M, Kliček R, Hrelec Patrlj M, Kolovrat M, Rakić M, Đuzel A. The surgical treatment of patients with colorectal cancer and liver metastases in the setting of the “liver first” approach. *Hepatobiliary Surg Nutr* 2014;3(5):324-329. doi: 10.3978/j.issn.2304-3881.2014.09.12