

Biological aspects of bone, cartilage and tendon regeneration

Pećina, Marko; Vukičević, Slobodan

Source / Izvornik: **International Orthopaedics, 2007, 31, 719 - 720**

Journal article, Accepted version

Rad u časopisu, Završna verzija rukopisa prihvaćena za objavljivanje (postprint)

<https://doi.org/10.1007/s00264-007-0425-7>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:105:445242>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-06-03**



Repository / Repozitorij:

[Dr Med - University of Zagreb School of Medicine
Digital Repository](#)





Središnja medicinska knjižnica

Pećina, M., Vukičević, S. (2007) *Biological aspects of bone, cartilage and tendon regeneration*. International Orthopaedics, 31 (6). pp. 719-720.

The original publication is available at www.springerlink.com

<http://www.springerlink.com/content/y553092505954178/>

<http://dx.doi.org/10.1007/s00264-007-0425-7>

<http://medlib.mef.hr/308>

University of Zagreb Medical School Repository

<http://medlib.mef.hr/>

EDITORIAL

Biological aspects of bone, cartilage and tendon regeneration

Marko Pecina and Slobodan Vukicevic

Current orthopedic procedures in supporting regeneration of bone, cartilage and tendon are dependant on our understanding of the molecular processes responsible for tissue repair. At present we know how to regenerate bone when physiological mechanisms of fracture repair fail [13]. Since the original description of the potential of demineralized bone matrix to induce bone at an ectopic site it took more than three decades to bring bone morphogenetic proteins (BMPs) to clinical use. By the end of 2007 nearly one million patients worldwide will be treated with BMPs for spinal fusions, non-unions, acute fractures and maxillofacial reconstruction. Use of animal models, genomics and proteomics deciphered new mechanisms and candidate molecules for the regeneration of joint cartilage and tendons, opening new avenues in regenerative orthopedics.

This special volume reviews novel strategies in the regeneration of bone, cartilage and tendon. Bishop and Einhorn [2] describe the clinical use of recombinant BMP-2 and BMP-7. They predict that current techniques in engineering bone for restoring defined skeletal defects represent a unique opportunity for BMPs in the future. McKay and colleagues [9] reviewed the clinical applications of BMP-2 (INFUSE Bone Graft), while Vaccaro, McKee and colleagues [14] describe the clinical applications of BMP-7 (OP-1 implant or Osigraft). Authors suggest that recombinant human BMP based devices, when properly applied, can eliminate the need to harvest autologous bone for grafting procedures benefiting both the surgeon and patients. Grgurevic and collaborators [6] describe molecules discovered by proteomic analysis in the plasma of patients with an acute bone fracture. The characterization and use of potential new biomarkers, like TGF- β induced protein IG-H3, cartilage acidic protein 1, procollagen C proteinase enhancer protein and TGF- β receptor III, for bone and cartilage regeneration are discussed. Sendak and colleagues [11] review the newly discovered role of thyroid stimulating hormone (TSH) and follicle stimulating hormone (FSH) in bone

remodeling. It has been recently found that low doses of TSH increase bone volume and improve bone microarchitecture and strength in aged osteoporotic rats, suggesting TSH directly affects bone remodeling in vivo [10]. Grasser and collaborators [5], by using a genomic approach, deciphered the potential mechanism of bone augmentation following systemic administration of BMP-6, suggesting that IGF-1 and EGF mediate, at least in part, the effects of BMP-6 on bone, which has been recently reported [12]. The role of prostaglandin E2 receptors and their analogs in bone formation is reviewed by Li and colleagues [8] demonstrating that the PGE2 receptor is essential for restoring bone defects in animal models. Cartilage regeneration and chronic back pain due to degenerative disc disease are among major obstacles in orthopedics. Chubinskaya and colleagues [4] summarized the role of osteogenic protein-1 (BMP-7) in animal models of osteochondral and chondral defects, osteoarthritis and degeneration in intervertebral disc cartilage. They show that OP-1 is exhibiting unique pro-anabolic and prominent anti-catabolic properties having a potential for treating cartilage and disc lesions.

Overuse tendon disorders are common and related to a significant morbidity. Tendon and ligament function engineering and repair are discussed in articles by Aspenberg [1] and by Hoffman and Gross [7]. Aspenberg found that the repair of subcutaneous tendon ruptures can be stimulated by a single application of one of several growth factors, including PDGF, TGF- β , IGF-1, VEGF and GDF-5,-6,-7, or by a thrombocyte concentrate. The response is dependent on the mechanical microenvironment, which is crucial for the repair process.

Hoffmann and Gross specifically discuss gene-therapeutic approaches using mesenchymal stem cells for a potential future application in tendon/ligament regeneration.

Borovecki and colleagues [3] discuss the potential of a genomic approach in analyzing bone homeostasis and diseases. Gene expression profiling studies yielded novel insights into the complex interplay of osteoblast and osteoclast regulation, paracrine and endocrine control of bone and cartilage remodeling, as well as the pathophysiology of osteoporosis and bone tumors.

We believe that this volume will provide basic and clinical scientists with the state of art in regenerative orthopedics.

References:

1. Aspenberg P (2007) Stimulation of tendon repair: Mechanical loading, GDFs and platelets. A mini-review. Int Orthop 31: -
2. Bishop GB, Einhorn TA (2007) Current and future clinical application of bone morphogenetic proteins in orthopaedic trauma surgery. Int Orthop 31: -
3. Borovecki F, Pecina-Slaus N, Vukicevic S (2007) Biological mechanisms of bone and cartilage remodeling - Genomic perspective. Int Orthop 31: -
4. Chubinskaya S, Hurtig M, Rueger D (2007) OP-1/BMP-7 in cartilage rapair. Int Orthop 31: -
5. Grasser WA, Orlic I, Borovecki F, Riccardi KA, Simic P, Vukicevic S and Paralkar VM (2007) BMP-6 exerts its osteoinductive effect through activation of IGF-I and EGF pathways. Int Orthop 31: -
6. Grgurevic L, Macek B, Durdevic D, Vukicevic S (2007) Detection of bone and cartilage-related proteins in plasma of patients with a bone fracture using liquid chromatography-mass spectrometry. Int Orthop 31: -
7. Hoffmann A, Gross G (2007) Tendon and ligament engineering in the adult organism: Mesenchymal stem cells and gene therapeutic approaches. Int Orthop 31: -
8. Li M, Paralkar VM, Thompson DD (2007) Prostaglandin E2 receptors in bone formation. Int Orthop 31: -
9. McKay WF, Peckham SM, Badura JM (2007) Comprehensive clinical review of recombinant human bone morphogenetic protein-2 (INFUSE[®] Bone Graft). Int Orthop 31: -
10. Sampath TK, Simic P, Sendak R, Draca N, Bowe AE, O'brien S, Schiavi SC, McPherson JM, Vukicevic S (2007) Thyroid stimulating hormone restores bone volume, microarchitecture and strength in aged ovariectomized rats. J Bone Miner Res 22:849-859
11. Sendak RA, Sampath KT, McPherson JM (2007) Newly reported roles of thyroid stimulating hormone and follicle stimulating hormone in bone remodeling. Int Orthop 31: -

12. Simic P, Buljan Culej J, Orlic I, Grgurevic L, Draca N, Spaventi R, Vukicevic S (2006) Systemically administered bone morphogenetic protein-6 restores bone in aged ovariectomized rats by increasing bone formation and suppressing bone resorption. *J Biol Chem* 281:25509-25521
13. Simic P, Vukicevic S (2007) Bone Morphogenetic Proteins: from developmental signals to tissue regeneration. Conference on bone morphogenetic proteins. *EMBO Rep* 8:327-331
14. White AP, Vaccaro AR, Hall JA, Whang PG, Friel BC, McKee MD (2007) Clinical applications of BMP-7/OP-1 in fractures, nonunions and spinal fusion. *Int Orthop* 31: -