

# First report on PVL-positive methicillin-resistant Staphylococcus aureus of SCCmec type V, spa type T441 in Croatia

---

**Budimir, Ana; Tićac, Brigita; Rukavina, Tomislav; Farkaš, Maja; Kalenić, Smilja**

*Source / Izvornik:* **Collegium Antropologicum, 2016, 40, 133 - 137**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

*Permanent link / Trajna poveznica:* <https://urn.nsk.hr/urn:nbn:hr:105:844109>

*Rights / Prava:* [In copyright](#) / [Zaštićeno autorskim pravom.](#)

*Download date / Datum preuzimanja:* **2024-08-01**



*Repository / Repozitorij:*

[Dr Med - University of Zagreb School of Medicine  
Digital Repository](#)



# First Report on PVL-Positive Methicillin-Resistant *Staphylococcus Aureus* of *Sccmec* Type V, *SPA* Type T441 in Croatia

Ana Budimir<sup>1,2</sup>, Brigita Tićac<sup>2</sup>, Tomislav Rukavina<sup>2</sup>, Maja Farkaš<sup>2</sup> and Smilja Kalenić<sup>3</sup>

<sup>1</sup> Department of Clinical and Molecular Microbiology, University Hospital Centre Zagreb, Zagreb, Croatia

<sup>2</sup> University of Rijeka, School of Medicine, Department of Microbiology, Teaching Institute of Public Health, Rijeka, Croatia

<sup>3</sup> University of Zagreb, School of Medicine, Zagreb, Croatia

## ABSTRACT

The aim of the study was to investigate the molecular epidemiology of MRSA in Primorsko-Goranska County of Croatia during a six-year period (2001–2007). In period from 2001 and 2007, 46 MRSA isolates were collected in Rijeka, strains were subjected to susceptibility testing according to CLSI guidelines, *mecA* gene detection and *SCCmec* typing as well as detection of PVL. Strains were typed by Pulse Field Gel Electrophoresis (PFGE) and *spa* typing. All isolates were susceptible to vancomycin, linezolid, mupirocin, nitrofurantoin, only one strain was resistant to fusidic acid and co-trimoxazole. Results of *SCCmec* typing showed the presence of *SCCmec* type IV in 26 MRSA strains, *SCCmec* type V in three strains, and 13 strains comprised *SCCmec* I. *SCCmec* type II and III were not observed. Four MRSA strains were non-typeable by applied *SCCmec* typing methods. PVL was detected in 4 strains, two *SCCmec* IV and two *SCCmec* V. PFGE analysis, grouped MRSA strains into six similarity groups and 18 singletons. Dominating *spa* types in this collection of strains were t015, with 15 strains, followed by t041 (N=7), t051, (N=2), t2850 (N=2), t008 (N=2) and single isolates t441, t002, t448, t018, t019, t355, t390, t026, t449, t148. We also detected two new *spa* types, t3510 and t3509, respectively. This is the first report on *SCCmec* type V in Croatia, and, to our knowledge, first report of PVL-positive methicillin-resistant *Staphylococcus aureus* *SCCmec* type V and T441 (ST59-MRSA-V) in this part of Europe.

**Key words:** MRSA, Croatia, *SCCmec*, *spa* type

## Introduction

*Staphylococcus aureus*, the most virulent *Staphylococcus* species, is also second most common isolate from patients in outpatient settings. *S. aureus* causes a wide range of syndromes, from minor skin and soft tissue infections to life-threatening pneumonia and toxicosis [1]. Rates of methicillin – resistance increased slowly, but progressively over decades. Increase in methicillin resistance worldwide was accompanied by isolation of MRSA isolates from community-acquired infections among previously healthy individuals with few or not traditional healthcare-associated risk factors<sup>1</sup>.

The increase in the rate of CA-MRSA infections is a reason for public health concerns, and a challenge for infection control, since MRSA is found in nursing homes,

kindergartens, and schools<sup>2</sup>. However, studies on the CA-MRSA prevalence are sporadic. HA-MRSA and CA-MRSA can be distinguished by their genetic background, the resistance determinant Staphylococcal Cassette Chromosome *mec* (*SCCmec*), and the presence of Pantone-Valentine leukocidin (PVL)<sup>3</sup>.

The aim of the present study was to investigate the molecular epidemiology of MRSA in Primorsko-Goranska County of Croatia during a six-year period (2001–2007).

Prevalence of MRSA in Croatia in blood culture isolates was 36,3 %, in 2009 as presented by European Resistance Surveillance System (EARSS) website<sup>4</sup>.

Primorsko-Goranska County is situated in western part of Croatia, as seen at Figure 1., covering an area of 3.582 km<sup>2</sup>. Largest city and county center is Rijeka, third largest city in Croatia. Public Health and Teaching Insti-

tute in Rijeka is leading public health and teaching institution in county with gravitating population of approximately 300 000 inhabitants (around 7% of Croatian population). Department of microbiology mainly process specimens from outpatient population and ambulatory care.

## Methods

In period from 2001 and 2007, all MRSA strains isolated (N=46) were collected and stored until further analysis. Strains originated from nasal swabs (N=20), 15 from wound swabs, and two or one isolate per following sites: skin infection sample, furunculosis, conjunctiva, tongue.

All isolates were identified based on colony morphology, catalase, coagulase and DNA-se presence. Isolates were screened for MRSA with oxacillin, cefoxitin disk and latex agglutination test.

All 46 MRSA strains were subjected to further analysis in Clinical Hospital Centre Zagreb, Department of Clinical and Molecular Microbiology and some non-typeable strains were submitted to University Hospital Maastricht, Maastricht Infection Center for additional typing.

Susceptibility testing was performed according to CLSI guidelines<sup>5</sup> for the following antibiotics: penicillin, oxacillin, cefoxitin, erythromycin, clindamycin, azithromycin, gentamycin, tetracycline, rifampicin, trimethoprim/sulfamethoxazole, linezolid, ciprofloxacin, vancomycin, nitrofurantoin, trimethoprim, except for fusidic acid and mupirocin, which were interpreted according to the criteria of the French Society of Microbiology (CAFM)<sup>6</sup>.

*mecA* gene detection and SCCmec typing was performed according to Oliveira et al.<sup>7</sup> with some modifications according to Deurenberg et al.<sup>8</sup> and for some strains, fluorescent PCR assay was used<sup>9</sup>, partially based on Zhang multiplex PCR method<sup>10</sup>. Detection of PVL was performed as previously described<sup>11</sup>.

Pulse Field Gel Electrophoresis (PFGE) was performed as previously described<sup>12</sup>.

PFGE patterns were analyzed and compared with use of GelCompar III software (Applied Maths, Sint-Martens-Latem, Belgium). Spa typing was performed according to SeqNet protocol, spa types were assigned by using Ridom Spa software (Ridom, GmbH, Germany) and synchronization with Ridom server<sup>13</sup>.

BURP alignment was used for spa CC alignment.

## Results

All strains were resistant to penicillin, oxacillin and cefoxitin, 35% were resistant to erythromycin, 30% to clindamycin, 35% to azithromycin, 30% to gentamycin, 2% to tetracycline (17,4 % were intermediate resistant to tetracycline), 13% to rifampicin (4% intermediate resistant to rifampicin), 37 % to ciprofloxacin (2% intermediate resistant to ciprofloxacin), 4,3% to trimethoprim (2% intermediate resistant to trimethoprim). All isolates were susceptible to vancomycin, linezolid, mupirocin, nitrofurantoin,

only one strain was resistant to fusidic acid and co-trimoxazole.

Typing results are shown in Figure 1.

All isolates were *mecA* positive. Results of SCCmec typing showed the presence of SCCmec type IV in 26 MRSA strains, SCCmec type V in three strains, and 13 strains comprised SCCmec I. SCCmec type II and III were not observed in study collection.

Four MRSA strains were non-typeable by applied SCCmec typing methods.

PVL was detected in 4 strains, two SCCmec IV and two SCCmec V.

PFGE analysis, using Tenover criteria, grouped MRSA strains into six similarity groups (A, B, C, D, E, F) containing 9, 9, 6, 2, and 2 isolates, respectively, and 18 singletons.

Dominating spa types in this collection of strains were t015, with 15 strains, followed by t041 (N=7), t051, (N=2), t2850 (N=2), t008 (N=2) and single isolates t441, t002, t448, t018, t019, t355, t390, t026, t449, t148. We also detected two new spa types, t3510 and t3509, respectively.

MRSA strains SCCmec type V are spa typed as t002 (assigned ST-5), t355 (assigned ST-377), t441 (assigned ST-59).

Isolates were classified into 2 clusters: CC 1 with t008, t024 and t051, and other cluster, CC41 consisting of t002, t041 and t2027.

## Discussion

With efforts to prevent the spread of MRSA in hospitals and raised awareness of possible routes of transmission, encouraged by hospital administration and, in some counties, media campaign and Government involvement, the MRSA rates decreased in some countries<sup>14</sup>.

Worrying are reports and suggestions that CA MRSA is more and more present in hospitals. Children, young without risk factors are carriers and sometimes, they develop clinical infection, ranging from mild skin infection to severe skin and soft-tissue infections, pneumonia<sup>15,16</sup> etc. CA-MRSA strains can cause infections with rapidly progressive, fatal disease including necrotizing pneumonia, severe sepsis and necrotizing fasciitis<sup>1</sup>.

There are few data on prevalence and structure of community-associated MRSA strains in Europe<sup>17–19</sup> although some reports are showing increase in this distinct population of bacteria. In USA, majority of bloodstream infections<sup>20</sup> and especially skin and soft tissue infections<sup>21</sup> are caused by CA MRSA.

Genetic backgrounds of CA-MRSA isolates are distinct in different geographic locations<sup>22,11</sup> and many different backgrounds can exist within a narrow geographic area.

The most widely used molecular typing method for study of MRSA epidemiology is Pulse Field Gel Electrophoresis, PFGE, suitable method for the determination of clonal relationships but not for long term epidemiological




Fig. 1. Dendrogram of MRSA strains from Rijeka.

investigation. Spa typing is proposed as an accurate method for typing of *S. aureus*<sup>23</sup>, and it has the advantages of speed, interpretation and interlaboratory comparison.

In this study, we combined sequence-based, single locus-oriented spa typing method with whole-genome, restriction-based methods in order to compare isolates within larger time frame. For complete nomenclature of MRSA it is important to perform SCCmec typing as well.

Our previous reports suggested the dominance of ST-247/250 MRSA-I among MRSA strains described<sup>24</sup> and in collection of strains in 2004 (data not shown).

There were sporadic reports on community-associated MRSA in Croatia<sup>25</sup>, isolated from non-infected body sites or skin and soft tissue infections, but this is the first report on SCCmec type V in Croatia, and, to our knowledge, in this part of Europe.

In this study we showed presence of PVL-positive strains, in four cases. Some epidemiologic and clinical

data provide compelling evidence that the high virulence potential of CA-MRSA is associated with genes lukS-PV and lukF-PV (pvl) encoding the subunits of the Panton-Valentine leukocidin.

Seems that clinical sequelae of pvl- positive infections are more severe than pvl-negative *S. aureus*<sup>26</sup>.

The interesting finding is that half of PVL-positive isolates is characterized as MRSA SCCmec V, which is not an common finding, described in severe infection in Greece<sup>27</sup>, but within different clonal lineage (ST-80).

SCCmec type V is typical for Australia but is usually PVL-negative. Strains causing infections with SCCmec V and PVL positive; ST 59 are observed in Taiwan, especially in children. This clone, observed in Taiwan is also present in hospitals in significant proportion[28].

This work is showing that there is a certain number of MRSA strains in the community of this area, with heterogenous background. This is in concordance with similar studies in some European countries<sup>29,17,18</sup>, but some clonal lineages are described in this region (Southern Eastern Europe) for the first time.

Predominant spa type in this collection of strains, t015 is observed in Croatia, in eastern, western parts of country, and Istria, area which is situated close to Primorsko-goranska county.

It is obvious that some strains, although not having typical and direct hospital background, ended originated in hospitals, which is well known for spa type t041.

t041 is an frequently encountered spa type in Croatia, and characterizing typical hospital-associated MRSA together with SCCmec type I, also found in Italy [30]. Two of predominating spa types are grouped in two major

similarity groups, B, C and D, based on PFGE typing. There are some overlaps, and singletons are not grouped in between genetically related strains.

t002 is common spa type, isolated in Germany, Austria, Spain, Israel, Canada, USA, UK,

Switzerland etc<sup>31–33,13,30</sup>.

spa type t355 is observed in Austria, Slovenia, Germany, Norway, Netherlands, Turkey, Israel, Sweden, Iceland, Finland, France<sup>13,34</sup>.

Strains typed as t441 were previously observed in Sweden, Norway, Netherlands, Iceland, Taiwan<sup>13</sup>.

In conclusion, CA-MRSA isolates have been emerging around the world in different *S. aureus* genetic lineages and are beginning to enter in healthcare environments where they could become endemic.

The strong epidemiologic link between PVL and CA-MRSA disease leaves little doubt that PVL must play an important role in the pathogenesis and course of disease; however, Croatian SCCmec IV MRSA isolates were mostly PVL-negative, and there were SCCmec V-PVL positive isolates which suggests that there is a lot of genetic exchange and variability among MRSA isolates from community.

## Acknowledgments

We would like to thank Dr. T. Ito (Department of Bacteriology, Juntendo University; Tokyo, Japan) and Dr. D. Oliveira (Laboratório de Genética Molecular, Instituto de Tecnologia Química e Biológica, Oeiras, Portugal) for providing the reference strains for SCCmec typing.

## REFERENCES

- BOYLE-VAVRA S, DAUMRS, Laboratory Investigation 87 (2007) 3. — 2. URTH T, JUUL G, SKOV R, SCHONHEYDER H, Spread of a methicillin-resistant staphylococcus aureus st80-iv clone in a danish community, Infection Control Hospital Epidemiology 252 (2005) 144 — 3. DEURENBERG RH, VINK C, KALENIC S, FRIEDRICH AW, BRUGGEMAN CA, STOBBERINGH EE, The molecular evolution of methicillin-resistant staphylococcus aureus. Clinical Microbiology and Infection 13 (3) (2006) 222. doi:10.1111/j.1469-0691.2006.01573.x — 4. http://www.rivm.nl/earss/database/. — 5. Clinical and laboratory standards institute N, Performace standards for antimicrobial susceptibility testing (2006) — 6. Comité de l'antibiogramme de la société française de microbiologie (2004). http://www.sfmasso.fr/nouv/general/php — 7. OLIVEIRA DC, DE LENCASTRE H Antimicrob Agents Chemother 46 (7) (2002) 2155. — 8. DEURENBERG RH, VINK C, OUDHUIS GJ, MOOIJ JE, DRIESSEN C, COPPENS G, CRAEGHS J, DE BRAUWER E, LEMMEN S, WAGENVOORT H, FRIEDRICH AW, SCHERES J, STOBBERINGH EE, Antimicrob Agents Chemother 49 (10) (2005) 4263. — 9. VALVATNE H, RIJNDERS MI, BUDIMIR A, BOUMANS ML, DE NEELING AJ, BEISSER PS, STOBBERINGH EE, DEURENBERG RH, A rapid, 2-well, multiplex real-time polymerase chain reaction assay for the detection of scmec types i to v in methicillin-resistant staphylococcus aureus. Diagn Microbiol Infect Dis 65 (4) (2009) 384. doi:S0732-8893(09)00335-6 [pii] 10.1016/j.diagmicrobio.2009.08.006 — 10. ZHANG K, MCCLURE JA, ELSAYED S, LOUIE T, CONLY JM, J Clin Microbiol 43 (10) (2005) 5026. — 11. LINA G, PIEMONT Y, GODAIL-GAMOT F, BES M, PETER MO, GAUDUCHON V, VANDENESCH F, ETIENNE J, Clin Infect Dis 29 (5) (1999) 1128. — 12. GOERING RV, Pulsed-field gel electrophoresis. In:

Persing DH (ed) Molecular microbiology: Diagnostic principles and practice. ASM Press, Washington D.C. (2004) — 13. http://spaserver.ridom.de/spaserver/spa-t441.shtml). — 14. LIEBOWITZ LD, Int J Antimicrob 34 (3) (2009) 11. doi:S0924-8579(09)70551-5 [pii] 10.1016/S0924-8579(09)70551-5 — 15. SALGADO CD, FARR BM, CALFEE DP, Clin Infect Dis 36 (2) (2003) 131. — 16. ALONSO-TARRES C, VILLEGAS ML, DE GISPERT FJ, CORTES-LLETGET MC, ROVIRA PLARROMANIA, ETIENNE J, Eur J Clin Microbiol Infect Dis 24 (11) (2005) 756. — 17. HANSEN AM, FOSSUM A, MIKALSEN J, HALVORSEN DS, BUKHOLM G, SOLLID JU, J Clin Microbiol 43 (5) (2005) 2118. — 18. HARBARTH S, FRANCOIS P, SHRENZEL J, FANKHAUSER-RODRIGUEZ C, HUGONNET S, KOESSLER T, HUYGHE A, PITTET D, Emerg Infect Dis 11 (6) (2005) 962-965 — 19. FARIA NA, OLIVEIRA DC, WESTH H, MONNET DL, LARSENAR, SKOV R, DE LENCASTRE H, J Clin Microbiol 43 (4) (2005) 1836. — 20. SEYBOLD U, KOURBATOVA EV, JOHNSON JG, HALVOSA SJ, WANG YF, KING MD, RAY SM, BLUMBERG HM, Clin Infect Dis 42 (5) (2006) 647. — 21. NOSKIN GA, RUBIN RJ, SCHENTAG JJ, KLUYTMANS J, HEDBLUM EC, SMULDERS M, LAPETINA E, GEMMEN E, Arch Intern Med 165 (2005) 1756. — 22. VANDENESCH F, NAIMI T, ENRIGHT MC, LINA G, NIMMO GR, HEFFERNAN H, LIASSINE N, BES M, GREENLAND T, REVERDY ME, ETIENNE J, Em erg Infect Dis 9 (8) (2003) 978-984 — 23. FRENAY HM, BUNSCHOTEN AE, SCHOULS LM, VAN LEEUWEN W, VANDENBROUCKE-GRAULS CM, VERHOEF J, MOOIJ J, Eur J Clin Microbiol Infect Dis 15 (1996) 60. — 24. BUDIMIR A, DEURENBERG RH, PLECKO V, VINK C, KALENIC S, STOBBERINGH EE, J Antimicrob Chemother 57 (2) (2006) 331. — 25. KRZYSTON-RUSSJAN

- J, TAMBIC-ANDRASEVIC A, BUKOVSKI S, SABATA, HRYNIEWICZ W, Clin Microbiol Infect 12 (7) (2006) 697. — 26. GILLET Y, ISSARTEL B, VANHEMS P, GODAIL-GAMOT F, Association between staphylococcus aureus strains carrying gene for panton-valentine leukocidin and highly lethal necrotising pneumonia in young immunocompetent patients. Lancet (2002) — 27. IRINI G, GEORGIOS M, KONSTANTINOS G, ANTONIOS M, IRIS S, EFI P, Diagn Microbiol Infect Dis — 28. HUANG YC, HO CF, CHEN CJ, SU LH, LIN TY, Clin Microbiol Infect 14 (2008) 1167. — 29. HALLIN M, DENIS O, DEPLANO A, DE MENDONCA R, DE RYCK R, ROTTIERS S, STRUELENS MJ, J Antimicrob Chemother 59 (2007) 465. — 30. GRUNDMANN H, AANENSEN DM, VAN DEN WIJNGAARD CC, SPRATT BG, HARMSSEN D, FRIEDRICH AW, PLoS Med 7 (1) (2007) :e1000215. doi:10.1371/journal.pmed.1000215 — 31. DEURENBERG R, VINK C, S K, FRIEDRICH A, BRUGGEMAN C, STOBBERINGH EE, Clin Microbiol Infect 13 (2008) 222. — 32. WU D, WANG Q, YANG Y, GENG W, YU S, YAO K, YUAN L, SHEN X, Diagn Microbiol Infect Dis 67 (1) (2008) 1. doi:S0732-8893(09)00483-0 [pii] — 10.1016/j.diagmicrobio.2009.12.006 — 33. WILDEMAUWE C, DE BROUWER D, GODARD C, BUYSENS P, DEWIT J, JOSEPH R, VANHOOF R, Pathol Biol 58 (1) (2008) 70. doi:S0369-8114(09)00162-X [pii] 10.1016/j.patbio.2009.07.024 — 34. KRZIWANEK K, LUGER C, SAMMER B, STUMVOLL S, STAMMLER M, METZ-GERCEKS, MITTERMAYER H, Eur J Clin Microbiol Infect Dis 26 (12) (2007) 931. doi:10.1007/s10096-007-0391-4

## **PRVI IZOLAT *STAPHYLOCOCCUS AUREUS* REZISTENTNOG NA METICILIN *SCCMEC* TIPA V, POZITIVAN NA PVL**

### **SAŽETAK**

Cilj studije je bio istražiti epidemiološke i molekularne karakteristike MRSA izolata iz Primorsko-Goranske županije tijekom šestogodišnjeg perioda. U periodu od 2001. Do 2007. godine, 46 MRSA izolata je prikupljeno, testirana je osjetljivost na antimikrobne lijekove prema CLSI smjernicama, također je provedena detekcija *mecA* gena, *SCCmec* tipizacija i detekcija PVL kodirajućih gena. Tipizacija sojeva dodatno je upotpunjena analizom elektroforeze u pulsirajućem polju (PFGE analizom) i tipizacijom spa lokusa. Svi MRSA izolati su osjetljivi na vankomicin, linezolid, mupirocin, nitrofurantoin, samo jedan soj je bio rezistentan na fucidinsku kiselinu i ko-trimoksazol. Rezultati *SCCmec* tipizacije pokazali su prisutnost *SCCmec* tipa IV u 26 MRSA sojeva, *SCCmec* tipa V kod tri izolata, a 13 sojeva je imalo *SCCmec* I. *SCCmec* tipovi II i III nisu nađeni. Četiri MRSA su bila netipabilna navedenom metodom. PVL je detektiran kod 4 soja, dva *SCCmec* IV i dva *SCCmec* V. PFGE analiza grupirala je MRSA sojeve u 6 grupa sličnosti 18 pojedinačnih, nesvrstanih sojeva. Dominantni spa tipovi u ovoj kolekciji sojeva su bili: t015, s 15 sojeva, te t041 (N=7), t051, (N=2), t2850 (N=2), t008 (N=2) i s po jednim izolatom tipiziranim kao: t441, t002, t448, t018, t019, t355, t390, t026, t449, t148. Otkrili smo i nove tipove, t3510 i t3509. Ovo je prvi spomen o prisutnosti *SCCmec* tipa V u Hrvatskoj, i, prema našem saznanju, prvo izvješće od PVL-pozitivnom MRSA s *SCCmec* tipom V i spa tipom t441 (ST59-MRSA-V) u ovom dijelu Europe.